

PRIMITIVE FORMS FOR AFFINE CUSP POLYNOMIALS

YOSHIHISA ISHIBASHI, YUUKI SHIRAISHI, AND ATSUSHI TAKAHASHI

ABSTRACT. We determine a primitive form for a universal unfolding of an affine cusp polynomial. Moreover, we prove that the resulting Frobenius manifold is isomorphic to the one constructed from the Gromov-Witten theory for an orbifold projective line with at most three orbifold points.

1. INTRODUCTION

The notion of primitive forms is introduced by K. Saito [15, 16] in his study of period mapping associated to a deformation theory of an isolated hypersurface singularity. Roughly speaking, a primitive form is a generalization of a differential of the first kind on an elliptic curve and the tools in order to define a primitive form are natural generalization of a polarized Hodge structure. One of the most important ingredients is a Frobenius structure, which he called the *flat structure*, on the deformation space.

The existence of primitive forms is proved for isolated hypersurface singularities [18] (for comprehensive guides, see also [7, 14]). However, explicit forms of primitive forms are rarely known except for a few cases; for ADE singularities [15], for simple elliptic singularities [15, 20] and for some Laurent polynomials [3, 4]. In the present paper, we shall add a new class of polynomials to them.

Let A be a triplet (a_1, a_2, a_3) of positive integers such that $a_1 \leq a_2 \leq a_3$. Set $\mu_A = a_1 + a_2 + a_3 - 1$ and $\chi_A := 1/a_1 + 1/a_2 + 1/a_3 - 1$. We can associate to A the polynomial $f_A(\mathbf{x}) \in \mathbb{C}[x_1, x_2, x_3]$ given as

$$f_A(\mathbf{x}) := x_1^{a_1} + x_2^{a_2} + x_3^{a_3} - q^{-1} \cdot x_1 x_2 x_3 \quad (1.1)$$

for some $q \in \mathbb{C} \setminus \{0\}$, which we shall call a cusp polynomial of type A , in particular, an affine cusp polynomial of type A if $\chi_A > 0$.

The purpose of this paper is to determine a primitive form for a universal unfolding F_A of an affine cusp polynomial f_A , a deformation F_A of f_A over a μ_A -dimensional manifold $M := \mathbb{C}^{\mu_A-1} \times (\mathbb{C} \setminus \{0\})$ given as

$$F_A(\mathbf{x}; \mathbf{s}, s_{\mu_A}) := x_1^{a_1} + x_2^{a_2} + x_3^{a_3} - s_{\mu_A}^{-1} \cdot x_1 x_2 x_3 + s_1 \cdot 1 + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} s_{i,j} \cdot x_i^j \quad (1.2)$$

which induces an $\mathcal{O}_M$ -isomorphism

$$\mathcal{T}_M \longrightarrow \mathcal{O}_M[x_1, x_2, x_3] \left/ \left(\frac{\partial F_A}{\partial x_1}, \frac{\partial F_A}{\partial x_2}, \frac{\partial F_A}{\partial x_3} \right) \right., \quad \delta \mapsto \delta F_A. \quad (1.3)$$

Moreover, we shall prove that the resulting Frobenius manifold is isomorphic to the one constructed from the Gromov-Witten theory for an orbifold projective line with at most three orbifold points.

First, we shall show the following theorem, which is our main theorem in this paper:

Theorem (Theorem 3.1). *The element $\zeta_A := [s_{\mu_A}^{-1} dx_1 \wedge dx_2 \wedge dx_3] \in \mathcal{H}_{F_A}^{(0)}$ is a primitive form for the tuple $(\mathcal{H}_{F_A}^{(0)}, \nabla, K_{F_A})$ with the minimal exponent $r = 1$.*

There is an $\mathcal{O}_M$ -algebra structure on $\mathcal{T}_M$ by the $\mathcal{O}_M$ -isomorphism (1.3), which we shall denote by $\circ$. Note also that we have special elements $[1]$ and $[F_A]$ on the right hand side of (1.3) which give elements $e \in \Gamma(M, \mathcal{T}_M)$ which is the identity with respect to the product $\circ$ and $E \in \Gamma(M, \mathcal{T}_M)$. As a corollary of our main theorem, we obtain a non-degenerate symmetric $\mathcal{O}_M$ -bilinear form on $\mathcal{T}_M$ such that the tuple $(\eta, \circ, e, E)$ defines on the deformation space M a Frobenius structure of rank μ_A and dimension one (cf. [19]).

Next, we show that this Frobenius structure satisfies the conditions in Theorem 3.1 of [9].

Theorem (Theorem 4.1). *For the Frobenius structure of rank μ_A and dimension one constructed from the pair (f_A, ζ_A) , there exists flat coordinates $t_1, t_{1,1}, \dots, t_{3,a_3-1}, t_{\mu_A}$ satisfying the following conditions:*

(i) *The unit vector field e and the Euler vector field E are given by*

$$e = \frac{\partial}{\partial t_1}, \quad E = t_1 \frac{\partial}{\partial t_1} + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} \frac{a_i - j}{a_i} t_{i,j} \frac{\partial}{\partial t_{i,j}} + \chi_A \frac{\partial}{\partial t_{\mu_A}}.$$

(ii) *The non-degenerate symmetric bilinear form η on $\mathcal{T}_M$ defined by*

$$\eta(\delta, \delta') := K_{F_A}^{(0)}(u \nabla_{\delta} \zeta_A, u \nabla_{\delta'} \zeta), \quad \delta, \delta' \in \mathcal{T}_M, \quad (1.4)$$

satisfies

$$\begin{aligned} \eta \left(\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_{\mu_A}} \right) &= \eta \left(\frac{\partial}{\partial t_{\mu_A}}, \frac{\partial}{\partial t_1} \right) = 1, \\ \eta \left(\frac{\partial}{\partial t_{i_1, j_1}}, \frac{\partial}{\partial t_{i_2, j_2}} \right) &= \begin{cases} \frac{1}{a_{i_1}} & i_1 = i_2 \text{ and } j_2 = a_{i_1} - j_1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

(iii) *The Frobenius potential $\mathcal{F}_{f_A, \zeta_A}$ satisfies $E \mathcal{F}_{f_A, \zeta_A}|_{t_1=0} = 2 \mathcal{F}_{f_A, \zeta_A}|_{t_1=0}$,*

$$\mathcal{F}_{f_A, \zeta_A}|_{t_1=0} \in \mathbb{C} [t_{1,1}, \dots, t_{1,a_1-1}, t_{2,1}, \dots, t_{2,a_2-1}, t_{3,1}, \dots, t_{3,a_3-1}, e^{t_{\mu_A}}].$$

- (iv) Assume the condition (iii). The restriction of the Frobenius potential $\mathcal{F}_{f_A, \zeta_A}$ to the submanifold $\{t_1 = e^{t_{\mu_A}} = 0\}$ is given as

$$\mathcal{F}_{f_A, \zeta_A}|_{t_1=e^{t_{\mu_A}}=0} = \mathcal{G}^{(1)} + \mathcal{G}^{(2)} + \mathcal{G}^{(3)},$$

where $\mathcal{G}^{(i)} \in \mathbb{C}[t_{i,1}, \dots, t_{i,a_i-1}]$, $i = 1, 2, 3$.

- (v) Assume the condition (iii). In the frame $\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_{1,1}}, \dots, \frac{\partial}{\partial t_{3,a_3-1}}, \frac{\partial}{\partial t_{\mu_A}}$ of $\mathcal{T}_M$, the product $\circ$ can be extended to the limit $t_1 = t_{1,1} = \dots = t_{3,a_3-1} = e^{t_{\mu_A}} = 0$. The $\mathbb{C}$ -algebra obtained in this limit is isomorphic to

$$\mathbb{C}[x_1, x_2, x_3] / (x_1 x_2, x_2 x_3, x_3 x_1, a_1 x_1^{a_1} - a_2 x_2^{a_2}, a_2 x_2^{a_2} - a_3 x_3^{a_3}),$$

where $\partial/\partial t_{1,1}, \partial/\partial t_{2,1}, \partial/\partial t_{3,1}$ are mapped to x_1, x_2, x_3 , respectively.

- (vi) The term

$$\begin{cases} e^{t_{\mu_A}} & \text{if } a_1 = a_2 = a_3 = 1, \\ t_{3,1} e^{t_{\mu_A}} & \text{if } 1 = a_1 = a_2 < a_3, \\ t_{2,1} t_{3,1} e^{t_{\mu_A}} & \text{if } 1 = a_1 < a_2, \\ t_{1,1} t_{2,1} t_{3,1} e^{t_{\mu_A}} & \text{if } a_1 = 2, \end{cases}$$

occurs with the coefficient 1 in $\mathcal{F}_{f_A, \zeta_A}$.

As a consequence, we obtain the mirror isomorphism as Frobenius manifolds:

Corollary (Corollary 4.5). *There exists an isomorphism of Frobenius manifolds between the one constructed from the Gromov–Witten theory for $\mathbb{P}_A^1$ and the one constructed from the pair (f_A, ζ_A) .*

This result is already obtained by Milanov–Tseng [11] for the case $a_1 = 1$ and by Rossi [13] for the case $\chi_A > 0$. The present work is the second in a series of papers to simplify and generalise their works. In our subsequent papers, we shall show that the Frobenius structure constructed from the pair (f_A, ζ_A) in this paper is isomorphic to the one constructed from the extended affine Weyl group $\widehat{W}_A$ by Dubrovin–Zhang in [5] and that the Frobenius structure constructed from cusp singularity (i.e., f_A with $\chi_A < 0$) with the “canonical” primitive form satisfies the conditions in Theorem 3.1 of [9].

Acknowledgement

The second named author is deeply grateful to Claus Hertling and Christian Sevenheck for their valuable discussions and encouragement. He is supported by the JSPS International Training Program (ITP). The third named author is supported by Grant-in Aid for Scientific Research grant numbers 24684005 from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

2. NOTATIONS AND TERMINOLOGIES

Let A be a triplet (a_1, a_2, a_3) of positive integers such that $a_1 \leq a_2 \leq a_3$. Set

$$\mu_A := a_1 + a_2 + a_3 - 1 \quad (2.1)$$

and

$$\chi_A := \frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} - 1. \quad (2.2)$$

2.1. Universal unfolding of affine cusp polynomials.

Definition 2.1. A polynomial $f_A(\mathbf{x}) \in \mathbb{C}[x_1, x_2, x_3]$ given as

$$f_A(\mathbf{x}) := x_1^{a_1} + x_2^{a_2} + x_3^{a_3} - q^{-1} \cdot x_1 x_2 x_3 \quad (2.3)$$

for some $q \in \mathbb{C} \setminus \{0\}$ is called the *cusp polynomial of type A* . In particular, we shall call $f_A(\mathbf{x})$ an *affine cusp polynomial of type A* if $\chi_A > 0$.

Assumption 2.2. *Throughout this paper, we shall assume that the triplet of integers A satisfies $\chi_A > 0$. In particular, we shall only consider affine cusp polynomials.*

Let $f_A = f_A(\mathbf{x})$ be an affine cusp polynomial. One can easily show that

$$\mathbb{C}[x_1, x_2, x_3] \left/ \left(\frac{\partial f_A}{\partial x_1}, \frac{\partial f_A}{\partial x_2}, \frac{\partial f_A}{\partial x_3} \right) \right.$$

is of rank μ_A . We can consider the *universal unfolding* of f_A , a deformation F_A of f_A defined on $\mathbb{C}^3 \times M$, $M := \mathbb{C}^{\mu_A-1} \times (\mathbb{C} \setminus \{0\})$ over a μ_A -dimensional parameters $(\mathbf{s}, s_{\mu_A}) \in M$ given as follows:

Definition 2.3. Define a function $F_A(\mathbf{x}; \mathbf{s}, s_{\mu_A})$ defined on $\mathbb{C}^3 \times M$ as follows;

$$F_A(\mathbf{x}; \mathbf{s}, s_{\mu_A}) := x_1^{a_1} + x_2^{a_2} + x_3^{a_3} - s_{\mu_A}^{-1} \cdot x_1 x_2 x_3 + s_1 \cdot 1 + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} s_{i,j} \cdot x_i^j. \quad (2.4)$$

Denote by

$$p : \mathbb{C}^3 \times M \longrightarrow M, \quad (\mathbf{x}; \mathbf{s}, s_{\mu_A}) \mapsto (\mathbf{s}, s_{\mu_A})$$

the projection map from the total space to the deformation space. Set

$$p_* \mathcal{O}_{\mathcal{C}} := \mathcal{O}_M[x_1, x_2, x_3] \left/ \left(\frac{\partial F_A}{\partial x_1}, \frac{\partial F_A}{\partial x_2}, \frac{\partial F_A}{\partial x_3} \right) \right. \quad (2.5)$$

$p_* \mathcal{O}_{\mathcal{C}}$ can be thought of as the direct image of the sheaf of relative *algebraic* functions on the relative critical set $\mathcal{C}$ of F_A with respect to the projection $p : \mathbb{C}^3 \times M \longrightarrow M$.

Proposition 2.4. *The function $F_A(\mathbf{x}; \mathbf{s}, s_{\mu_A})$ satisfies the following conditions:*

- (i) $F_A(\mathbf{x}; \mathbf{0}, q) = f_A(\mathbf{x})$.

(ii) The $\mathcal{O}_M$ -homomorphism ρ called the Kodaira–Spencer map defined as

$$\rho : \mathcal{T}_M \longrightarrow p_*\mathcal{O}_C, \quad \delta \mapsto \delta F_A, \quad (2.6)$$

is an isomorphism.

Proof. Since F_A is tame at any point on the deformation space M , namely, there are no critical points coming from infinity. Therefore, the statement follows from the same argument in Section 2.a of [3]. $\square$

Note that the tangent bundle $\mathcal{T}_M$ naturally obtains an $\mathcal{O}_M$ -algebra structure.

Definition 2.5. We shall denote by $\circ$ the induced product structure on $\mathcal{T}_M$ by the $\mathcal{O}_M$ -isomorphism (2.6). Namely, for $\delta, \delta' \in \mathcal{T}_M$, we have

$$(\delta \circ \delta')F_A = \delta F_A \cdot \delta' F_A \text{ in } p_*\mathcal{O}_C. \quad (2.7)$$

Set $\overline{M} := \mathbb{C}^{\mu_A-1} \times \mathbb{C}$. Let ι be the natural embedding

$$\iota : M \hookrightarrow \overline{M}, \quad (\mathbf{s}, s_{\mu_A}) \mapsto (\mathbf{s}, s_{\mu_A}), \quad (2.8)$$

and let M_∞ be the divisor $\overline{M} \setminus \iota(M) := \{(\mathbf{s}, s_{\mu_A}) \in \overline{M} \mid s_{\mu_A} = 0\}$. Denote again by p the projection map from the total space to the deformation space:

$$p : \mathbb{C}^3 \times \overline{M} \longrightarrow \overline{M}, \quad (\mathbf{x}; \mathbf{s}, s_{\mu_A}) \mapsto (\mathbf{s}, s_{\mu_A})$$

Set

$$p_*\mathcal{O}_{\overline{C}} := \mathcal{O}_{\overline{M}}[x_1, x_2, x_3] \left/ \left(s_{\mu_A} \frac{\partial F_A}{\partial x_1}, s_{\mu_A} \frac{\partial F_A}{\partial x_2}, s_{\mu_A} \frac{\partial F_A}{\partial x_3}, H_1(\mathbf{x}, \mathbf{s}), H_2(\mathbf{x}, \mathbf{s}) \right) \right. . \quad (2.9)$$

where

$$H_i(\mathbf{x}, \mathbf{s}) := a_i x_i^{a_i} - a_{i+1} x_{i+1}^{a_{i+1}} + \sum_{j=1}^{a_i-1} j \cdot s_{i,j} \cdot x_i^j - \sum_{j=1}^{a_{i+1}-1} j \cdot s_{i+1,j} \cdot x_{i+1}^j, \quad i = 1, 2. \quad (2.10)$$

Proposition 2.6. As an $\mathcal{O}_{\overline{M}}$ -module, $p_*\mathcal{O}_{\overline{C}}$ is free of rank μ_A . As an $\mathcal{O}_{\overline{M}}$ -algebra, it is isomorphic to the $\mathcal{O}_{\overline{M}}$ -subalgebra of $\iota_*(p_*\mathcal{O}_C)$ whose $\mathcal{O}_{\overline{M}}$ -basis is given by the set of residue classes of monomials

$$\{1, x_i^j \ (i = 1, 2, 3, \ j = 1, \dots, a_i - 1), s_{\mu_A}^{-1} x_1 x_2 x_3\}. \quad (2.11)$$

In particular, we have the isomorphism of $\mathcal{O}_M$ -modules

$$\iota^*(p_*\mathcal{O}_{\overline{C}}) \simeq p_*\mathcal{O}_C. \quad (2.12)$$

Namely, $p_*\mathcal{O}_{\overline{C}}$ is an extension of $p_*\mathcal{O}_C$ as a free $\mathcal{O}_{\overline{M}}$ -module.

Proof. It is almost obvious that $\iota^*(p_*\mathcal{O}_{\overline{C}}) \simeq p_*\mathcal{O}_C$.

Lemma 2.7. *We have*

$$p_*\mathcal{O}_{\overline{\mathcal{C}}}/s_{\mu_A}p_*\mathcal{O}_{\overline{\mathcal{C}}} \simeq \mathcal{O}_{M_\infty}[x_1, x_2, x_3] / \left(x_2x_3, x_3x_1, x_1x_2, H_1(\mathbf{x}, \mathbf{s}), H_2(\mathbf{x}, \mathbf{s}) \right), \quad (2.13)$$

is a free $\mathcal{O}_{M_\infty}$ -module of rank μ_A . In particular, we have

$$p_*\mathcal{O}_{\overline{\mathcal{C}}}/\mathfrak{m}_{(0,0)}p_*\mathcal{O}_{\overline{\mathcal{C}}} \simeq \mathbb{C}[x_1, x_2, x_3] / \left(x_2x_3, x_3x_1, x_1x_2, a_1x_1^{a_1} - a_2x_2^{a_2}, a_2x_2^{a_2} - a_3x_3^{a_3} \right), \quad (2.14)$$

where $\mathfrak{m}_{(0,0)}$ is the maximal ideal of $\mathcal{O}_{\overline{M}}$ corresponding to the point $(0,0)$.

Proof. Some elementary calculations yield the statement. $\square$

By this lemma, we see that $p_*\mathcal{O}_{\overline{\mathcal{C}}}$ is free of rank μ_A as an $\mathcal{O}_{\overline{M}}$ -module. The rest follows from the following equality in $\iota_*(p_*\mathcal{O}_{\mathcal{C}})$:

$$[s_{\mu_A}^{-1}x_1x_2x_3] = \left[a_i x_i^{a_i} + \sum_{j=1}^{a_i-1} j \cdot s_{i,j} \cdot x_i^j \right], \quad i = 1, 2, 3. \quad (2.15)$$

$\square$

Therefore, $p_*\mathcal{O}_{\overline{\mathcal{C}}}$ can be thought of as the direct image of the sheaf of relative algebraic functions on $\overline{\mathcal{C}}$, a closure of the relative critical set $\mathcal{C}$ in $\mathbb{C}^3 \times \overline{M}$, which is flat and finite over $\overline{M}$.

Proposition 2.8. *Denote by $\mathcal{T}_{\overline{M}}(-\log M_\infty)$ the sheaf of holomorphic vector fields on $\overline{M}$ with logarithmic zeros along the divisor M_∞ defined as*

$$\mathcal{T}_{\overline{M}}(-\log M_\infty) := \{ \delta \in \mathcal{T}_{\overline{M}} \mid \delta s_{\mu_A} \in s_{\mu_A} \cdot \mathcal{O}_{\overline{M}} \}. \quad (2.16)$$

Then, $\mathcal{T}_{\overline{M}}(-\log M_\infty)$ is a free $\mathcal{O}_{\overline{M}}$ -module of rank μ_A and we have the isomorphism of $\mathcal{O}_M$ -modules

$$\iota^*(\mathcal{T}_{\overline{M}}(-\log M_\infty)) \simeq \mathcal{T}_M. \quad (2.17)$$

Namely, $\mathcal{T}_{\overline{M}}(-\log M_\infty)$ is an extension of $\mathcal{T}_M$ as a free $\mathcal{O}_{\overline{M}}$ -module.

Proof. The statement follows from the fact that

$$\mathcal{T}_{\overline{M}}(-\log M_\infty) \simeq \mathcal{O}_{\overline{M}} \frac{\partial}{\partial s_1} \bigoplus_{\substack{1 \leq i \leq 3, \\ 1 \leq j \leq a_i-1}} \mathcal{O}_{\overline{M}} \frac{\partial}{\partial s_{i,j}} \bigoplus \mathcal{O}_{\overline{M}} s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}. \quad (2.18)$$

$\square$

Proposition 2.9. *The Kodaira–Spencer map ρ induces the $\mathcal{O}_{\overline{M}}$ -isomorphism $\overline{\rho}$*

$$\overline{\rho} : \mathcal{T}_{\overline{M}}(-\log M_\infty) \longrightarrow p_*\mathcal{O}_{\overline{\mathcal{C}}}. \quad (2.19)$$

Proof. The statement easily follows since the Kodaira–Spencer map ρ induces the $\mathcal{O}_{\overline{M}}$ -isomorphism between $\mathcal{T}_{\overline{M}}(-\log M_\infty)$ and the free $\mathcal{O}_{\overline{M}}$ -submodule of $\iota_*(p_*\mathcal{O}_C)$ spanned by the residue classes of monomials

$$1, x_i^j \ (i = 1, 2, 3, \ j = 1, \dots, a_i - 1), \ s_{\mu_A}^{-1}x_1x_2x_3. \quad (2.20)$$

□

2.2. Primitive vector field and Euler vector field.

Definition 2.10. The vector field e and E on M corresponding to the unit 1 and F by the $\mathcal{O}_M$ -isomorphism (2.6) is called the *primitive vector field* and the *Euler vector field*, respectively. That is,

$$eF_A = 1 \text{ and } EF_A = F_A \text{ in } p_*\mathcal{O}_C. \quad (2.21)$$

Proposition 2.11. *The primitive vector field e and the Euler vector field E on M are given by*

$$e = \frac{\partial}{\partial s_1}, \quad E = s_1 \frac{\partial}{\partial s_1} + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} \frac{a_i - j}{a_i} s_{i,j} \frac{\partial}{\partial s_{i,j}} + \chi_A s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}. \quad (2.22)$$

In particular, they naturally define elements of $\Gamma(\overline{M}, \mathcal{T}_{\overline{M}}(-\log M_\infty))$.

Proof. This is clear. □

Lemma 2.12. *We have the “Euler’s identity”:*

$$F_A = EF_A + \sum_{i=1}^3 \frac{1}{a_i} x_i \frac{\partial F_A}{\partial x_i}. \quad (2.23)$$

Proof. This is also clear. □

Definition 2.13. An element $g \in \mathcal{O}_M$ is of *degree l* for some $k \in \mathbb{Q}$ if it satisfies the equation $Eg = lg$ where E is the Euler vector field. If $Eg = lg$, then l is denoted by $\deg(g)$.

Lemma 2.14. *Suppose that $g \in \mathcal{O}_{\overline{M}}$ satisfies $Eg = \deg(g) \cdot g$. Then, we have $\deg(g) \geq 0$ and $\deg(g) = 0$ if and only if g is a constant function.*

Proof. It follows from our assumption that $\chi_A > 0$. □

2.3. Filtered de Rham cohomology. For any non-negative integer i , we set

$$p_*\Omega_{\mathbb{A}^3 \times M/M}^i := \Gamma(\mathbb{A}^3, \Omega_{\mathbb{A}^3}^i) \otimes_{\mathbb{C}} \mathcal{O}_M, \quad (2.24)$$

where $\Omega_{\mathbb{A}^3}^i$ denotes the sheaf of *algebraic* differential i -forms on $\mathbb{A}^3$, the algebraic variety underlying the complex manifold $\mathbb{C}^3$. $p_*\Omega_{\mathbb{A}^3 \times M/M}^i$ can be thought of as the direct image of the sheaf of relative algebraic differential i -forms with respect to the projection $p : \mathbb{C}^3 \times M \rightarrow M$.

Definition 2.15. Define Ω_{F_A} as

$$\Omega_{F_A} := p_*\Omega_{\mathbb{A}^3 \times M/M}^3 / dF_A \wedge p_*\Omega_{\mathbb{A}^3 \times M/M}^2. \quad (2.25)$$

Proposition 2.16. Ω_{F_A} is a free $p_*\mathcal{O}_C$ -module of rank one and hence a free $\mathcal{O}_M$ -module of rank μ_A .

Proof. A $p_*\mathcal{O}_C$ -free base of Ω_{F_A} can be chosen as $[\omega]$, $\omega := dx_1 \wedge dx_2 \wedge dx_3$, and hence an $\mathcal{O}_M$ -free basis of Ω_{F_A} can be chosen as

$$\{[\omega], [x_i^j \omega] \ (i = 1, 2, 3, \ j = 1, \dots, a_i - 1), [x_1 x_2 x_3 \omega]\}. \quad (2.26)$$

□

Definition 2.17. We set

$$\mathcal{H}_{F_A} := \mathbb{R}^3 p_*(\Omega_{\mathbb{A}^3 \times M/M}^\bullet \otimes_{\mathcal{O}_M} \mathcal{O}_M((u)), ud + dF_A \wedge) \quad (2.27)$$

and call it the *filtered de Rham cohomology group* of the universal unfolding F_A .

For any $k \in \mathbb{Z}$, put

$$\mathcal{H}_{F_A}^{(-k)} := \mathbb{R}^3 p_*(\Omega_{\mathbb{A}^3 \times M/M}^\bullet \otimes_{\mathcal{O}_M} \mathcal{O}_M[[u]]u^k, ud + dF_A \wedge).$$

Obviously, one has an $\mathcal{O}_M[[u]]$ -isomorphism for all $i \in \mathbb{Z}$

$$\mathcal{H}_{F_A}^{(0)} \simeq \mathcal{H}_{F_A}^{(-k)}, \quad \omega \mapsto u^k \omega, \quad (2.28)$$

and an $\mathcal{O}_M((u))$ -isomorphism

$$\mathcal{H}_{F_A}^{(-k)} \otimes_{\mathcal{O}_M[[u]]} \mathcal{O}_M((u)) \simeq \mathcal{H}_{F_A}. \quad (2.29)$$

Furthermore, $\mathcal{H}_{F_A}^{(-k)}$ is naturally a submodule of $\mathcal{H}_{F_A}$ so that $\{\mathcal{H}_{F_A}^{(-k)}\}_{i \in \mathbb{Z}}$ form an increasing and exhaustive filtration of $\mathcal{H}_{F_A}$:

$$\dots \subset \mathcal{H}_{F_A}^{(-k-1)} \subset \mathcal{H}_{F_A}^{(-k)} \subset \dots \subset \mathcal{H}_{F_A}^{(-1)} \subset \mathcal{H}_{F_A}^{(0)} \subset \dots \subset \mathcal{H}_{F_A}, \quad (2.30)$$

such that $\mathcal{H}_{F_A}$ is complete with respect to the filtration in the following sense:

$$\bigcup_{k \in \mathbb{Z}} \mathcal{H}_{F_A}^{(-k)} = \mathcal{H}_{F_A} \text{ and } \bigcap_{k \in \mathbb{Z}} \mathcal{H}_{F_A}^{(-k)} = \{0\}. \quad (2.31)$$

Proposition 2.18. *For any $k \in \mathbb{Z}$, $\mathcal{H}_{F_A}^{(-k)}$ is an $\mathcal{O}_M[[u]]$ -free module of rank μ_A . In particular, we have the following short exact sequence of $\mathcal{O}_M$ -modules*

$$0 \rightarrow \mathcal{H}_{F_A}^{(-1)} \hookrightarrow \mathcal{H}_{F_A}^{(0)} \xrightarrow{r^{(0)}} \Omega_{F_A} \rightarrow 0. \quad (2.32)$$

Proof. One can choose an $\mathcal{O}_M[[u]]$ -free basis of $\mathcal{H}_{F_A}^{(0)}$ as

$$\{[\omega], [x_i^j \omega] \ (i = 1, 2, 3, \ j = 1, \dots, a_i - 1), [x_1 x_2 x_3 \omega]\} \quad (2.33)$$

where $\omega = dx_1 \wedge dx_2 \wedge dx_3$. The rest is clear. $\square$

Definition 2.19. Define an element ζ_A of $\Gamma(M, \mathcal{H}_{F_A}^{(0)})$ as

$$\zeta_A := [s_{\mu_A}^{-1} dx_1 \wedge dx_2 \wedge dx_3]. \quad (2.34)$$

Definition 2.20. For any $k \in \mathbb{Z}$, define an $\mathcal{O}_{\overline{M}}[[u]]$ -free module $\overline{\mathcal{H}}_{F_A}^{(-k)}$ of rank μ_A as

$$\overline{\mathcal{H}}_{F_A}^{(k)} := \mathcal{O}_{\overline{M}}[[u]] \cdot u^k \zeta_A \bigoplus_{\substack{1 \leq i \leq 3, \\ 1 \leq j \leq a_i - 1}} \mathcal{O}_{\overline{M}}[[u]] \cdot u^k x_i^j \zeta_A \bigoplus \mathcal{O}_{\overline{M}}[[u]] \cdot u^k s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A, \quad (2.35)$$

where we regard $\zeta_A, x_i^j \zeta_A, s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A$ as elements of $\Gamma(\overline{M}, \iota_* \mathcal{H}_{F_A}^{(0)})$.

Proposition 2.21. *We have the isomorphism of $\mathcal{O}_M[[u]]$ -modules*

$$\iota^* \left(\overline{\mathcal{H}}_{F_A}^{(-k)} \right) \simeq \mathcal{H}_{F_A}^{(-k)}, \quad i \in \mathbb{Z}. \quad (2.36)$$

Namely, $\overline{\mathcal{H}}_{F_A}^{(-k)}$ is an extension of $\mathcal{H}_{F_A}^{(-k)}$ as a free $\mathcal{O}_{\overline{M}}[[u]]$ -module.

Proof. It is clear by Proposition 2.18. $\square$

Definition 2.22. Define an $\mathcal{O}_{\overline{M}}$ -free module $\overline{\Omega}_{F_A}$ of rank μ_A as

$$\overline{\Omega}_{F_A} := \mathcal{O}_{\overline{M}} \cdot r^{(0)}(\zeta_A) \bigoplus_{\substack{1 \leq i \leq 3, \\ 1 \leq j \leq a_i - 1}} \mathcal{O}_{\overline{M}} \cdot r^{(0)}(x_i^j \zeta_A) \bigoplus \mathcal{O}_{\overline{M}} \cdot r^{(0)}(s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A), \quad (2.37)$$

where we regard $r^{(0)}(\zeta_A), r^{(0)}(x_i^j \zeta_A), r^{(0)}(s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A)$ as elements of $\Gamma(\overline{M}, \iota_* \Omega_{F_A})$.

Proposition 2.23. *We have the isomorphism of $\mathcal{O}_M$ -modules*

$$\iota^* \left(\overline{\Omega}_{F_A} \right) \simeq \Omega_{F_A}. \quad (2.38)$$

Namely, $\overline{\Omega}_{F_A}$ is an extension of Ω_{F_A} as a free $\mathcal{O}_{\overline{M}}$ -module.

Proof. It is almost clear. $\square$

Proposition 2.24. *$\overline{\Omega}_{F_A}$ is a free $p_* \mathcal{O}_{\overline{C}}$ -module of rank one.*

Proof. A $p_* \mathcal{O}_{\overline{C}}$ -free base of $\overline{\Omega}_{F_A}$ can be chosen as $r^{(0)}(\zeta_A)$ in Definition 2.22. $\square$

Proposition 2.25. *We have the following short exact sequence of $\mathcal{O}_{\overline{M}}$ -modules*

$$0 \rightarrow \overline{\mathcal{H}}_{F_A}^{(-1)} \hookrightarrow \overline{\mathcal{H}}_{F_A}^{(0)} \xrightarrow{\overline{r}^{(0)}} \overline{\Omega}_{F_A} \rightarrow 0. \quad (2.39)$$

Proof. It is clear by their definitions. $\square$

2.4. Gauß–Manin connection. We define the free $\mathcal{O}_M[[u]]$ -module $\mathcal{T}_{\mathbb{C}_u \times M}$ of rank $\mu_A + 1$ as follows:

$$\mathcal{T}_{\mathbb{C}_u \times M} := \mathcal{O}_M[[u]] \frac{d}{du} \oplus \mathcal{O}_M[[u]] \otimes_{\mathcal{O}_M} \mathcal{T}_M. \quad (2.40)$$

Definition 2.26. We define a connection, called the *Gauß–Manin connection*,

$$\nabla : \mathcal{T}_{\mathbb{C}_u \times M} \otimes_{\mathcal{O}_M} \mathcal{H}_{F_A} \rightarrow \mathcal{H}_{F_A} \quad (2.41)$$

by letting; for $\delta \in \mathcal{T}_M$ and $\zeta = [\phi dx_1 \wedge \cdots \wedge dx_n] \in \mathcal{H}_{F_A}$,

$$\nabla_{\delta} \zeta := \left[\left(\frac{1}{u} (\delta F_A) \phi + \delta(\phi) \right) dx_1 \wedge \cdots \wedge dx_n \right], \quad (2.42a)$$

$$\nabla_{\frac{d}{du}} \zeta := \left[\left(-\frac{1}{u^2} F_A \phi + \frac{d\phi}{du} \right) dx_1 \wedge \cdots \wedge dx_n \right]. \quad (2.42b)$$

Proposition 2.27. *Gauß–Manin connection $\nabla : \mathcal{T}_{\mathbb{C}_u \times M} \otimes_{\mathcal{O}_M} \mathcal{H}_{F_A} \longrightarrow \mathcal{H}_{F_A}$ satisfies following;*

(i) ∇ is integrable:

$$\left[\nabla_{\frac{d}{du}}, \nabla_{\frac{d}{du}} \right] = 0, \quad \left[\nabla_{\frac{d}{du}}, \nabla_{\delta} \right] = 0, \quad [\nabla_{\delta}, \nabla_{\delta'}] = \nabla_{[\delta, \delta']}, \quad \delta, \delta' \in \mathcal{T}_M.$$

(ii) ∇ satisfies Griffith transversality: that is,

$$\nabla : \mathcal{T}_M \otimes_{\mathcal{O}_M} \mathcal{H}_{F_A}^{(-k)} \longrightarrow \mathcal{H}_{F_A}^{(-k+1)}, \quad k \in \mathbb{Z}. \quad (2.43)$$

(iii) The covariant differentiation $\nabla_{\frac{d}{du}}$ satisfies

$$\nabla_{u \frac{d}{du}} (\mathcal{H}_{F_A}^{(-k)}) \subset \mathcal{H}_{F_A}^{(-k+1)}, \quad k \in \mathbb{Z}. \quad (2.44)$$

Proof. See Proposition 4.5 of [19] and reference there in. $\square$

We shall consider the extension the Gauß–Manin connection ∇ on $\mathcal{H}_{F_A}$ to the one on $\overline{\mathcal{H}}_{F_A}$. Define the free $\mathcal{O}_{\overline{M}}[[u]]$ -module $\mathcal{T}_{\mathbb{C}_u \times \overline{M}}(-\log M_{\infty})$ of rank $\mu_A + 1$ as follows:

$$\mathcal{T}_{\mathbb{C}_u \times \overline{M}}(-\log M_{\infty}) := \mathcal{O}_{\overline{M}}[[u]] \frac{d}{du} \oplus \mathcal{O}_{\overline{M}}[[u]] \otimes_{\mathcal{O}_{\overline{M}}} \mathcal{T}_{\overline{M}}(-\log M_{\infty}). \quad (2.45)$$

Proposition 2.28. *The Gauß–Manin connection ∇ on $\mathcal{H}_{F_A}$ extends to $\overline{\mathcal{H}}_{F_A}$ with logarithmic poles along M_{∞} , namely, ∇ induces the connection*

$$\overline{\nabla} : \mathcal{T}_{\mathbb{C}_u \times \overline{M}}(-\log M_{\infty}) \otimes_{\mathcal{O}_{\overline{M}}} \overline{\mathcal{H}}_{F_A} \longrightarrow \overline{\mathcal{H}}_{F_A}, \quad (2.46)$$

satisfying the following conditions:

(i) $\overline{\nabla}$ is integrable:

$$\left[\overline{\nabla}_{\frac{d}{du}}, \overline{\nabla}_{\frac{d}{du}}\right] = 0, \quad \left[\overline{\nabla}_{\frac{d}{du}}, \overline{\nabla}_{\delta}\right] = 0, \quad [\overline{\nabla}_{\delta}, \overline{\nabla}_{\delta'}] = \overline{\nabla}_{[\delta, \delta']}, \quad \delta, \delta' \in \mathcal{T}_{\overline{M}}(-\log M_{\infty}).$$

(ii) $\overline{\nabla}$ satisfies Griffith transversality: that is,

$$\overline{\nabla} : \mathcal{T}_{\overline{M}}(-\log M_{\infty}) \otimes_{\mathcal{O}_{\overline{M}}} \overline{\mathcal{H}}_{F_A}^{(-k)} \longrightarrow \overline{\mathcal{H}}_{F_A}^{(-k+1)}, \quad k \in \mathbb{Z}. \quad (2.47)$$

(iii) The covariant differentiation $\overline{\nabla}_{\frac{d}{du}}$ satisfies

$$\overline{\nabla}_{u \frac{d}{du}}(\overline{\mathcal{H}}_{F_A}^{(-k)}) \subset \overline{\mathcal{H}}_{F_A}^{(-k+1)}, \quad k \in \mathbb{Z}. \quad (2.48)$$

Proof. We shall check that

$$s_{\mu_A} \nabla_{\frac{\partial}{\partial s_{\mu_A}}}(\overline{\mathcal{H}}_{F_A}) \subset \overline{\mathcal{H}}_{F_A}. \quad (2.49)$$

$$\nabla_{\frac{\partial}{\partial s_1}}(\overline{\mathcal{H}}_{F_A}) \subset \overline{\mathcal{H}}_{F_A}, \quad \nabla_{\frac{\partial}{\partial s_{i,j}}}(\overline{\mathcal{H}}_{F_A}) \subset \overline{\mathcal{H}}_{F_A}, \quad i = 1, 2, 3, \quad j = 1, \dots, a_i - 1. \quad (2.50)$$

$$u \nabla_{\frac{d}{du}}(\overline{\mathcal{H}}_{F_A}) \subset \overline{\mathcal{H}}_{F_A}. \quad (2.51)$$

First we shall check the condition (2.49). One has

$$s_{\mu_A} \nabla_{\frac{\partial}{\partial s_{\mu_A}}} \zeta_A = \frac{1}{u} s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A - \zeta_A, \quad (2.52)$$

$$s_{\mu_A} \nabla_{\frac{\partial}{\partial s_{\mu_A}}} x_i^j \zeta_A = \frac{1}{u} s_{\mu_A}^{-1} x_1 x_2 x_3 \cdot x_i^j \zeta_A - x_i^j \zeta_A, \quad (2.53)$$

$$s_{\mu_A} \nabla_{\frac{\partial}{\partial s_{\mu_A}}} s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A = \frac{1}{u} (s_{\mu_A}^{-1} x_1 x_2 x_3)^2 \zeta_A - 2 s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A. \quad (2.54)$$

The images of $s_{\mu_A}^{-1} x_1 x_2 x_3 \cdot x_i^j \zeta_A$ and $(s_{\mu_A}^{-1} x_1 x_2 x_3)^2 \zeta_A$ by $r^{(0)}$ can be extended to $\overline{\Omega}_{F_A}$. By Proposition 2.24 and Proposition 2.25, we can show that $1/u \cdot s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A$, $1/u \cdot (s_{\mu_A}^{-1} x_1 x_2 x_3)^2 \zeta_A \in \overline{\mathcal{H}}_{F_A}$.

Next we shall check that the condition (2.51). The assertion for s_1 is obvious. One has

$$\nabla_{\frac{\partial}{\partial s_{i',j'}}} \phi \cdot \zeta_A = \frac{1}{u} x_{i'}^{j'} \cdot \phi \cdot \zeta_A, \quad (2.55)$$

where ϕ is an element of the set $\{1, \partial F_A / \partial s_{i,j} \mid (i = 1, 2, 3, j = 1, \dots, a_i - 1), \partial F_A / \partial s_{\mu_A}\}$. The image of $x_{i'}^{j'} \cdot \phi \cdot \zeta_A$ by $r^{(0)}$ can be extended to $\overline{\Omega}_{F_A}$. By Proposition 2.24 and Proposition 2.25, we can show that $1/u \cdot x_{i'}^{j'} \cdot \phi \cdot \zeta_A \in \overline{\mathcal{H}}_{F_A}$.

Finally we shall check the condition (2.51). One has

$$u \nabla_{\frac{d}{du}} \phi \cdot \zeta_A = -\frac{1}{u} F_A \cdot \phi \cdot \zeta_A, \quad (2.56)$$

where ϕ is an element of the set $\{1, \partial F_A / \partial s_{i,j} \mid (i = 1, 2, 3, j = 1, \dots, a_i - 1), \partial F_A / \partial s_{\mu_A}\}$. The image of $F_A \cdot \phi \cdot \zeta_A$ by $r^{(0)}$ can be extended to $\overline{\Omega}_{F_A}$ since we have

$$F_A = E F_A + \sum_{i=1}^3 \frac{1}{a_i} x_i \frac{\partial F_A}{\partial x_i}. \quad (2.57)$$

By Proposition 2.24 and Proposition 2.25, we can show that $-1/u \cdot F_A \cdot \phi \cdot \zeta_A \in \overline{\mathcal{H}}_{F_A}$. The conditions (i), (ii) and (iii) follow from Proposition 2.27. $\square$

2.5. Higher residue pairing.

Definition 2.29. Define an $\mathcal{O}_M$ -bilinear form J_A on Ω_{F_A} by

$$J_{F_A}(\omega_1, \omega_2) := -\text{Res}_{\mathbb{C}^3 \times M/M} \left[\begin{array}{c} \phi_1 \phi_2 dx_1 \wedge dx_2 \wedge dx_3 \\ \frac{\partial F_A}{\partial x_1} \frac{\partial F_A}{\partial x_2} \frac{\partial F_A}{\partial x_3} \end{array} \right] \quad (2.58)$$

$$= \frac{-1}{(2\pi\sqrt{-1})^3} \int_C \frac{\phi_1 \phi_2 dx_1 \wedge dx_2 \wedge dx_3}{\frac{\partial F_A}{\partial x_1} \frac{\partial F_A}{\partial x_2} \frac{\partial F_A}{\partial x_3}} \quad (2.59)$$

where $\omega_1 = [\phi_1 dx_1 \wedge dx_2 \wedge dx_3]$ and $\omega_2 = [\phi_2 dx_1 \wedge dx_2 \wedge dx_3]$, C is the distinguished boundary of a polydisc containing all the critical points of F_A (see (3.3.7) and (3.3.8) of [14]).

Proposition 2.30. *The $\mathcal{O}_M$ -bilinear form J_{F_A} on Ω_{F_A} is non-degenerate.*

Proof. This is a well-known fact. In our case, we can show this by a direct calculation:

Lemma 2.31. *We have*

$$\text{Res}_{\mathbb{C}^3 \times M/M} \left[\begin{array}{c} 1 \cdot dx_1 \wedge dx_2 \wedge dx_3 \\ \frac{\partial F_A}{\partial x_1} \frac{\partial F_A}{\partial x_2} \frac{\partial F_A}{\partial x_3} \end{array} \right] = 0, \quad (2.60a)$$

$$\text{Res}_{\mathbb{C}^3 \times M/M} \left[\begin{array}{c} x_i^j \cdot dx_1 \wedge dx_2 \wedge dx_3 \\ \frac{\partial F_A}{\partial x_1} \frac{\partial F_A}{\partial x_2} \frac{\partial F_A}{\partial x_3} \end{array} \right] = 0, \quad i = 1, 2, 3, \quad j = 1, \dots, a_i - 1, \quad (2.60b)$$

and

$$\text{Res}_{\mathbb{C}^3 \times M/M} \left[\begin{array}{c} x_1 x_2 x_3 \cdot dx_1 \wedge dx_2 \wedge dx_3 \\ \frac{\partial F_A}{\partial x_1} \frac{\partial F_A}{\partial x_2} \frac{\partial F_A}{\partial x_3} \end{array} \right] = s_{\mu_A}^3. \quad (2.60c)$$

Proof. Some elementary calculations of residues yield the statement. $\square$

Now, the statement of Proposition 2.30 easily follows. $\square$

Proposition 2.32. *The pairing J_{F_A} on Ω_{F_A} induces a non-degenerate symmetric $\mathcal{O}_{\overline{M}}$ -bilinear form $\overline{J}_{F_A}$ on $\overline{\Omega}_{F_A}$ whose restriction to $\iota^*(\overline{\Omega}_{F_A}) \simeq \Omega_{F_A}$ coincides with J_{F_A} .*

Proof. Since the elements $r^{(0)}(\zeta_A), r^{(0)}(x_i^j \zeta_A), r^{(0)}(s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A)$ form an $\mathcal{O}_{\overline{M}}$ -basis of $\overline{\Omega}_{F_A}$, it is clear from Lemma 2.31. $\square$

In order to define the higher residue pairing, we prepare a notation. For $P = \sum_{i \in \mathbb{Z}} p_i u^i \in \mathcal{O}_M((u))$, set $P^* := \sum_{i \in \mathbb{Z}} p_i (-u)^i$ such that $(P^*)^* = P$.

Definition 2.33. An $\mathcal{O}_M$ -bilinear form

$$K_{F_A} : \mathcal{H}_{F_A} \otimes_{\mathcal{O}_M} \mathcal{H}_{F_A} \rightarrow \mathcal{O}_M((u)) \quad (2.61)$$

is called the *higher residue pairing* if it satisfies the following properties:

(i) For all $\omega_1, \omega_2 \in \mathcal{H}_{F_A}$,

$$K_{F_A}(\omega_1, \omega_2) = (-1)^3 K_{F_A}(\omega_2, \omega_1)^*. \quad (K1)$$

(ii) For all $P \in \mathcal{O}_M((u))$ and $\omega_1, \omega_2 \in \mathcal{H}_F$,

$$PK_{F_A}(\omega_1, \omega_2) = K_{F_A}(P\omega_1, \omega_2) = K_{F_A}(\omega_1, P^*\omega_2). \quad (K2)$$

(iii) For all $\omega_1, \omega_2 \in \mathcal{H}_{F_A}^{(0)}$,

$$K_{F_A}(\omega_1, \omega_2) \in u^3 \mathcal{O}_M[[u]]. \quad (K3)$$

(iv) The following diagram is commutative:

$$\begin{array}{ccc} K_{F_A} : \mathcal{H}_{F_A}^{(0)} \times \mathcal{H}_{F_A}^{(0)} & \longrightarrow & u^3 \mathcal{O}_M[[u]] \\ \downarrow & & \downarrow \text{mod } u^4 \mathcal{O}_M[[u]] \\ J_{F_A} : \Omega_{F_A} \times \Omega_{F_A} & \longrightarrow & u^3 \mathcal{O}_M. \end{array}$$

(v) For all $\omega_1, \omega_2 \in \mathcal{H}_F$ and $\delta \in \mathcal{T}_M$,

$$\delta K_{F_A}(\omega_1, \omega_2) = K_{F_A}(\nabla_\delta \omega_1, \omega_2) + K_{F_A}(\omega_1, \nabla_\delta \omega_2). \quad (K4)$$

(vi) For all $\omega_1, \omega_2 \in \mathcal{H}_F$,

$$u \frac{d}{du} K_{F_A}(\omega_1, \omega_2) = K_{F_A}(u \nabla_{\frac{d}{du}} \omega_1, \omega_2) + K_{F_A}(\omega_1, u \nabla_{\frac{d}{du}} \omega_2). \quad (K5)$$

Definition 2.34. Define $K_{F_A}^{(k)}$ for $k \in \mathbb{Z}$ by the coefficient of the expansion of K_{F_A} in u

$$K_{F_A}(\omega_1, \omega_2) := \sum_{k \in \mathbb{Z}} K_{F_A}^{(k)}(\omega_1, \omega_2) u^{k+3}, \quad (2.62)$$

and call it the k -th *higher residue pairing*.

Remark 2.35. In the paper [19], there were errors concerning to the signs in (K1) and (K5). We fix them here.

Proposition 2.36. *There exists a unique higher residue pairing K_{F_A} .*

Proof. Since F_A is tame at any point on the deformation space M , namely, there are no critical points coming from infinity. Therefore, the statement follows from the same argument in Theorem 2.14 and Remark 2.15 in [3]. $\square$

We shall consider the extension $\overline{K}_{F_A}$ of K_{F_A} on $\overline{\mathcal{H}}_{F_A}$.

Proposition 2.37. *The pairing K_{F_A} on $\mathcal{H}_{F_A}^{(0)}$ induces an $\mathcal{O}_{\overline{M}}$ -bilinear form*

$$\overline{K}_{F_A} : \overline{\mathcal{H}}_{F_A}^{(0)} \otimes_{\mathcal{O}_{\overline{M}}} \overline{\mathcal{H}}_{F_A}^{(0)} \longrightarrow u^3 \mathcal{O}_{\overline{M}}[[u]] \quad (2.63)$$

whose restriction to $\iota^* \left(\overline{\mathcal{H}}_{F_A}^{(0)} \right) \simeq \mathcal{H}_{F_A}^{(0)}$ coincides with K_{F_A} .

Proof. This follows from Lemma 3.4 of [8], where $\overline{M}$, M , M_∞ , $\mathcal{H}_{F_A}^{(0)}$, K_{F_A} , $\overline{\mathcal{H}}_{F_A}^{(0)}$ correspond to X , Y , D , $\mathcal{H}$, P , ${}_1\mathcal{F}$ in [8], respectively. $\square$

2.6. Primitive form.

Definition 2.38. An element $\zeta \in \Gamma(M, \mathcal{H}_{F_A}^{(0)})$ is called a *primitive form* for the tuple $(\mathcal{H}_{F_A}^{(0)}, \nabla, K_{F_A})$ if it satisfies following five conditions;

(i) $u\nabla_e \zeta = \zeta$ and ζ induces $\mathcal{O}_M$ -isomorphism:

$$\mathcal{T}_M[[u]] \simeq \mathcal{H}_{F_A}^{(0)}, \quad \sum_{k=0}^{\infty} \delta_k u^k \mapsto \sum_{k=0}^{\infty} u^k (u\nabla_{\delta_k} \zeta). \quad (\text{P1})$$

(ii) We have

$$K_{F_A}(u\nabla_{\delta} \zeta, u\nabla_{\delta'} \zeta) \in \mathbb{C} \cdot u^3, \quad \text{for all } \delta, \delta' \in \mathcal{T}_M. \quad (\text{P2})$$

(iii) There exists $r \in \mathbb{C}$ such that

$$\nabla_{u \frac{d}{du} + E} \zeta = r \zeta. \quad (\text{P3})$$

(iv) There exists a connection $\nabla / : \mathcal{T}_M \times \mathcal{T}_M \longrightarrow \mathcal{T}_M$ such that

$$u\nabla_{\delta} \nabla_{\delta'} \zeta = \nabla_{\delta \circ \delta'} \zeta + u\nabla_{\nabla_{\delta} \delta'} \zeta, \quad \delta, \delta' \in \mathcal{T}_M. \quad (\text{P4})$$

(v) There exists an $\mathcal{O}_M$ -endomorphism $N : \mathcal{T}_M \longrightarrow \mathcal{T}_M$ such that

$$u\nabla_{\frac{d}{du}} (u\nabla_{\delta} \zeta) = -\nabla_{E \circ \delta} \zeta + u\nabla_{N\delta} \zeta, \quad \delta \in \mathcal{T}_M. \quad (\text{P5})$$

In particular, the constant r of (P3) is called the *minimal exponent*.

3. MAIN THEOREM

Theorem 3.1. *The element $\zeta_A := [s_{\mu_A}^{-1} dx_1 \wedge dx_2 \wedge dx_3] \in \mathcal{H}_{F_A}^{(0)}$ is a primitive form for the tuple $(\mathcal{H}_{F_A}^{(0)}, \nabla, K_{F_A})$ with the minimal exponent $r = 1$.*

Once we have a primitive form ζ_A , we obtain a Frobenius structure on M by the general theory developed by K. Saito.

Corollary 3.2. *The primitive form ζ_A determines a Frobenius structure of rank μ_A and dimension one on the deformation space M of the universal unfolding of f_A . More precisely, the non-degenerate symmetric bilinear form η on $\mathcal{T}_M$ defined by*

$$\eta(\delta, \delta') := K_{F_A}^{(0)}(u\nabla_\delta \zeta_A, u\nabla_{\delta'} \zeta_A) = J_{F_A}(r^{(0)}(u\nabla_\delta \zeta_A), r^{(0)}(u\nabla_{\delta'} \zeta_A)), \quad \delta, \delta' \in \mathcal{T}_M, \quad (3.1)$$

together with the product $\circ$ on $\mathcal{T}_M$, the primitive vector field $e \in \Gamma(M, \mathcal{T}_M)$ and the Euler vector field $E \in \Gamma(M, \mathcal{T}_M)$ define a Frobenius structure on M of rank μ_A and dimension one.

Proof. This is a direct consequence of Theorem 7.5 of [19]. $\square$

Definition 3.3. For simplicity, we shall denote by M_{f_A, ζ_A} the deformation space M together with the Frobenius structure on M obtained in Corollary 3.2 and call it *the Frobenius manifold constructed from the pair (f_A, ζ_A) .*

For a proof of Theorem 3.1, we first give some useful lemmas.

Lemma 3.4. *For $\delta \in \mathcal{T}_M$ such that $[\delta, E] = l\delta$ for some $l \in \mathbb{Q}$, we have*

$$\nabla_{u\frac{d}{du}+E}(u\nabla_\delta \zeta_A) = (2-l) \cdot u\nabla_\delta \zeta_A, \quad (3.2)$$

In particular, we have the following equations:

$$\nabla_{u\frac{d}{du}+E}(u\nabla_e \zeta_A) = 1 \cdot u\nabla_e \zeta_A, \quad (3.3a)$$

$$\nabla_{u\frac{d}{du}+E}\left(u\nabla_{\frac{\partial}{\partial s_{i,j}}} \zeta_A\right) = \left(1 + \frac{j}{a_i}\right) \cdot u\nabla_{\frac{\partial}{\partial s_{i,j}}} \zeta_A, \quad i = 1, 2, 3, \quad j = 1, \dots, a_i - 1. \quad (3.3b)$$

$$\nabla_{u\frac{d}{du}+E}\left(u\nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A\right) = 2 \cdot u\nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A. \quad (3.3c)$$

Proof. Note that

$$\begin{aligned} (ud + dF_A) & \left(\frac{x_1}{a_1} dx_2 \wedge dx_3 - \frac{x_2}{a_2} dx_1 \wedge dx_3 + \frac{x_3}{a_3} dx_1 \wedge dx_2 \right) \\ &= \left\{ u \left(\sum_{i=1}^3 \frac{1}{a_i} \right) + \sum_{i=1}^3 \frac{1}{a_i} x_i \frac{\partial F_A}{\partial x_i} \right\} dx_1 \wedge dx_2 \wedge dx_3. \end{aligned}$$

For $\delta \in \mathcal{T}_M$ such that $[\delta, E] = l\delta$ for some $l \in \mathbb{Q}$, one has

$$\begin{aligned}
& \nabla_{u\frac{d}{du}+E}(u\nabla_\delta\zeta_A) \\
&= \nabla_{u\frac{d}{du}+E}[(\delta(F_A)s_{\mu_A}^{-1} + u\delta(s_{\mu_A}^{-1}))dx_1 \wedge dx_2 \wedge dx_3] \\
&= \left[\left\{ -\frac{1}{u}F_A\delta(F_A)s_{\mu_A}^{-1} - F_A\delta(s_{\mu_A}^{-1}) + u\delta(s_{\mu_A}^{-1}) \right\} dx_1 \wedge dx_2 \wedge dx_3 \right] \\
&\quad + \left[\left\{ \frac{1}{u}EF_A(\delta(F_A)s_{\mu_A}^{-1} + u\delta(s_{\mu_A}^{-1})) + E(\delta(F_A)s_{\mu_A}^{-1} + u\delta(s_{\mu_A}^{-1})) \right\} dx_1 \wedge dx_2 \wedge dx_3 \right] \\
&= \left[\left\{ \frac{1}{u}(EF_A - F_A)\delta(F_A)s_{\mu_A}^{-1} + E\delta(F_A)s_{\mu_A}^{-1} + \delta(F_A)E(s_{\mu_A}^{-1}) \right\} dx_1 \wedge dx_2 \wedge dx_3 \right] \\
&\quad + [\{(EF_A - F_A)\delta(s_{\mu_A}^{-1}) + uE\delta(s_{\mu_A}^{-1}) + u\delta(s_{\mu_A}^{-1})\}dx_1 \wedge dx_2 \wedge dx_3] \\
&= \left[\left\{ u \cdot \frac{1}{u} \left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} \right) + (1-l) - \chi_A \right\} \delta(F_A)s_{\mu_A}^{-1} dx_1 \wedge dx_2 \wedge dx_3 \right] \\
&\quad + \left[\left\{ u \left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} \right) - u\chi_A - ul + u \right\} \delta(s_{\mu_A}^{-1}) dx_1 \wedge dx_2 \wedge dx_3 \right] \\
&= (2-l) \cdot u\nabla_\delta\zeta_A.
\end{aligned}$$

Therefore we have Lemma 3.4. $\square$

Lemma 3.5. *Let $\delta_1, \delta_2, \delta_3$ be elements of $\mathcal{T}_M$ satisfying $[\delta_i, E] = l_i\delta_i$, $i = 1, 2, 3$ for some $l_1, l_2, l_3 \in \mathbb{Q}$. Then, we have*

$$\deg \left(K_{F_A}^{(k)} (u^2 \nabla_{\delta_1} \nabla_{\delta_2} \zeta_A, u \nabla_{\delta_3} \zeta_A) \right) = 2 - l_1 - l_2 - l_3 - k, \quad k \in \mathbb{Z}. \quad (3.4)$$

Proof. The equations (K4) and (K5) yields

$$\begin{aligned}
& \left(u \frac{d}{du} + E \right) K_{F_A} (u^2 \nabla_{\delta_1} \nabla_{\delta_2} \zeta_A, u \nabla_{\delta_3} \zeta_A) \\
&= K_{F_A} \left(\nabla_{u\frac{d}{du}+E}(u\nabla_{\delta_1}(u\nabla_{\delta_2}\zeta_A)), u\nabla_{\delta_3}\zeta_A \right) + K_{F_A} \left(u^2 \nabla_{\delta_1} \nabla_{\delta_2} \zeta_A, \nabla_{u\frac{d}{du}+E}(u\nabla_{\delta_3}\zeta_A) \right).
\end{aligned}$$

By the integrability of the Gauß–Manin connection, we have

$$\left[\nabla_{u\frac{d}{du}+E}, u\nabla_{\delta_1} \right] = (1-l_1)u\nabla_{\delta_1}.$$

The statement now easily follows from the definition of $K_{F_A}^{(k)}$ and Lemma 3.4. $\square$

3.1. Condition (i). Condition (i) is satisfied since for all $\delta \in \mathcal{T}_M$ we have

$$u\nabla_\delta[s_{\mu_A}^{-1}dx_1 \wedge dx_2 \wedge dx_3] = [(\delta F + us_{\mu_A}\delta(s_{\mu_A}^{-1}))s_{\mu_A}^{-1}dx_1 \wedge dx_2 \wedge dx_3]. \quad (3.5)$$

3.2. **Condition (ii).** It is enough to show the following:

Lemma 3.6. *Let δ_1, δ_2 be elements of the set of vector fields*

$$\left\{ e = \frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_{1,1}}, \dots, \frac{\partial}{\partial s_{i,j}}, \dots, \frac{\partial}{\partial s_{3,a_3-1}}, s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}} \right\}.$$

Then, for all $k \geq 1$, $K_{F_A}^{(k)}(u\nabla_{\delta_1}\zeta_A, u\nabla_{\delta_2}\zeta_A) = 0$.

Proof. By Lemma 3.5, we have

$$\deg \left(K_{F_A}^{(k)}(u\nabla_{\delta_1}\zeta_A, u\nabla_{\delta_2}\zeta_A) \right) = 1 - l_1 - l_2 - k, \quad k \in \mathbb{Z}.$$

Therefore, $\deg \left(K_{F_A}^{(k)}(u\nabla_{\delta_1}\zeta_A, u\nabla_{\delta_2}\zeta_A) \right)$ is zero if $\delta_1 = \delta_2 = s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}$, $k = 1$ and is negative otherwise. If the degree is negative, then since $K_{F_A}^{(k)}(u\nabla_{\delta_1}\zeta_A, u\nabla_{\delta_2}\zeta_A) \in \mathcal{O}_{\overline{M}}$ by Proposition 2.37, it must be zero. Since the first higher residue pairing $K_{F_A}^{(1)}$ is skew-symmetric by the equation (K1), we have

$$K_{F_A}^{(1)} \left(u\nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta, u\nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta \right) = 0.$$

□

3.3. **Condition (iii).** This is already proven in Lemma 3.4.

3.4. **Condition (iv).** We shall prepare the following useful lemma to check Condition (iv):

Lemma 3.7. *Under the conditions (P1) and (P2), the following two conditions are equivalent for $\delta_1, \delta_2 \in \mathcal{T}_M$:*

- (i) *The equation (P4) holds.*
- (ii) *For all $\delta_3 \in \mathcal{T}_M$ and $k \geq 2$, $K_{F_A}^{(k)}(u^2\nabla_{\delta_1}\nabla_{\delta_2}\zeta_A, u\nabla_{\delta_3}\zeta_A) = 0$.*

Proof. (i) $\Rightarrow$ (ii). It immediately follows from Lemma 3.6.

(ii) $\Rightarrow$ (i). Recall that the $K^{(0)}(u\nabla_{\delta}\zeta_A, u\nabla_{\delta'}\zeta_A) = J_{F_A}(r^{(0)}(u\nabla_{\delta}\zeta_A), r^{(0)}(u\nabla_{\delta'}\zeta_A))$ by (K3). Since J_{F_A} is non-degenerate, the conditions (P1) and (P2) yield the statement.

□

By Lemma 3.7, it is enough to show the following:

Lemma 3.8. *Let $\delta_1, \delta_2, \delta_3$ be elements of the set of vector fields*

$$\left\{ e = \frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_{1,1}}, \dots, \frac{\partial}{\partial s_{i,j}}, \dots, \frac{\partial}{\partial s_{3,a_3-1}}, s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}} \right\}.$$

Then, $K_{F_A}^{(k)}(u^2\nabla_{\delta_1}\nabla_{\delta_2}\zeta_A, u\nabla_{\delta_3}\zeta_A) = 0$ for all $\delta_1, \delta_2, \delta_3 \in \mathcal{T}_M$ and $k \geq 2$ except for one case when $\delta_1 = \delta_2 = \delta_3 = s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}$, $k = 2$.

Proof. By Lemma 3.5, for all $k \geq 2$, $\deg \left(K_{FA}^{(k)} (u^2 \nabla_{\delta_1} \nabla_{\delta_2} \zeta_A, u \nabla_{\delta_3} \zeta_A) \right) = 0$ if $\delta_1 = \delta_2 = \delta_3 = s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}$, $k = 2$ and is negative otherwise. If the degree is negative, then since $K_{FA}^{(k)} (u^2 \nabla_{\delta_1} \nabla_{\delta_2} \zeta_A, u \nabla_{\delta_3} \zeta_A) \in \mathcal{O}_{\overline{M}}$ by Proposition 2.37, it must be zero.

If $\delta_1 = \delta_2 = \delta_3 = s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}$ and $k = 2$, $K_{FA}^{(2)} \left(u^2 \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A, u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right)$ is a constant. Then we only have to calculate this value along the submanifold $\{\mathbf{s} = \mathbf{0}\}$.

By Lemma 3.4, one has $\nabla_{u \frac{d}{du} + E} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) = 2u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A$. This implies

$$\begin{aligned} \chi_A \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) &= 2u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A - \nabla_{u \frac{d}{du}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \\ &\quad - \nabla_{E - \chi_A s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right). \end{aligned}$$

Then, by $\chi_A \neq 0$ and Lemma 3.6, one has

$$\begin{aligned} &K_{FA}^{(2)} \left(u^2 \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A, u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \Big|_{\mathbf{s}=\mathbf{0}} \\ &= \frac{1}{\chi_A} K_{FA}^{(2)} \left(2u^2 \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A, u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \Big|_{\mathbf{s}=\mathbf{0}} \\ &\quad - \frac{1}{\chi_A} K_{FA}^{(2)} \left(u^2 \nabla_{u \frac{d}{du}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right), u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \Big|_{\mathbf{s}=\mathbf{0}} \\ &= - \frac{1}{\chi_A} K_{FA}^{(0)} \left(\nabla_{u \frac{d}{du}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right), u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \Big|_{\mathbf{s}=\mathbf{0}}. \end{aligned}$$

By the symmetry of $K_{FA}^{(0)}$ and the equation (K5), one also has

$$\begin{aligned} &K_{FA}^{(0)} \left(\nabla_{u \frac{d}{du}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right), u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \\ &= \frac{1}{2} \left\{ K_{FA}^{(0)} \left(\nabla_{u \frac{d}{du}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right), u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \right. \\ &\quad \left. + K_{FA}^{(0)} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A, \nabla_{u \frac{d}{du}} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) \right) \right\} \\ &= \frac{3}{2} K_{FA}^{(0)} \left(u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A, u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A \right) = 0. \end{aligned}$$

Therefore we have Lemma 3.8. $\square$

3.5. Condition (v). Under the Condition (iv), it is easy to see that Condition (v) follows from Lemma 3.4.

4. MIRROR SYMMETRY

From now on, we shall denote by $\mathcal{F}_{f_A, \zeta_A}$ the Frobenius potential for the Frobenius structure on M constructed from the pair (f_A, ζ_A) in Corollary 3.2.

Theorem 4.1. *For the Frobenius manifold M_{f_A, ζ_A} constructed from the pair (f_A, ζ_A) , there exists flat coordinates $t_1, t_{1,1}, \dots, t_{3,a_3-1}, t_{\mu_A}$ satisfying the following conditions:*

(i) *The unit vector field e and the Euler vector field E are given by*

$$e = \frac{\partial}{\partial t_1}, \quad E = t_1 \frac{\partial}{\partial t_1} + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} \frac{a_i - j}{a_i} t_{i,j} \frac{\partial}{\partial t_{i,j}} + \chi_A \frac{\partial}{\partial t_{\mu_A}}.$$

(ii) *The non-degenerate symmetric bilinear form η on $\mathcal{T}_M$ defined by*

$$\eta(\delta, \delta') := K_{F_A}^{(0)}(u \nabla_\delta \zeta_A, u \nabla_{\delta'} \zeta), \quad \delta, \delta' \in \mathcal{T}_M, \quad (4.1)$$

satisfies

$$\begin{aligned} \eta\left(\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_{\mu_A}}\right) &= \eta\left(\frac{\partial}{\partial t_{\mu_A}}, \frac{\partial}{\partial t_1}\right) = 1, \\ \eta\left(\frac{\partial}{\partial t_{i_1, j_1}}, \frac{\partial}{\partial t_{i_2, j_2}}\right) &= \begin{cases} \frac{1}{a_{i_1}} & i_1 = i_2 \text{ and } j_2 = a_{i_1} - j_1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

(iii) *The Frobenius potential $\mathcal{F}_{f_A, \zeta_A}$ satisfies $E\mathcal{F}_{f_A, \zeta_A}|_{t_1=0} = 2\mathcal{F}_{f_A, \zeta_A}|_{t_1=0}$,*

$$\mathcal{F}_{f_A, \zeta_A}|_{t_1=0} \in \mathbb{C}[t_{1,1}, \dots, t_{1,a_1-1}, t_{2,1}, \dots, t_{2,a_2-1}, t_{3,1}, \dots, t_{3,a_3-1}, e^{t_{\mu_A}}].$$

(iv) *Assume the condition (iii). The restriction of the Frobenius potential $\mathcal{F}_{f_A, \zeta_A}$ to the submanifold $\{t_1 = e^{t_{\mu_A}} = 0\}$ is given as*

$$\mathcal{F}_{f_A, \zeta_A}|_{t_1=e^{t_{\mu_A}}=0} = \mathcal{G}^{(1)} + \mathcal{G}^{(2)} + \mathcal{G}^{(3)},$$

where $\mathcal{G}^{(i)} \in \mathbb{C}[t_{i,1}, \dots, t_{i,a_i-1}]$, $i = 1, 2, 3$.

(v) *Assume the condition (iii). In the frame $\frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_{1,1}}, \dots, \frac{\partial}{\partial t_{3,a_3-1}}, \frac{\partial}{\partial t_{\mu_A}}$ of $\mathcal{T}_M$, the product $\circ$ can be extended to the limit $t_1 = t_{1,1} = \dots = t_{3,a_3-1} = e^{t_{\mu_A}} = 0$. The $\mathbb{C}$ -algebra obtained in this limit is isomorphic to*

$$\mathbb{C}[x_1, x_2, x_3] / (x_1 x_2, x_2 x_3, x_3 x_1, a_1 x_1^{a_1} - a_2 x_2^{a_2}, a_2 x_2^{a_2} - a_3 x_3^{a_3}),$$

where $\partial/\partial t_{1,1}, \partial/\partial t_{2,1}, \partial/\partial t_{3,1}$ are mapped to x_1, x_2, x_3 , respectively.

(vi) *The term*

$$\begin{cases} e^{t_{\mu_A}} & \text{if } a_1 = a_2 = a_3 = 1, \\ t_{3,1} e^{t_{\mu_A}} & \text{if } 1 = a_1 = a_2 < a_3, \\ t_{2,1} t_{3,1} e^{t_{\mu_A}} & \text{if } 1 = a_1 < a_2, \\ t_{1,1} t_{2,1} t_{3,1} e^{t_{\mu_A}} & \text{if } a_1 = 2, \end{cases}$$

occurs with the coefficient 1 in $\mathcal{F}_{f_A, \zeta_A}$.

We shall check all conditions in the above theorem.

4.1. Condition (i).

Lemma 4.2. *We have*

$$K_{FA}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial s_1}} \zeta_A) = 0, \quad K_{FA}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial s_{i,j}}} \zeta_A) = 0, \quad i = 1, 2, 3, j = 1, \dots, a_i - 1, \quad (4.2)$$

and

$$K_{FA}^{(0)}(\zeta_A, u \nabla_{s_{\mu_A} \frac{\partial}{\partial s_{\mu_A}}} \zeta_A) = 1. \quad (4.3)$$

Proof. The statement follows from Lemma 2.31. $\square$

Lemma 4.3. *The one form $\theta \in \Gamma(M, \Omega_M^1)$ defined by*

$$\begin{aligned} \theta := & K_{FA}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial s_1}} \zeta_A) ds_1 \\ & + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} K_{FA}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial s_{i,j}}} \zeta_A) ds_{i,j} + K_{FA}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial s_{\mu_A}}} \zeta_A) ds_{\mu_A}, \end{aligned} \quad (4.4)$$

is a closed form which is independent from the choice of coordinates on M . Moreover, there exists a flat coordinate t such that $\theta = dt$.

Proof. See section 3.3 3) of [16]. $\square$

Combining these two Lemmas, we have $dt = ds_{\mu_A}/s_{\mu_A}$. Therefore, we can choose $t_{\mu_A} := \log s_{\mu_A}$ as a flat coordinate. Since $\zeta_A, x_i^j \zeta_A (i = 1, 2, 3, j = 1, \dots, a_i - 1), s_{\mu_A}^{-1} x_1 x_2 x_3 \zeta_A$ form a $\mathbb{C}[[u]]$ -basis of $\overline{\mathcal{H}}_{FA}^{(0)}$ at $(\mathbf{0}, 0) \in \overline{M}$, by Lemma 3.4, one can choose other flat coordinates $t_1, t_{1,1}, \dots, t_{3,a_3-1}$ such that

$$t_1|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = t_{1,1}|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = \dots = t_{3,a_3-1}|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = 0$$

together with the following normalization;

$$\frac{\partial t_1}{\partial s_1} \Big|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = 1, \quad \frac{\partial t_1}{\partial s_{i,j}} \Big|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = 0, \quad \frac{\partial t_1}{\partial s_{\mu_A}} \Big|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = 0, \quad (4.5a)$$

$$\frac{\partial t_{i,j}}{\partial s_1} \Big|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = 0, \quad \frac{\partial t_{i,j}}{\partial s_{i',j'}} \Big|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = \delta_{ii'} \delta_{jj'}, \quad \frac{\partial t_{i,j}}{\partial s_{\mu_A}} \Big|_{(s, s_{\mu_A})=(\mathbf{0}, 0)} = 0, \quad (4.5b)$$

where $\delta_{ii'}$ and $\delta_{jj'}$ are Kronecker's deltas. In particular, the flat coordinates $t_1, t_{1,1}, \dots, t_{3,a_3-1}, t_{\mu_A}$ satisfy

$$e = \frac{\partial}{\partial t_1}, \quad E = t_1 \frac{\partial}{\partial t_1} + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} \frac{a_i - j}{a_i} t_{i,j} \frac{\partial}{\partial t_{i,j}} + \chi_A \frac{\partial}{\partial t_{\mu_A}},$$

which is the condition (i).

4.2. **Condition (ii).**

Lemma 4.4. *We have*

$$K_{F_A}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial t_1}} \zeta_A) = 0, \quad K_{F_A}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial t_{i,j}}} \zeta_A) = 0, \quad i = 1, 2, 3, j = 1, \dots, a_i - 1, \quad (4.6)$$

and

$$K_{F_A}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial t_{\mu_A}}} \zeta_A) = 1. \quad (4.7)$$

Proof. By Lemma 4.3, we have

$$dt_{\mu_A} = K_{F_A}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial t_1}} \zeta_A) dt_1 + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} K_{F_A}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial t_{i,j}}} \zeta_A) dt_{i,j} + K_{F_A}^{(0)}(\zeta_A, u \nabla_{\frac{\partial}{\partial t_{\mu_A}}} \zeta_A) dt_{\mu_A}.$$

The statement follows. $\square$

Note that the pairings to consider are constant since we are dealing with flat coordinates. Therefore, we can evaluate them at $(\mathbf{0}, 0) \in \overline{M}$. Moreover, by the normalization (4.5), we have

$$\begin{aligned} & \lim_{s_{\mu_A} \rightarrow 0} K_{F_A}^{(0)} \left(u \nabla_{\frac{\partial}{\partial t_{i,j}}} \zeta_A, u \nabla_{\frac{\partial}{\partial t_{i',j'}}} \zeta_A \right) \Big|_{\mathbf{s}=0} \\ &= \overline{K}_{F_A}^{(0)} \left(u \nabla_{\frac{\partial}{\partial t_{i,j}}} \zeta_A, u \nabla_{\frac{\partial}{\partial t_{i',j'}}} \zeta_A \right) \Big|_{(\mathbf{s}, s_{\mu_A}) = (\mathbf{0}, 0)} \\ &= \overline{K}_{F_A}^{(0)} \left(u \nabla_{\frac{\partial}{\partial s_{i,j}}} \zeta_A, u \nabla_{\frac{\partial}{\partial s_{i',j'}}} \zeta_A \right) \Big|_{(\mathbf{s}, s_{\mu_A}) = (\mathbf{0}, 0)} \\ &= \lim_{s_{\mu_A} \rightarrow 0} K_{F_A}^{(0)} \left(u \nabla_{\frac{\partial}{\partial s_{i,j}}} \zeta_A, u \nabla_{\frac{\partial}{\partial s_{i',j'}}} \zeta_A \right) \Big|_{\mathbf{s}=0} \\ &= \lim_{s_{\mu_A} \rightarrow 0} K_{F_A}^{(0)} \left(\zeta_A, u \nabla_{\frac{\partial}{\partial s_{i,j}} \circ \frac{\partial}{\partial s_{i',j'}}} \zeta_A \right) \Big|_{\mathbf{s}=0}. \end{aligned}$$

The statement follows from Lemma 2.7.

4.3. **Condition (iii).** Note that we have an $\mathcal{O}_{\overline{M}}$ -isomorphism

$$\tau_{\overline{M}}(-\log M_{\infty}) \simeq \mathcal{O}_{\overline{M}} \frac{\partial}{\partial t_1} \bigoplus_{\substack{1 \leq i \leq 3, \\ 1 \leq j \leq a_i - 1}} \mathcal{O}_{\overline{M}} \frac{\partial}{\partial t_{i,j}} \bigoplus \mathcal{O}_{\overline{M}} \frac{\partial}{\partial t_{\mu_A}}. \quad (4.8)$$

The condition (iii) follows from this together with Proposition 2.9 and the fact that all flat coordinates have positive degrees.

4.4. **Condition** (iv). Recall that the ideal in the equation (2.9) restricted to M_∞ is given by

$$(x_2x_3, x_3x_1, x_1x_2, H_1(\mathbf{x}, \mathbf{s}), H_2(\mathbf{x}, \mathbf{s})),$$

where

$$H_i(\mathbf{x}, \mathbf{s}) := a_i x_i^{a_i} - a_{i+1} x_{i+1}^{a_{i+1}} + \sum_{j=1}^{a_i-1} j \cdot s_{i,j} \cdot x_i^j - \sum_{j=1}^{a_{i+1}-1} j \cdot s_{i+1,j} \cdot x_{i+1}^j, \quad i = 1, 2.$$

In particular, we have

$$\eta \left(\frac{\partial}{\partial s_{i,j}}, \frac{\partial}{\partial s_{i',j'}} \right) \Big|_{s_{\mu_A}=0} = 0, \quad \text{if } i \neq i'. \quad (4.9)$$

Note that the connection ∇ on $\mathcal{T}_M$ in (P4) is the Levi-Civita connection with respect to η (see Proposition 7.9 and Proposition 7.16 of [19]) and that $t_{i,j}$ is the solution of the system

$$\begin{aligned} \nabla^* dt_{i,j} &= 0, \quad t_{i,j}|_{(\mathbf{s}, s_{\mu_A})=(\mathbf{0},0)} = 0, \\ \frac{\partial t_{i,j}}{\partial s_1} \Big|_{(\mathbf{s}, s_{\mu_A})=(\mathbf{0},0)} &= 0, \quad \frac{\partial t_{i,j}}{\partial s_{i',j'}} \Big|_{(\mathbf{s}, s_{\mu_A})=(\mathbf{0},0)} = \delta_{ii'} \delta_{jj'}, \quad \frac{\partial t_{i,j}}{\partial s_{\mu_A}} \Big|_{(\mathbf{s}, s_{\mu_A})=(\mathbf{0},0)} = 0, \end{aligned}$$

where ∇^* is a connection on Ω_M^1 dual to ∇ . Therefore, by (4.9), we have

$$\frac{\partial t_{i,j}}{\partial s_{i',j'}} \Big|_{s_{\mu_A}=0} = 0, \quad \text{if } i \neq i'. \quad (4.10)$$

The third derivatives of the Frobenius potential with respect to flat coordinates are given by residues. For example, we have

$$\frac{\partial^3 \mathcal{F}_{f_A, \zeta_A}}{\partial t_{i_1, j_1} \partial t_{i_2, j_2} \partial t_{i_3, j_3}} = -e^{-2t_{\mu_A}} \text{Res}_{\mathbb{C}^3 \times M/M} \left[\frac{\frac{\partial F_A}{\partial t_{i_1, j_1}} \frac{\partial F_A}{\partial t_{i_2, j_2}} \frac{\partial F_A}{\partial t_{i_3, j_3}} dx_1 \wedge dx_2 \wedge dx_3}{\frac{\partial F_A}{\partial x_1} \frac{\partial F_A}{\partial x_2} \frac{\partial F_A}{\partial x_3}} \right]. \quad (4.11)$$

Therefore, by using the above description of the ideal, the normalization (4.5) and (4.10), we can show that

$$\lim_{e^{t_{\mu_A}} \rightarrow 0} \frac{\partial^n \mathcal{F}_{f_A, \zeta_A}}{\partial t_{i_1, j_1} \cdots \partial t_{i_n, j_n}} \Big|_{t_1=t_{1,1}=\cdots=t_{3,a_3-1}=0} \neq 0 \text{ only if } i_1 = \cdots = i_n,$$

by induction on n .

4.5. **Condition** (v). Taking into account the normalization (4.5), the condition (v) easily follows from Lemma 2.7.

4.6. **Condition (vi).** Note that the coefficient of the term $e^{t_{\mu_A}}$ (if $a_1 = a_2 = a_3 = 1$), $t_{3,1}e^{t_{\mu_A}}$ (if $1 = a_1 = a_2 < a_3$), $t_{2,1}t_{3,1}e^{t_{\mu_A}}$ (if $1 = a_1 < a_2$), $t_{1,1}t_{2,1}t_{3,1}e^{t_{\mu_A}}$ (if $a_1 \geq 2$) is given by the limit

$$\begin{aligned} & \lim_{e^{t_{\mu_A}} \rightarrow 0} \left(e^{-t_{\mu_A}} \cdot \frac{\partial^3 \mathcal{F}_{f_A, \zeta_A}}{\partial t_{\mu_A} \partial t_{\mu_A} \partial t_{\mu_A}} \Big|_{t_1=0} \right) \text{ if } a_1 = a_2 = a_3 = 1, \\ & \lim_{e^{t_{\mu_A}} \rightarrow 0} \left(e^{-t_{\mu_A}} \cdot \frac{\partial^3 \mathcal{F}_{f_A, \zeta_A}}{\partial t_{\mu_A} \partial t_{\mu_A} \partial t_{3,1}} \Big|_{t_1=t_{3,1}=\dots=t_{3,a_3-1}=0} \right) \text{ if } 1 = a_1 = a_2 < a_3, \\ & \lim_{e^{t_{\mu_A}} \rightarrow 0} \left(e^{-t_{\mu_A}} \cdot \frac{\partial^3 \mathcal{F}_{f_A, \zeta_A}}{\partial t_{\mu_A} \partial t_{2,1} \partial t_{3,1}} \Big|_{t_1=t_{2,1}=\dots=t_{3,a_3-1}=0} \right) \text{ if } 1 = a_1 < a_2, \\ & \lim_{e^{t_{\mu_A}} \rightarrow 0} \left(e^{-t_{\mu_A}} \cdot \frac{\partial^3 \mathcal{F}_{f_A, \zeta_A}}{\partial t_{1,1} \partial t_{2,1} \partial t_{3,1}} \Big|_{t_1=t_{1,1}=\dots=t_{3,a_3-1}=0} \right) \text{ if } a_1 \geq 2. \end{aligned}$$

Since the third derivatives of the Frobenius potential with respect to flat coordinates are given by residues of the form (4.11), it is easy to see that the above limits are always reduced to the following limit

$$\lim_{s_{\mu_A} \rightarrow 0} \left(s_{\mu_A}^{-1} \cdot s_{\mu_A}^{-2} \text{Res}_{\mathbb{C}^3 \times M/M} \left[\frac{x_1 x_2 x_3 dx_1 \wedge dx_2 \wedge dx_3}{\frac{\partial F_A}{\partial x_1} \frac{\partial F_A}{\partial x_2} \frac{\partial F_A}{\partial x_3}} \right] \Big|_{\mathbf{s}=0} \right)$$

by using the relations in $p_*\mathcal{O}_C$ and the normalization (4.5), which is one.

4.7. **Mirror isomorphism.** This theorem enables us to simplify the proofs given by Milanov–Tseng [11] and Rossi [13].

Corollary 4.5. *There exists an isomorphism of Frobenius manifolds between the one constructed from the Gromov–Witten theory for $\mathbb{P}_A^1$ and the one constructed from the pair (f_A, ζ_A) .*

5. ANOTHER PROOF OF OUR MAIN THEOREM

Denote by $(\eta_A, \circ_A, e_A, E_A)$ the Frobenius structure for a triplet of positive integers $A = (a_1, a_2, a_3)$ with $\chi_A > 0$ uniquely determined by Theorem 3.1 of [9]. In particular, we shall denote by $(\mathbf{t}, t_{\mu_A}) := (t_1, t_2, \dots, t_{\mu_A-1}, t_{\mu_A}) := (t_1, t_{1,1}, \dots, t_{3,a_3-1}, t_{\mu_A})$ the flat coordinates on M and also denote by ∂_i the basis of $\mathcal{T}_M$ corresponding to the coordinate t_i for simplicity.

In this section, we shall show the way to determine not only the Frobenius structure $(\eta, \circ, e, E)$ on M constructed from the pair (f_A, ζ_A) but also the primitive form ζ_A itself from the Frobenius structure $(\eta_A, \circ_A, e_A, E_A)$. It is very important to note that the

Frobenius potential $\mathcal{F}_A$ for $(\eta_A, \circ_A, e_A, E_A)$ is isomorphic to the one on M constructed from the Gromov-Witten theory of the orbifold $\mathbb{P}_A^1$.

5.1. Key lemma. We can determine a primitive form by the following Lemma, which is essentially Lemma 3.3 of [16].

Lemma 5.1. *Suppose that there exists a local coordinates $(t_1, t_{1,1}, \dots, t_{3,a_3-1}, t_{\mu_A})$ on M and an element $\zeta \in \Gamma(M, \mathcal{H}_{F_A}^{(0)})$ satisfying the following properties:*

(i) *The unit vector field e and the Euler vector field E are given by*

$$e = \frac{\partial}{\partial t_1}, \quad E = t_1 \frac{\partial}{\partial t_1} + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} \frac{a_i - j}{a_i} t_{i,j} \frac{\partial}{\partial t_{i,j}} + \chi_A \frac{\partial}{\partial t_{\mu_A}}.$$

(ii) *ζ satisfies*

$$u \nabla_e \zeta = \zeta, \quad \nabla_{u \frac{d}{du} + E} \zeta = \zeta.$$

(iii) *We have an $\mathcal{O}_M[[u]]$ -isomorphism*

$$\mathcal{H}_{F_A}^{(0)} \simeq \mathcal{O}_M[[u]] \cdot u \nabla_{\frac{\partial}{\partial t_1}} \zeta \bigoplus_{\substack{1 \leq i \leq 3, \\ 1 \leq j \leq a_i - 1}} \mathcal{O}_M[[u]] \cdot u \nabla_{\frac{\partial}{\partial t_{i,j}}} \zeta \bigoplus \mathcal{O}_M[[u]] \cdot u \nabla_{\frac{\partial}{\partial t_{\mu_A}}} \zeta.$$

(iv) *Let V be the $\mathbb{C}$ -vector space given by*

$$V := \mathbb{C} \frac{\partial}{\partial t_1} \bigoplus_{\substack{1 \leq i \leq 3, \\ 1 \leq j \leq a_i - 1}} \mathbb{C} \frac{\partial}{\partial t_{i,j}} \bigoplus \mathbb{C} \frac{\partial}{\partial t_{\mu_A}}.$$

We have

$$K_{F_A}^{(0)}(u \nabla_{\delta_1} \zeta, u \nabla_{\delta_2} \zeta) \in \mathbb{C}, \quad \text{for all } \delta_1, \delta_2 \in V.$$

(v) *We have*

$$u \nabla_{\delta_1} \nabla_{\delta_2} \zeta = \nabla_{\delta_1 \circ \delta_2} \zeta, \quad \text{for all } \delta_1, \delta_2 \in V.$$

Then, the $\mathcal{O}_M$ -bilinear form

$$K'_{F_A} : \mathcal{H}_{F_A} \otimes_{\mathcal{O}_M} \mathcal{H}_{F_A} \longrightarrow \mathcal{O}_M((u)) \quad (5.1)$$

defined by

$$K'_{F_A}(u^{i_1+1} \nabla_{\delta_1} \zeta, u^{i_2+1} \nabla_{\delta_2} \zeta) := (-1)^{i_2} \cdot K_{F_A}^{(0)}(u \nabla_{\delta_1} \zeta, u \nabla_{\delta_2} \zeta) \cdot u^{i_1+i_2+3}, \quad \delta_1, \delta_2 \in \mathcal{T}_M, \quad (5.2)$$

coincides with K_{F_A} . In particular, ζ is a primitive form for the tuple $(\mathcal{H}_{F_A}^{(0)}, \nabla, K_{F_A})$ with the minimal exponent $r = 1$.

Proof. The pairing K'_{F_A} obviously satisfies the same conditions (K1), (K2), (K3) as the higher residue pairing K_{F_A} does. We can easily show by a direct calculation that the conditions (iv) and (v) imply the condition (K4) for K'_{F_A} . We have

$$\nabla_{u \frac{d}{du} + E} \left(u \nabla_{\frac{\partial}{\partial t_{i,j}}} \zeta \right) = \left(\frac{j}{a_i} + 1 \right) \cdot u \nabla_{\frac{\partial}{\partial t_{i,j}}} \zeta, \quad i = 1, 2, 3, \quad j = 1, \dots, a_i - 1,$$

$$\nabla_{u \frac{d}{du} + E} \left(u \nabla_{\frac{\partial}{\partial t_{\mu_A}}} \zeta \right) = 2 \cdot u \nabla_{\frac{\partial}{\partial t_{\mu_A}}} \zeta,$$

which implies the condition (P5) for ζ . By using this, we can also show the condition (K5) for K'_{F_A} . Therefore, by the uniqueness of the higher residue pairings in Proposition 2.36, we have $K'_{F_A} = K_{F_A}$.

ζ obviously satisfies the conditions (P1), (P3) and (P4) by the assumption. Since

$$K_{F_A}(u \nabla_{\delta_1} \zeta, u \nabla_{\delta_2} \zeta) = K'_{F_A}(u \nabla_{\delta_1} \zeta, u \nabla_{\delta_2} \zeta) := K_{F_A}^{(0)}(u \nabla_{\delta_1} \zeta, u \nabla_{\delta_2} \zeta) \cdot u^3, \quad \delta_1, \delta_2 \in \mathcal{T}_M,$$

ζ also satisfies the condition (P2). The statement follows. $\square$

5.2. From the Frobenius potential $\mathcal{F}_A$ to the primitive form ζ_A . In our case, the conditions for $\zeta = \zeta_A$ in Lemma 5.1 can be reduced to the assumption in the following lemma:

Lemma 5.2. *Assume that we have a homogeneous change of coordinates $(\mathbf{t}, e^{t_{\mu_A}}) \mapsto (\mathbf{s}, s_{\mu_A}) = (\mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})$ on $\overline{M}$,*

$$E_A s_1(\mathbf{t}, e^{t_{\mu_A}}) = s_1(\mathbf{t}, e^{t_{\mu_A}}), \quad (5.3a)$$

$$E_A s_{i,j}(\mathbf{t}, e^{t_{\mu_A}}) = \frac{a_i - j}{a_i} s_{i,j}(\mathbf{t}, e^{t_{\mu_A}}), \quad i = 1, 2, 3, \quad j = 1, \dots, j = a_i - 1, \quad (5.3b)$$

such that there exist functions

$$\phi_{ij}^a(\mathbf{x}; \mathbf{t}, e^{t_{\mu_A}}), \psi_a(\mathbf{x}; \mathbf{t}, e^{t_{\mu_A}}) \in \Gamma(M, \mathcal{O}_M)[x_1, x_2, x_3], \quad a = 1, 2, 3, \quad i, j = 1, \dots, \mu_A$$

satisfying the following equations:

$$\begin{aligned} & \frac{\partial F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial t_i} \cdot \frac{\partial F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial t_j} - \sum_{k=1}^{\mu_A} C_{ij}^k \cdot \frac{\partial F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial t_k} \\ &= \sum_{a=1}^3 \phi_{ij}^a(\mathbf{x}; \mathbf{t}, e^{t_{\mu_A}}) \cdot \frac{\partial F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial x_a}, \quad (5.4a) \end{aligned}$$

$$\begin{aligned}
& \frac{\partial^2 F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial t_i \partial t_j} + \eta_{ij} - \delta_{i\mu} \cdot \frac{\partial F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial t_j} \\
& - \delta_{j\mu} \cdot \frac{\partial F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial t_i} - \sum_{a=1}^3 \frac{\partial \phi_{ij}^a(\mathbf{x}; \mathbf{t}, e^{t_{\mu_A}})}{\partial x_a} \\
& = \delta_{i\mu} \cdot \delta_{j\mu} \cdot \sum_{a=1}^3 \psi^a(\mathbf{x}; \mathbf{t}, e^{t_{\mu_A}}) \cdot \frac{\partial F_A(\mathbf{x}; \mathbf{s}(\mathbf{t}, e^{t_{\mu_A}}), e^{t_{\mu_A}})}{\partial x_a}, \quad (5.4b)
\end{aligned}$$

$$\sum_{a=1}^3 \frac{\partial \psi_a(\mathbf{x}; \mathbf{t}, e^{t_{\mu_A}})}{\partial x_a} = 1, \quad (5.4c)$$

where $(\eta^{ij}) := (\eta_A(\partial_i, \partial_j))^{-1}$ and $C_{ij}^k = \sum_{l=1}^{\mu_A} \eta^{lk} \partial_l \partial_i \partial_j \mathcal{F}_A$.

Then, we have $e = e_A$, $E = E_A$, where e and E are the primitive vector field and the Euler vector field defined in Definition 2.10, and we have the equations

$$u \nabla_{\frac{\partial}{\partial t_i}} \nabla_{\frac{\partial}{\partial t_j}} \zeta_A = \sum_{k=1}^{\mu_A} C_{ij}^k \nabla_{\frac{\partial}{\partial t_k}} \zeta_A, \quad i, j = 1, \dots, \mu_A.$$

In particular, the product $\circ_A$ on $\mathcal{T}_M$ coincides with the product $\circ$ on $\mathcal{T}_M$ defined in Definition 2.5.

Proof. Note that the degrees of $t_1, \dots, t_{\mu_A-1}$ and $e^{t_{\mu_A}}$ are all positive since we assume $\chi_A > 0$. Therefore, the homogeneity (5.3) implies that

$$e_A = \frac{\partial}{\partial s_1}, \quad E_A = s_1 \frac{\partial}{\partial s_1} + \sum_{i=1}^3 \sum_{j=1}^{a_i-1} \frac{a_i - j}{a_i} s_{i,j} \frac{\partial}{\partial s_{i,j}} + \chi_A \frac{\partial}{\partial t_{\mu_A}}.$$

Hence, we have $e = e_A$ and $E = E_A$.

Based on the definition of the Gauß–Manin connection ∇ , one can show that the equation $u \nabla_{\partial_i} \nabla_{\partial_j} \zeta_A = \sum_{k=1}^{\mu_A} C_{ij}^k \nabla_{\partial_k} \zeta_A$ is satisfied if and only if

$$\begin{aligned}
& \left[\left(\partial_i F_A \partial_j F_A - \sum_{k=1}^{\mu} C_{ij}^k \partial_k F_A \right) e^{-t_{\mu_A}} dx_1 \wedge dx_2 \wedge dx_3 \right] \\
& + u \left[(\partial_i \partial_j F_A + \eta_{ij} - \delta_{i\mu} \partial_j F_A - \delta_{j\mu} \partial_i F_A) e^{-t_{\mu_A}} dx_1 \wedge dx_2 \wedge dx_3 \right] \\
& + u^2 \left[\delta_{i\mu} \delta_{j\mu} e^{-t_{\mu_A}} dx_1 \wedge dx_2 \wedge dx_3 \right] = 0. \quad (5.5)
\end{aligned}$$

The rest is straightforward. $\square$

In our case, Lemma 5.2 also enables us to compare η_A and η defined in Corollary 3.2.

Lemma 5.3. *Suppose that Lemma 5.2 holds. Then, we have the equation*

$$K_{F_A}^{(0)}(u\nabla_{\partial_i}\zeta_A, u\nabla_{\partial_j}\zeta_A) = \eta_A(\partial_i, \partial_j), \quad i, j = 1, \dots, \mu_A.$$

Proof. It is easy to calculate that $K_{F_A}^{(0)}(u\nabla_{\partial_i}\zeta_A, u\nabla_{\partial_j}\zeta_A) = \sum_{k=1}^{\mu_A} C_{ij}^k K_{F_A}^{(0)}(\zeta_A, u\nabla_{\partial_k}\zeta_A)$. We can check that $K_{F_A}^{(0)}(\zeta_A, u\nabla_{\partial_k}\zeta_A) = 0$ if $k \neq \mu_A$ and $K_{F_A}^{(0)}(\zeta_A, u\nabla_{\partial_{\mu_A}}\zeta_A) = 1$ by Lemma 4.2 and by the first statement of Lemma 4.3. $\square$

The functions ϕ, ψ and the homogenous change of coordinates in Lemma 5.2 should exist for any type A with $\chi_A > 0$ due to the existence of the primitive form by Theorem 3.1. We would, conversely, reconstruct the primitive form from the Gromov-Witten potential $\mathcal{F}_0^{\mathbb{P}^1 A}$ via the existence of them. Indeed, for a given A , they can be figured out by hand. We hope to give a proof of the existence of them for any A with $\chi_A > 0$ elsewhere in the near future.

In the present paper, we shall consider particular cases by giving all data explicitly in Appendix:

Theorem 5.4. *Let A be one of $(2, 3, 3)$, $(2, 3, 4)$, $(2, 3, 5)$. Then, the assumption in Lemma 5.2 is satisfied. Therefore, it turns out that $\zeta_A = [e^{-t_{\mu_A}} dx_1 \wedge dx_2 \wedge dx_3] \in \mathcal{H}_{F_A}^{(0)}$ is a primitive form for the tuple $(\mathcal{H}_{F_A}^{(0)}, \nabla, K_{F_A})$ with the minimal exponent $r = 1$.*

APPENDIX A. DATA FOR THE PROOF OF THEOREM 5.4

A.1. Frobenius potential for $A = (2, 3, 3)$.

$$\begin{aligned} & \mathcal{F}_{(2,3,3)}(t_1, t_{1,1}, t_{2,1}, t_{2,2}, t_{3,1}, t_{3,2}, t_{\mu_A}) \\ &= \frac{t_1^2 t_{\mu_A}}{2} + t_1 \left(\frac{t_{1,1}^2}{4} + \frac{t_{2,1} t_{2,2}}{3} + \frac{t_{3,1} t_{3,2}}{3} \right) + \frac{1}{18} (t_{2,1}^3 + t_{3,1}^3) - \left(\frac{t_{1,1}^4}{96} + \frac{1}{36} (t_{2,1}^2 t_{2,2}^2 + t_{3,1}^2 t_{3,2}^2) \right) \\ &+ \frac{1}{648} (t_{2,1} t_{2,2}^4 + t_{3,1} t_{3,2}^4) - \frac{1}{19440} (t_{2,2}^6 + t_{3,2}^6) + e^{t_{\mu_A}} \left(t_{1,1} t_{2,1} t_{3,1} + \frac{1}{6} t_{1,1} (t_{2,2}^2 t_{3,1} + t_{2,1} t_{3,2}^2) + \frac{1}{36} t_{1,1} t_{2,2}^2 t_{3,2}^2 \right) \\ &+ e^{2t_{\mu_A}} \left(\frac{1}{2} t_{1,1}^2 t_{2,2} t_{3,2} + \frac{1}{2} (t_{2,2} t_{3,1}^2 + t_{2,1}^2 t_{3,2}) + \frac{1}{6} (t_{2,1} t_{2,2}^2 t_{3,2} + t_{2,2} t_{3,1} t_{3,2}^2) + \frac{1}{72} (t_{2,2}^4 t_{3,2} + t_{2,2} t_{3,2}^4) \right) \\ &+ e^{3t_{\mu_A}} \left(\frac{t_{1,1}^3}{3} + t_{1,1} (t_{2,1} t_{2,2} + t_{3,1} t_{3,2}) + \frac{1}{6} t_{1,1} (t_{2,2}^3 + t_{1,1} t_{3,2}^3) \right) \\ &+ e^{4t_{\mu_A}} \left(t_{2,1} t_{3,1} + \frac{1}{6} (t_{2,2}^2 t_{3,1} + t_{2,1} t_{3,2}^2) + \frac{5 t_{2,2}^2 t_{3,2}^2}{18} \right) + e^{5t_{\mu_A}} t_{1,1} t_{2,2} t_{3,2} + e^{6t_{\mu_A}} \left(\frac{t_{1,1}^2}{2} + \frac{1}{6} (t_{2,2}^3 + t_{3,2}^3) \right) \\ &+ \frac{1}{2} e^{8t_{\mu_A}} t_{2,2} t_{3,2} + \frac{e^{12t_{\mu_A}}}{12}. \end{aligned}$$

A.2. Coordinate change for $A = (2, 3, 3)$.

$$\begin{aligned} s_1(\mathbf{t}, e^{t\mu_A}) &= t_1 - 2e^{2t\mu_A} t_{2,2} t_{3,2} - 6e^{3t\mu_A} t_{1,1} + 18e^{6t\mu_A}, \quad s_{1,1}(\mathbf{t}, e^{t\mu_A}) = t_{1,1} - 8e^{3t\mu_A}, \\ s_{2,1}(\mathbf{t}, e^{t\mu_A}) &= t_{2,1} + \frac{t_{2,2}^2}{6} - 3e^{2t\mu_A} t_{3,2}, \quad s_{2,2}(\mathbf{t}, e^{t\mu_A}) = t_{2,2}, \\ s_{3,1}(\mathbf{t}, e^{t\mu_A}) &= t_{3,1} + \frac{t_{3,2}^2}{6} - 3e^{2t\mu_A} t_{2,2}, \quad s_{3,2}(\mathbf{t}, e^{t\mu_A}) = t_{3,2}. \end{aligned}$$

A.3. ϕ for $A = (2, 3, 3)$.

$$\begin{aligned} \phi_{ij}^a &= 0, \quad a = 1, 2, 3, i = 1, j = 1, \dots, 7. \\ \phi_{22}^1 &= \frac{x_1}{2}, \quad \phi_{22}^2 = 0, \quad \phi_{22}^3 = 0. \quad \phi_{23}^1 = 0, \quad \phi_{23}^2 = 0, \quad \phi_{23}^3 = -e^{t\mu_A}. \\ \phi_{24}^1 &= e^{2t\mu_A} (-2t_{3,2} - 3x_3), \quad \phi_{24}^2 = -3e^{3t\mu_A}, \quad \phi_{24}^3 = e^{t\mu_A} \left(-\frac{t_{2,2}}{3} - x_2 \right). \\ \phi_{25}^1 &= 0, \quad \phi_{25}^2 = -e^{t\mu_A}, \quad \phi_{25}^3 = 0. \\ \phi_{26}^1 &= e^{2t\mu_A} (-2t_{2,2} - 3x_2), \quad \phi_{26}^2 = e^{t\mu_A} \left(-\frac{t_{3,2}}{3} - x_3 \right), \quad \phi_{26}^3 = -3e^{3t\mu_A}. \\ \phi_{27}^1 &= e^{2t\mu_A} (-4t_{2,2} t_{3,2} - 6t_{3,2} x_2 - 6t_{2,2} x_3 - 9x_2 x_3) + e^{3t\mu_A} (-9t_{1,1} - 12x_1) + 72e^{6t\mu_A}, \\ \phi_{27}^2 &= e^{t\mu_A} \left(-t_{3,1} - \frac{t_{3,2}^2}{6} - 2t_{3,2} x_3 - 3x_3^2 \right) - 3e^{3t\mu_A} t_{2,2}, \quad \phi_{27}^3 = -x_1 x_3 - 6e^{3t\mu_A} t_{3,2}. \\ \phi_{33}^1 &= 0, \quad \phi_{33}^2 = 0, \quad \phi_{33}^3 = 0. \quad \phi_{34}^1 = 3e^{3t\mu_A}, \quad \phi_{34}^2 = \frac{x_2}{3}, \quad \phi_{34}^3 = 0. \\ \phi_{35}^1 &= -e^{t\mu_A}, \quad \phi_{35}^2 = 0, \quad \phi_{35}^3 = 0. \quad \phi_{36}^1 = e^{t\mu_A} \left(-\frac{t_{3,2}}{3} - x_3 \right), \quad \phi_{36}^2 = -2e^{2t\mu_A}, \quad \phi_{36}^3 = 0. \\ \phi_{37}^1 &= -x_1 x_2 + e^{3t\mu_A} (-6t_{2,2} - 18x_2), \quad \phi_{37}^2 = e^{2t\mu_A} (-4t_{3,2} - 6x_3), \quad \phi_{37}^3 = e^{t\mu_A} (-t_{1,1} - 2x_1) - 4e^{4t\mu_A}. \\ \phi_{44}^1 &= -\frac{1}{3} x_1 x_2 - 2e^{3t\mu_A} t_{2,2}, \quad \phi_{44}^2 = \frac{x_2^2}{3} + e^{2t\mu_A} \left(-\frac{4t_{3,2}}{3} - 2x_3 \right), \quad \phi_{44}^3 = e^{t\mu_A} \left(-\frac{t_{1,1}}{3} - \frac{2x_1}{3} \right) + \frac{8}{3} e^{4t\mu_A}. \\ \phi_{45}^1 &= e^{t\mu_A} \left(-\frac{t_{2,2}}{3} - x_2 \right), \quad \phi_{45}^2 = 0, \quad \phi_{45}^3 = -2e^{2t\mu_A}. \\ \phi_{46}^1 &= e^{t\mu_A} \left(-\frac{1}{9} t_{2,2} t_{3,2} - \frac{1}{3} t_{3,2} x_2 - \frac{1}{3} t_{2,2} x_3 - x_2 x_3 \right) - e^{2t\mu_A} t_{1,1} - e^{5t\mu_A}, \\ \phi_{46}^2 &= e^{2t\mu_A} \left(-\frac{2t_{2,2}}{3} - x_2 \right), \quad \phi_{46}^3 = e^{2t\mu_A} \left(-\frac{2t_{3,2}}{3} - x_3 \right). \\ \phi_{47}^1 &= -\frac{1}{3} t_{2,2} x_1 x_2 - x_1 x_2^2 + e^{2t\mu_A} (-2t_{1,1} t_{3,2} - 3t_{1,1} x_3 - 3x_1 x_3) - 2e^{3t\mu_A} t_{2,2}^2 + e^{5t\mu_A} (10t_{3,2} + 42x_3), \\ \phi_{47}^2 &= e^{2t\mu_A} \left(-\frac{4}{3} t_{2,2} t_{3,2} - 2t_{3,2} x_2 - 2t_{2,2} x_3 \right) + e^{3t\mu_A} (-9t_{1,1} - 6x_1) + 36e^{6t\mu_A}, \\ \phi_{47}^3 &= e^{t\mu_A} \left(-\frac{1}{3} t_{1,1} t_{2,2} - \frac{2}{3} t_{2,2} x_1 - t_{1,1} x_2 - 2x_1 x_2 \right) + e^{2t\mu_A} \left(-2t_{3,1} - \frac{t_{3,2}^2}{3} \right) + e^{4t\mu_A} \left(\frac{14t_{2,2}}{3} + 14x_2 \right). \\ \phi_{55}^1 &= 0, \quad \phi_{55}^2 = 0, \quad \phi_{55}^3 = 0. \quad \phi_{56}^1 = 3e^{3t\mu_A}, \quad \phi_{56}^2 = 0, \quad \phi_{56}^3 = \frac{x_3}{3}. \\ \phi_{57}^1 &= -x_1 x_3 + e^{3t\mu_A} (-6t_{3,2} - 18x_3), \quad \phi_{57}^2 = e^{t\mu_A} (-t_{1,1} - 2x_1) - 4e^{4t\mu_A}, \quad \phi_{57}^3 = e^{2t\mu_A} (-4t_{2,2} - 6x_2). \\ \phi_{66}^1 &= -\frac{1}{3} x_1 x_3 - 2e^{3t\mu_A} t_{3,2}, \quad \phi_{66}^2 = e^{t\mu_A} \left(-\frac{t_{1,1}}{3} - \frac{2x_1}{3} \right) + \frac{8}{3} e^{4t\mu_A}, \quad \phi_{66}^3 = \frac{x_3^2}{3} + e^{2t\mu_A} \left(-\frac{4t_{2,2}}{3} - 2x_2 \right). \\ \phi_{67}^1 &= -\frac{1}{3} t_{3,2} x_1 x_3 - x_1 x_3^2 + e^{2t\mu_A} (-2t_{1,1} t_{2,2} - 3t_{1,1} x_2 - 3x_1 x_2) - 2e^{3t\mu_A} t_{3,2}^2 + e^{5t\mu_A} (10t_{2,2} + 42x_2), \\ \phi_{67}^2 &= e^{t\mu_A} \left(-\frac{1}{3} t_{1,1} t_{3,2} - \frac{2}{3} t_{3,2} x_1 - t_{1,1} x_3 - 2x_1 x_3 \right) + e^{2t\mu_A} \left(-2t_{2,1} - \frac{t_{2,2}^2}{3} \right) + e^{4t\mu_A} \left(\frac{14t_{3,2}}{3} + 14x_3 \right), \\ \phi_{67}^3 &= e^{2t\mu_A} \left(-\frac{4}{3} t_{2,2} t_{3,2} - 2t_{3,2} x_2 - 2t_{2,2} x_3 \right) + e^{3t\mu_A} (-9t_{1,1} - 6x_1) + 36e^{6t\mu_A}. \end{aligned}$$

$$\begin{aligned}
\phi_{77}^1 &= -e^{-t_{\mu A}} x_1^2 x_2 x_3 + e^{2t_{\mu A}} (-4t_{1,1} t_{2,2} t_{3,2} - 6t_{1,1} t_{3,2} x_2 - 12t_{3,2} x_1 x_2 - 6t_{1,1} t_{2,2} x_3 \\
&\quad - 12t_{2,2} x_1 x_3 - 9t_{1,1} x_2 x_3 - 54x_1 x_2 x_3) + e^{3t_{\mu A}} (-9t_{1,1}^2 + 24t_{2,2}^2 x_2 + 72t_{2,2} x_2^2 \\
&\quad + 54x_2^3 + 24t_{3,2}^2 x_3 + 72t_{3,2} x_3^2 + 54x_3^3) + e^{5t_{\mu A}} (8t_{2,2} t_{3,2} + 120t_{3,2} x_2 + 120t_{2,2} x_3 \\
&\quad + 180x_2 x_3) + e^{6t_{\mu A}} (252t_{1,1} + 288x_1) - 1440e^{9t_{\mu A}}, \\
\phi_{77}^2 &= e^{t_{\mu A}} \left(-t_{1,1} t_{3,1} - \frac{1}{6} t_{1,1} t_{3,2}^2 - 2t_{3,1} x_1 - \frac{1}{3} t_{3,2}^2 x_1 - 2t_{1,1} t_{3,2} x_3 - 4t_{3,2} x_1 x_3 \right. \\
&\quad \left. - 3t_{1,1} x_3^2 - 6x_1 x_3^2 \right) + e^{2t_{\mu A}} \left(-4t_{2,1} t_{3,2} - \frac{2}{3} t_{2,2}^2 t_{3,2} - 6t_{2,1} x_3 - t_{2,2}^2 x_3 \right. \\
&\quad \left. + 12t_{2,2} x_2 x_3 + 18x_2^2 x_3 \right) + e^{3t_{\mu A}} (-21t_{1,1} t_{2,2} - 18t_{2,2} x_1 - 18t_{1,1} x_2) \\
&\quad + e^{4t_{\mu A}} \left(-52t_{3,1} + \frac{10t_{3,2}^2}{3} - 14t_{3,2} x_3 - 48x_3^2 \right) + e^{6t_{\mu A}} (204t_{2,2} + 144x_2), \\
\phi_{77}^3 &= -t_{1,1} x_1 x_3 - 2x_1^2 x_3 + e^{2t_{\mu A}} \left(-4t_{2,2} t_{3,1} - \frac{2}{3} t_{2,2} t_{3,2}^2 - 6t_{3,1} x_2 - t_{3,2}^2 x_2 \right. \\
&\quad \left. + 12t_{3,2} x_2 x_3 + 18x_2 x_3^2 \right) + e^{3t_{\mu A}} (-24t_{1,1} t_{3,2} - 24t_{3,2} x_1 - 18t_{1,1} x_3 - 52x_1 x_3) \\
&\quad + e^{4t_{\mu A}} (12t_{2,2}^2 + 90t_{2,2} x_2 + 108x_2^2) + e^{6t_{\mu A}} (48t_{3,2} + 144x_3).
\end{aligned}$$

A.4. ψ for $A = (2, 3, 3)$.

$$\psi^1 = x_1 + 18e^{3t_{\mu A}}, \quad \psi^2 = 0, \quad \psi^3 = 0.$$

A.5. Frobenius potential for $A = (2, 3, 4)$.

$$\begin{aligned}
& \mathcal{F}_{(2,3,4)}(t_1, t_{1,1}, t_{2,1}, t_{2,2}, t_{3,1}, t_{3,2}, t_{3,3}, t_{\mu_A}) \\
&= \frac{t_1^2 t_{\mu_A}}{2} + t_1 \left(\frac{t_{1,1}^2}{4} + \frac{t_{2,1} t_{2,2}}{3} + \frac{t_{3,2}^2}{8} + \frac{t_{3,1} t_{3,3}}{4} \right) + \left(\frac{t_{2,1}^3}{18} + \frac{t_{3,1}^2 t_{3,2}}{8} \right) \\
&+ \left(-\frac{t_{1,1}^4}{96} - \frac{t_{2,1}^2 t_{2,2}^2}{36} - \frac{t_{3,2}^4}{192} - \frac{1}{32} t_{3,1} t_{3,2}^2 t_{3,3} - \frac{t_{3,1}^2 t_{3,3}^2}{64} \right) + \left(\frac{t_{2,1} t_{2,2}^4}{648} + \frac{t_{3,2}^3 t_{3,3}^2}{384} + \frac{1}{384} t_{3,1} t_{3,2} t_{3,3}^3 \right) \\
&+ \left(-\frac{t_{2,2}^6}{19440} - \frac{t_{3,2}^2 t_{3,3}^4}{3072} - \frac{t_{3,1} t_{3,3}^5}{30720} \right) + \frac{t_{3,2} t_{3,3}^6}{73728} - \frac{t_{3,3}^8}{4128768} + e^{t_{\mu_A}} \left(t_{1,1} t_{2,1} t_{3,1} + \frac{1}{6} t_{1,1} t_{2,2}^2 t_{3,1} + \frac{1}{4} t_{1,1} t_{2,1} t_{3,2} t_{3,3} \right. \\
&+ \frac{1}{24} t_{1,1} t_{2,2}^2 t_{3,2} t_{3,3} + \frac{1}{96} t_{1,1} t_{2,1} t_{3,3}^3 + \frac{1}{576} t_{1,1} t_{2,2}^2 t_{3,3}^3 \left. \right) + e^{2t_{\mu_A}} \left(\frac{t_{2,2} t_{3,1}^2}{2} + \frac{t_{2,1}^2 t_{3,2}}{2} + \frac{1}{2} t_{1,1}^2 t_{2,2} t_{3,2} \right. \\
&+ \frac{1}{6} t_{2,1} t_{2,2}^2 t_{3,2} + \frac{t_{2,2}^4 t_{3,2}}{72} + \frac{1}{4} t_{2,2} t_{3,1} t_{3,2} t_{3,3} + \frac{t_{2,1}^2 t_{3,3}^2}{8} + \frac{1}{8} t_{1,1}^2 t_{2,2} t_{3,3}^2 + \frac{1}{24} t_{2,1} t_{2,2}^2 t_{3,3}^2 + \frac{t_{2,2}^4 t_{3,3}^2}{288} \\
&+ \frac{1}{32} t_{2,2} t_{3,2}^2 t_{3,3}^2 + \frac{1}{96} t_{2,2} t_{3,1} t_{3,3}^3 + \frac{1}{384} t_{2,2} t_{3,2} t_{3,3}^4 + \frac{t_{2,2} t_{3,3}^6}{18432} \left. \right) + e^{3t_{\mu_A}} \left(t_{1,1} t_{3,1} t_{3,2} + \frac{t_{1,1}^3 t_{3,3}}{3} \right. \\
&+ t_{1,1} t_{2,1} t_{2,2} t_{3,3} + \frac{1}{6} t_{1,1} t_{2,2}^3 t_{3,3} + \frac{1}{4} t_{1,1} t_{3,2}^2 t_{3,3} + \frac{1}{4} t_{1,1} t_{3,1} t_{3,3}^2 + \frac{7}{96} t_{1,1} t_{3,2} t_{3,3}^3 + \frac{t_{1,1} t_{3,3}^5}{384} \left. \right) \\
&+ e^{4t_{\mu_A}} \left(t_{1,1}^2 t_{2,1} + \frac{t_{2,1}^2 t_{2,2}}{2} + \frac{2t_{1,1}^2 t_{2,2}^2}{3} + \frac{t_{2,1} t_{2,2}^3}{6} + \frac{t_{2,2}^5}{72} + \frac{t_{2,2}^2 t_{3,2}^2}{4} + t_{2,1} t_{3,1} t_{3,3} + \frac{1}{6} t_{2,2}^2 t_{3,1} t_{3,3} \right. \\
&+ \frac{1}{4} t_{2,1} t_{3,2} t_{3,3}^2 + \frac{1}{6} t_{2,2}^2 t_{3,2} t_{3,3}^2 + \frac{t_{2,1} t_{3,3}^4}{96} + \frac{5t_{2,2}^2 t_{3,3}^4}{288} \left. \right) + e^{5t_{\mu_A}} \left(t_{1,1} t_{2,2} t_{3,1} + \frac{5}{4} t_{1,1} t_{2,2} t_{3,2} t_{3,3} + \frac{25}{96} t_{1,1} t_{2,2} t_{3,3}^3 \right. \\
&+ e^{6t_{\mu_A}} \left(\frac{t_{3,1}^2}{2} + \frac{t_{1,1}^2 t_{3,2}}{2} + t_{2,1} t_{2,2} t_{3,2} + \frac{t_{2,2}^3 t_{3,2}}{6} + \frac{t_{3,2}^3}{6} + \frac{t_{3,1} t_{3,2} t_{3,3}}{4} + \frac{5t_{1,1}^2 t_{3,3}^2}{8} + \frac{1}{4} t_{2,1} t_{2,2} t_{3,3}^2 \right. \\
&+ \frac{5t_{2,2}^3 t_{3,3}^2}{24} + \frac{5t_{3,2}^2 t_{3,3}^2}{32} + \frac{t_{3,1} t_{3,3}^3}{96} + \frac{13t_{3,2} t_{3,3}^4}{384} + \frac{49t_{3,3}^6}{18432} \left. \right) + e^{7t_{\mu_A}} \left(t_{1,1} t_{2,1} t_{3,3} + \frac{7}{6} t_{1,1} t_{2,2}^2 t_{3,3} \right) \\
&+ e^{8t_{\mu_A}} \left(\frac{t_{2,1}^2}{2} + t_{1,1}^2 t_{2,2} + \frac{t_{2,1} t_{2,2}^2}{6} + \frac{5t_{2,2}^4}{36} + \frac{1}{2} t_{2,2} t_{3,2} t_{3,3}^2 + \frac{t_{2,2} t_{3,3}^4}{8} \right) + e^{9t_{\mu_A}} \left(t_{1,1} t_{3,2} t_{3,3} + \frac{t_{1,1} t_{3,3}^3}{4} \right) \\
&+ e^{10t_{\mu_A}} \left(\frac{t_{2,2}^2 t_{3,2}}{2} + \frac{5t_{2,2}^2 t_{3,3}^2}{8} \right) + e^{11t_{\mu_A}} t_{1,1} t_{2,2} t_{3,3} + e^{12t_{\mu_A}} \left(\frac{t_{1,1}^2}{2} + \frac{t_{2,2}^3}{6} + \frac{t_{3,2}^2}{4} + \frac{t_{3,2} t_{3,3}^2}{8} + \frac{19t_{3,3}^4}{192} \right) \\
&+ \frac{1}{2} e^{14t_{\mu_A}} t_{2,2} t_{3,3}^2 + \frac{1}{4} e^{16t_{\mu_A}} t_{2,2}^2 + \frac{1}{6} e^{18t_{\mu_A}} t_{3,3}^2 + \frac{e^{24t_{\mu_A}}}{24}.
\end{aligned}$$

A.6. Coordinate change for $A = (2, 3, 4)$.

$$\begin{aligned}
s_1(\mathbf{t}) &= t_1 + e^{2t_{\mu_A}} \left(-2t_{2,2} t_{3,2} - \frac{1}{2} t_{2,2} t_{3,3}^2 \right) - 6e^{3t_{\mu_A}} t_{1,1} t_{3,3} - 12e^{4t_{\mu_A}} t_{2,1} \\
&+ e^{6t_{\mu_A}} \left(30t_{3,2} + \frac{51t_{3,3}^2}{2} \right) + 52e^{8t_{\mu_A}} t_{2,2} - 702e^{12t_{\mu_A}}, \\
s_{1,1}(\mathbf{t}) &= t_{1,1} - 8e^{3t_{\mu_A}} t_{3,3}, \\
s_{2,1}(\mathbf{t}) &= t_{2,1} + \frac{t_{2,2}^2}{6} + e^{2t_{\mu_A}} \left(-3t_{3,2} - \frac{3t_{3,3}^2}{4} \right) - 15e^{4t_{\mu_A}} t_{2,2} + 240e^{8t_{\mu_A}}, \quad s_{2,2}(\mathbf{t}) = t_{2,2} - 27e^{4t_{\mu_A}}, \\
s_{3,1}(\mathbf{t}) &= t_{3,1} + \frac{1}{4} t_{3,2} t_{3,3} + \frac{t_{3,3}^3}{96} - 3e^{2t_{\mu_A}} t_{2,2} t_{3,3} - 8e^{3t_{\mu_A}} t_{1,1} + 45e^{6t_{\mu_A}} t_{3,3}, \\
s_{3,2}(\mathbf{t}) &= t_{3,2} + \frac{t_{3,3}^2}{4} - 4e^{2t_{\mu_A}} t_{2,2} + 28e^{6t_{\mu_A}}, \quad s_{3,3}(\mathbf{t}) = t_{3,3}.
\end{aligned}$$

A.7. ϕ for $A = (2, 3, 4)$.

$$\begin{aligned}
\phi_{ij}^a &= 0, \quad a = 1, 2, 3, i = 1, j = 1, \dots, 8. \\
\phi_{22}^1 &= \frac{x_1}{2}, \quad \phi_{22}^2 = 16e^{4t_{\mu_A}}, \quad \phi_{22}^3 = 0, \quad \phi_{23}^1 = 8e^{4t_{\mu_A}}, \quad \phi_{23}^2 = 0, \quad \phi_{23}^3 = -e^{t_{\mu_A}}.
\end{aligned}$$

$$\begin{aligned}
\phi_{24}^1 &= e^{2t\mu_A} \left(-2t_{3,2} - \frac{t_{3,3}^2}{2} - 3t_{3,3}x_3 - 4x_3^2 \right) + e^{4t\mu_A} \left(\frac{8t_{2,2}}{3} - 4x_2 \right) + 40e^{8t\mu_A}, \\
\phi_{24}^2 &= e^{3t\mu_A} (-3t_{3,3} - 4x_3), \quad \phi_{24}^3 = e^{t\mu_A} \left(-\frac{t_{2,2}}{3} - x_2 \right) + 7e^{5t\mu_A}, \\
\phi_{25}^1 &= 0, \quad \phi_{25}^2 = -e^{t\mu_A}, \quad \phi_{25}^3 = 0. \\
\phi_{26}^1 &= e^{2t\mu_A} (-2t_{2,2} - 3x_2) + 30e^{6t\mu_A}, \quad \phi_{26}^2 = e^{t\mu_A} \left(-\frac{t_{3,3}}{4} - x_3 \right), \quad \phi_{26}^3 = -3e^{3t\mu_A}. \\
\phi_{27}^1 &= e^{2t\mu_A} \left(-t_{2,2}t_{3,3} - \frac{3}{2}t_{3,3}x_2 - 2t_{2,2}x_3 - 3x_2x_3 \right) + e^{3t\mu_A} (-3t_{1,1} - 4x_1) + e^{6t\mu_A} (39t_{3,3} + 54x_3), \\
\phi_{27}^2 &= e^{t\mu_A} \left(-\frac{t_{3,2}}{4} - \frac{t_{3,3}^2}{32} - \frac{1}{2}t_{3,3}x_3 - x_3^2 \right) - e^{3t\mu_A} t_{2,2} + -e^{7t\mu_A}, \quad \phi_{27}^3 = e^{3t\mu_A} \left(-\frac{3t_{3,3}}{2} - 2x_3 \right). \\
\phi_{28}^1 &= e^{2t\mu_A} \left(-4t_{2,2}t_{3,2} - t_{2,2}t_{3,3}^2 - 6t_{3,2}x_2 - \frac{3}{2}t_{3,3}^2x_2 - 6t_{2,2}t_{3,3}x_3 - 9t_{3,3}x_2x_3 \right. \\
&\quad \left. - 8t_{2,2}x_3^2 - 12x_2x_3^2 \right) + e^{3t\mu_A} (-9t_{1,1}t_{3,3} - 12t_{3,3}x_1 - 12t_{1,1}x_3 - 16x_1x_3) \\
&\quad + e^{6t\mu_A} (84t_{3,2} + 93t_{3,3}^2 + 294t_{3,3}x_3 + 264x_3^2) + e^{8t\mu_A} (32t_{2,2} + 48x_2) - 384e^{12t\mu_A}, \\
\phi_{28}^2 &= e^{t\mu_A} \left(-t_{3,1} - \frac{1}{4}t_{3,2}t_{3,3} - \frac{t_{3,3}^3}{96} - 2t_{3,2}x_3 - \frac{1}{2}t_{3,3}^2x_3 - 3t_{3,3}x_3^2 - 4x_3^3 \right) \\
&\quad - 3e^{3t\mu_A} t_{2,2}t_{3,3} + e^{4t\mu_A} (-8t_{1,1} - 32x_1) + e^{7t\mu_A} (209t_{3,3} + 304x_3), \\
\phi_{28}^3 &= -x_1x_3 + e^{3t\mu_A} \left(-6t_{3,2} - \frac{3t_{3,3}^2}{2} \right) + e^{5t\mu_A} (-4t_{2,2} + 12x_2) - 96e^{9t\mu_A}. \\
\phi_{33}^1 &= 0, \quad \phi_{33}^2 = 0, \quad \phi_{33}^3 = 0. \quad \phi_{34}^1 = e^{3t\mu_A} (3t_{3,3} + 4x_3), \quad \phi_{34}^2 = \frac{x_2}{3} + 8e^{4t\mu_A}, \quad \phi_{34}^3 = 0. \\
\phi_{35}^1 &= -e^{t\mu_A}, \quad \phi_{35}^2 = 0, \quad \phi_{35}^3 = 0. \quad \phi_{36}^1 = e^{t\mu_A} \left(-\frac{t_{3,3}}{4} - x_3 \right), \quad \phi_{36}^2 = -2e^{2t\mu_A}, \quad \phi_{36}^3 = 0. \\
\phi_{37}^1 &= e^{t\mu_A} \left(-\frac{t_{3,2}}{4} - \frac{t_{3,3}^2}{32} - \frac{1}{2}t_{3,3}x_3 - x_3^2 \right) + e^{3t\mu_A} (-t_{2,2} - 6x_2) + 63e^{7t\mu_A}, \\
\phi_{37}^2 &= e^{2t\mu_A} (-t_{3,3} - 2x_3), \quad \phi_{37}^3 = -4e^{4t\mu_A}. \\
\phi_{38}^1 &= -x_1x_2 + e^{3t\mu_A} (-6t_{2,2}t_{3,3} - 18t_{3,3}x_2 - 8t_{2,2}x_3 - 24x_2x_3) + e^{7t\mu_A} (246t_{3,3} + 264x_3), \\
\phi_{38}^2 &= e^{2t\mu_A} (-4t_{3,2} - t_{3,3}^2 - 6t_{3,3}x_3 - 8x_3^2) - 36e^{4t\mu_A} x_2 + 416e^{8t\mu_A}, \\
\phi_{38}^3 &= e^{t\mu_A} (-t_{1,1} - 2x_1) - 4e^{4t\mu_A} t_{3,3}. \\
\phi_{44}^1 &= -\frac{1}{3}x_1x_2 + e^{3t\mu_A} \left(-2t_{2,2}t_{3,3} - \frac{8}{3}t_{2,2}x_3 \right) + e^{7t\mu_A} (18t_{3,3} + 24x_3), \\
\phi_{44}^2 &= \frac{x_2^2}{3} + e^{2t\mu_A} \left(-\frac{4t_{3,2}}{3} - \frac{t_{3,3}^2}{3} - 2t_{3,3}x_3 - \frac{8x_3^2}{3} \right) - 4e^{4t\mu_A} x_2 + \frac{32}{3}e^{8t\mu_A}, \\
\phi_{44}^3 &= e^{t\mu_A} \left(-\frac{t_{1,1}}{3} - \frac{2x_1}{3} \right) + e^{4t\mu_A} \left(\frac{8t_{3,3}}{3} + 4x_3 \right). \\
\phi_{45}^1 &= e^{t\mu_A} \left(-\frac{t_{2,2}}{3} - x_2 \right) + 15e^{5t\mu_A}, \quad \phi_{45}^2 = 0, \quad \phi_{45}^3 = -2e^{2t\mu_A}. \\
\phi_{46}^1 &= e^{t\mu_A} \left(-\frac{1}{12}t_{2,2}t_{3,3} - \frac{1}{4}t_{3,3}x_2 - \frac{1}{3}t_{2,2}x_3 - x_2x_3 \right) - e^{2t\mu_A} t_{1,1} + e^{5t\mu_A} \left(\frac{11t_{3,3}}{4} + 3x_3 \right), \\
\phi_{46}^2 &= e^{2t\mu_A} \left(-\frac{2t_{2,2}}{3} - x_2 \right) + 6e^{6t\mu_A}, \quad \phi_{46}^3 = e^{2t\mu_A} \left(-\frac{t_{3,3}}{2} - x_3 \right). \\
\phi_{47}^1 &= e^{t\mu_A} \left(-\frac{1}{12}t_{2,2}t_{3,2} - \frac{1}{96}t_{2,2}t_{3,3}^2 - \frac{1}{4}t_{3,2}x_2 - \frac{1}{32}t_{3,3}^2x_2 - \frac{1}{6}t_{2,2}t_{3,3}x_3 - \frac{1}{2}t_{3,3}x_2x_3 - \frac{1}{3}t_{2,2}x_3^2 - x_2x_3^2 \right) \\
&\quad + e^{3t\mu_A} \left(-\frac{1}{3}t_{2,2}^2 + t_{2,2}x_2 \right) + e^{5t\mu_A} \left(\frac{31t_{3,2}}{4} + \frac{31t_{3,3}^2}{32} + \frac{31}{2}t_{3,3}x_3 + 23x_3^2 \right) + e^{7t\mu_A} (-12t_{2,2} - 3x_2) + 31e^{11t\mu_A}, \\
\phi_{47}^2 &= e^{2t\mu_A} \left(-\frac{1}{3}t_{2,2}t_{3,3} - \frac{1}{2}t_{3,3}x_2 - \frac{2}{3}t_{2,2}x_3 \right) + e^{3t\mu_A} (-3t_{1,1} - 2x_1) + e^{6t\mu_A} (15t_{3,3} + 14x_3),
\end{aligned}$$

$$\begin{aligned}
\phi_{47}^3 &= e^{2t_{\mu_A}} \left(-\frac{t_{3,2}}{2} - \frac{t_{3,3}^2}{16} - \frac{1}{2}t_{3,3}x_3 - x_3^2 \right) + e^{4t_{\mu_A}} \left(\frac{2t_{2,2}}{3} + 2x_2 \right) - 18e^{8t_{\mu_A}}. \\
\phi_{48}^1 &= -\frac{1}{3}t_{2,2}x_1x_2 - x_1x_2^2 + e^{2t_{\mu_A}} \left(-2t_{1,1}t_{3,2} - \frac{1}{2}t_{1,1}t_{3,3}^2 - 3t_{1,1}t_{3,3}x_3 - 3t_{3,3}x_1x_3 - 4t_{1,1}x_3^2 - 4x_1x_3^2 \right) \\
&\quad + e^{3t_{\mu_A}} \left(-2t_{2,2}^2t_{3,3} - \frac{8}{3}t_{2,2}^2x_3 \right) + e^{4t_{\mu_A}} (-8t_{1,1}t_{2,2} - 12t_{1,1}x_2 + 27x_1x_2) \\
&\quad + e^{5t_{\mu_A}} \left(10t_{3,2}t_{3,3} + \frac{5t_{3,3}^3}{2} - 24t_{3,2}x_3 + 36t_{3,3}^2x_3 + 56t_{3,3}x_3^2 \right) \\
&\quad + e^{7t_{\mu_A}} (56t_{2,2}t_{3,3} + 36t_{3,3}x_2 + 160t_{2,2}x_3 + 48x_2x_3) + 280e^{8t_{\mu_A}}t_{1,1} + e^{11t_{\mu_A}}(-1130t_{3,3} - 1464x_3), \\
\phi_{48}^2 &= e^{2t_{\mu_A}} \left(-\frac{4}{3}t_{2,2}t_{3,2} - \frac{1}{3}t_{2,2}t_{3,3}^2 - 2t_{3,2}x_2 - \frac{1}{2}t_{3,3}^2x_2 - 2t_{2,2}t_{3,3}x_3 - \frac{8}{3}t_{2,2}x_3^2 \right) \\
&\quad + e^{3t_{\mu_A}} (-9t_{1,1}t_{3,3} - 6t_{3,3}x_1 - 12t_{1,1}x_3 - 8x_1x_3) + e^{4t_{\mu_A}} \left(-8t_{2,1} - \frac{4t_{2,2}^2}{3} - 8t_{2,2}x_2 - 36x_2^2 \right) \\
&\quad + e^{6t_{\mu_A}} (84t_{3,2} + 57t_{3,3}^2 + 210t_{3,3}x_3 + 216x_3^2) + e^{8t_{\mu_A}} \left(-\frac{88t_{2,2}}{3} + 532x_2 \right) - 1824e^{12t_{\mu_A}}, \\
\phi_{48}^3 &= e^{t_{\mu_A}} \left(-\frac{1}{3}t_{1,1}t_{2,2} - \frac{2}{3}t_{2,2}x_1 - t_{1,1}x_2 - 2x_1x_2 \right) + e^{2t_{\mu_A}} \left(-2t_{3,1} - \frac{1}{2}t_{3,2}t_{3,3} - \frac{t_{3,3}^3}{48} \right) \\
&\quad + e^{4t_{\mu_A}} \left(\frac{14}{3}t_{2,2}t_{3,3} + 14t_{3,3}x_2 + 8t_{2,2}x_3 \right) + e^{5t_{\mu_A}} (19t_{1,1} + 54x_1) + e^{8t_{\mu_A}} (-270t_{3,3} - 168x_3). \\
\phi_{55}^1 &= 0, \quad \phi_{55}^2 = 0, \quad \phi_{55}^3 = 0. \quad \phi_{56}^1 = 3e^{3t_{\mu_A}}, \quad \phi_{56}^2 = 0, \quad \phi_{56}^3 = 0. \\
\phi_{57}^1 &= \frac{3}{2}e^{3t_{\mu_A}}t_{3,3}, \quad \phi_{57}^2 = 8e^{4t_{\mu_A}}, \quad \phi_{57}^3 = \frac{x_3}{4}. \\
\phi_{58}^1 &= -x_1x_3 + e^{3t_{\mu_A}} \left(-6t_{3,2} - \frac{3t_{3,3}^2}{2} - 18t_{3,3}x_3 - 24x_3^2 \right) + e^{5t_{\mu_A}} (12t_{2,2} - 36x_2) + 336e^{9t_{\mu_A}}, \\
\phi_{58}^2 &= e^{t_{\mu_A}} (-t_{1,1} - 2x_1) + e^{4t_{\mu_A}} (-4t_{3,3} - 48x_3), \\
\phi_{58}^3 &= e^{2t_{\mu_A}} (-4t_{2,2} - 6x_2) + 36e^{6t_{\mu_A}}. \\
\phi_{66}^1 &= e^{3t_{\mu_A}} \left(\frac{3t_{3,3}}{2} + 6x_3 \right), \quad \phi_{66}^2 = 12e^{4t_{\mu_A}}, \quad \phi_{66}^3 = \frac{x_3}{4}. \\
\phi_{67}^1 &= -\frac{1}{4}x_1x_3 + e^{3t_{\mu_A}} \left(-\frac{9t_{3,2}}{4} - \frac{9t_{3,3}^2}{32} - \frac{3}{2}t_{3,3}x_3 - 3x_3^2 \right) + e^{5t_{\mu_A}} (7t_{2,2} + 6x_2) - 57e^{9t_{\mu_A}}, \\
\phi_{67}^2 &= e^{t_{\mu_A}} \left(-\frac{t_{1,1}}{4} - \frac{x_1}{2} \right) + e^{4t_{\mu_A}} (t_{3,3} + 2x_3), \\
\phi_{67}^3 &= \frac{x_3^2}{4} + e^{2t_{\mu_A}} \left(-t_{2,2} - \frac{3x_2}{2} \right) + 15e^{6t_{\mu_A}}. \\
\phi_{68}^1 &= -\frac{1}{4}t_{3,3}x_1x_3 - x_1x_3^2 + e^{2t_{\mu_A}} (-2t_{1,1}t_{2,2} - 3t_{1,1}x_2 - 3x_1x_2) \\
&\quad + e^{3t_{\mu_A}} \left(-\frac{3}{2}t_{3,2}t_{3,3} - \frac{3t_{3,3}^3}{8} + 6t_{3,2}x_3 - 3t_{3,3}^2x_3 - 6t_{3,3}x_3^2 \right) \\
&\quad + e^{5t_{\mu_A}} (13t_{2,2}t_{3,3} + 33t_{3,3}x_2 - 12t_{2,2}x_3 + 36x_2x_3) + 18e^{6t_{\mu_A}}t_{1,1} + e^{9t_{\mu_A}}(-150t_{3,3} - 120x_3), \\
\phi_{68}^2 &= e^{t_{\mu_A}} \left(-\frac{1}{4}t_{1,1}t_{3,3} - \frac{1}{2}t_{3,3}x_1 - t_{1,1}x_3 - 2x_1x_3 \right) + e^{2t_{\mu_A}} \left(-2t_{2,1} - \frac{t_{2,2}^2}{3} \right) \\
&\quad + e^{4t_{\mu_A}} \left(6t_{3,2} + \frac{t_{3,3}^2}{2} + 2t_{3,3}x_3 - 24x_3^2 \right) + e^{6t_{\mu_A}} (54t_{2,2} + 108x_2) - 1056e^{10t_{\mu_A}}, \\
\phi_{68}^3 &= e^{2t_{\mu_A}} \left(-t_{2,2}t_{3,3} - \frac{3}{2}t_{3,3}x_2 - 2t_{2,2}x_3 \right) + e^{3t_{\mu_A}} (-9t_{1,1} - 6x_1) + e^{6t_{\mu_A}} (45t_{3,3} + 42x_3).
\end{aligned}$$

$$\begin{aligned}
\phi_{77}^1 &= -\frac{1}{16}t_{3,3}x_1x_3 - \frac{1}{4}x_1x_3^2 + e^{2t_{\mu A}} \left(-\frac{1}{2}t_{1,1}t_{2,2} - \frac{3}{4}t_{1,1}x_2 - \frac{3}{2}x_1x_2 \right) + e^{3t_{\mu A}} \left(-\frac{3}{32}t_{3,3}^3 + \frac{3}{8}t_{3,3}^2x_3 + \frac{3}{2}t_{3,3}x_3^2 \right) \\
&\quad + e^{5t_{\mu A}} (2t_{2,2}t_{3,3} + 12t_{3,3}x_2 + 8t_{2,2}x_3 + 12x_2x_3) + e^{6t_{\mu A}} \left(\frac{51t_{1,1}}{2} + 32x_1 \right) + e^{9t_{\mu A}} (-198t_{3,3} - 216x_3), \\
\phi_{77}^2 &= e^{t_{\mu A}} \left(-\frac{1}{16}t_{1,1}t_{3,3} - \frac{1}{8}t_{3,3}x_1 - \frac{1}{4}t_{1,1}x_3 - \frac{1}{2}x_1x_3 \right) + e^{2t_{\mu A}} \left(-\frac{t_{2,1}}{2} - \frac{t_{2,2}^2}{12} \right) \\
&\quad + e^{4t_{\mu A}} \left(-\frac{t_{3,2}}{2} + \frac{5t_{3,3}^2}{8} + 4t_{3,3}x_3 + 4x_3^2 \right) - \frac{1}{2}e^{6t_{\mu A}} t_{2,2}, \\
\phi_{77}^3 &= \frac{1}{16}t_{3,3}x_3^2 + \frac{x_3^3}{4} + e^{2t_{\mu A}} \left(-\frac{1}{4}t_{2,2}t_{3,3} - \frac{3}{8}t_{3,3}x_2 - t_{2,2}x_3 \right) + e^{3t_{\mu A}} (-3t_{1,1} - 3x_1) + e^{6t_{\mu A}} \left(\frac{75t_{3,3}}{4} + 19x_3 \right). \\
\phi_{78}^1 &= -\frac{1}{4}t_{3,2}x_1x_3 - \frac{1}{32}t_{3,3}^2x_1x_3 - \frac{1}{2}t_{3,3}x_1x_3^2 - x_1x_3^3 + e^{2t_{\mu A}} \left(-t_{1,1}t_{2,2}t_{3,3} - \frac{3}{2}t_{1,1}t_{3,3}x_2 - \frac{3}{2}t_{3,3}x_1x_2 \right. \\
&\quad \left. - 2t_{1,1}t_{2,2}x_3 - 3t_{2,2}x_1x_3 - 3t_{1,1}x_2x_3 - 12x_1x_2x_3 \right) + e^{3t_{\mu A}} \left(-3t_{1,1}^2 - \frac{3t_{3,2}^2}{2} - \frac{9}{16}t_{3,2}t_{3,3}^2 \right. \\
&\quad \left. - \frac{3t_{3,3}^4}{64} + 8t_{2,2}^2x_2 + 24t_{2,2}x_2^2 + 18x_2^3 - \frac{3}{2}t_{3,2}t_{3,3}x_3 + \frac{3}{16}t_{3,3}^3x_3 + \frac{3}{4}t_{3,3}^2x_3^2 \right) \\
&\quad + e^{5t_{\mu A}} \left(5t_{2,2}t_{3,2} + \frac{47}{8}t_{2,2}t_{3,3}^2 + 3t_{3,2}x_2 + \frac{183}{8}t_{3,3}^2x_2 + 52t_{2,2}t_{3,3}x_3 + 78t_{3,3}x_2x_3 + 52t_{2,2}x_3^2 + 60x_2x_3^2 \right) \\
&\quad + e^{6t_{\mu A}} (93t_{1,1}t_{3,3} + 96t_{3,3}x_1 + 114t_{1,1}x_3 + 249x_1x_3) + e^{7t_{\mu A}} (-84t_{2,2}^2 - 540t_{2,2}x_2 - 648x_2^2) \\
&\quad + e^{9t_{\mu A}} (102t_{3,2} - 582t_{3,3}^2 - 1674t_{3,3}x_3 - 1128x_3^2) + e^{11t_{\mu A}} (2780t_{2,2} + 7284x_2) - 26064e^{15t_{\mu A}}, \\
\phi_{78}^2 &= e^{t_{\mu A}} \left(-\frac{1}{4}t_{1,1}t_{3,2} - \frac{1}{32}t_{1,1}t_{3,3}^2 - \frac{1}{2}t_{3,2}x_1 - \frac{1}{16}t_{3,3}^2x_1 - \frac{1}{2}t_{1,1}t_{3,3}x_3 - t_{3,3}x_1x_3 - t_{1,1}x_3^2 - 2x_1x_3^2 \right) \\
&\quad + e^{2t_{\mu A}} \left(-t_{2,1}t_{3,3} - \frac{1}{6}t_{2,2}^2t_{3,3} - 2t_{2,1}x_3 - \frac{1}{3}t_{2,2}^2x_3 + 4t_{2,2}x_2x_3 + 6x_2^2x_3 \right) + e^{3t_{\mu A}} (-7t_{1,1}t_{2,2} - 6t_{2,2}x_1 - 6t_{1,1}x_2) \\
&\quad + e^{4t_{\mu A}} \left(-16t_{3,1} - 2t_{3,2}t_{3,3} + \frac{11t_{3,3}^3}{24} - 26t_{3,2}x_3 + 2t_{3,3}^2x_3 - 28t_{3,3}x_3^2 - 64x_3^3 \right) \\
&\quad + e^{6t_{\mu A}} (95t_{2,2}t_{3,3} + 102t_{3,3}x_2 + 198t_{2,2}x_3 - 108x_2x_3) + e^{7t_{\mu A}} (413t_{1,1} + 498x_1) + e^{10t_{\mu A}} (-3228t_{3,3} - 2288x_3), \\
\phi_{78}^3 &= e^{2t_{\mu A}} \left(-t_{2,2}t_{3,2} - \frac{1}{8}t_{2,2}t_{3,3}^2 - \frac{3}{2}t_{3,2}x_2 - \frac{3}{16}t_{3,3}^2x_2 - t_{2,2}t_{3,3}x_3 - 2t_{2,2}x_3^2 \right) \\
&\quad + e^{3t_{\mu A}} \left(-\frac{9}{2}t_{1,1}t_{3,3} - 3t_{3,3}x_1 - 6t_{1,1}x_3 - 16x_1x_3 \right) + e^{4t_{\mu A}} (4t_{2,2}^2 + 30t_{2,2}x_2 + 36x_2^2) \\
&\quad + e^{6t_{\mu A}} \left(-15t_{3,2} + \frac{105t_{3,3}^2}{8} + 57t_{3,3}x_3 + 42x_3^2 \right) + e^{8t_{\mu A}} (-240t_{2,2} - 666x_2) + 3036e^{12t_{\mu A}}.
\end{aligned}$$

$$\begin{aligned}
\phi_{88}^1 = & -e^{-t_{\mu A}} x_1^2 x_2 x_3 + e^{2t_{\mu A}} \left(-4t_{1,1} t_{2,2} t_{3,2} - t_{1,1} t_{2,2} t_{3,3}^2 - 6t_{1,1} t_{3,2} x_2 - \frac{3}{2} t_{1,1} t_{3,3}^2 x_2 - 12t_{3,2} x_1 x_2 - 3t_{3,3}^2 x_1 x_2 \right. \\
& - 6t_{1,1} t_{2,2} t_{3,3} x_3 - 12t_{2,2} t_{3,3} x_1 x_3 - 9t_{1,1} t_{3,3} x_2 x_3 - 54t_{3,3} x_1 x_2 x_3 - 8t_{1,1} t_{2,2} x_3^2 - 16t_{2,2} x_1 x_3^2 - 12t_{1,1} x_2 x_3^2 \\
& \left. - 72x_1 x_2 x_3^2 \right) + e^{3t_{\mu A}} \left(-9t_{1,1}^2 t_{3,3} + 24t_{2,2}^2 t_{3,3} x_2 + 72t_{2,2} t_{3,3} x_2^2 + 54t_{3,3} x_2^3 - 12t_{1,1}^2 x_3 + 24t_{3,3}^2 x_3 \right. \\
& + 12t_{3,2} t_{3,3}^2 x_3 + \frac{3}{2} t_{3,3}^4 x_3 - 24t_{1,1} x_1 x_3 + 32t_{2,2}^2 x_2 x_3 + 96t_{2,2} x_2^2 x_3 + 72x_2^3 x_3 + 72t_{3,2} t_{3,3} x_3^2 \\
& + 18t_{3,3}^3 x_3^2 + 96t_{3,2} x_3^3 + 78t_{3,3}^2 x_3^3 + 144t_{3,3} x_3^4 + 96x_3^5 \Big) + e^{5t_{\mu A}} \left(8t_{2,2} t_{3,2} t_{3,3} + 2t_{2,2} t_{3,3}^3 + 120t_{3,2} t_{3,3} x_2 + 30t_{3,3}^3 x_2 \right. \\
& - 96t_{2,2} t_{3,2} x_3 + 96t_{2,2} t_{3,3}^2 x_3 + 288t_{3,2} x_2 x_3 + 252t_{3,3}^2 x_2 x_3 + 160t_{2,2} t_{3,3} x_3^2 + 672t_{3,3} x_2 x_3^2 + 576x_2 x_3^3 \Big) \\
& + e^{4t_{\mu A}} \left(-16t_{1,1} t_{2,2}^2 + 24t_{1,1} t_{2,2} x_2 + 24t_{2,2} x_1 x_2 + 72t_{1,1} x_2^2 - 72x_1 x_2^2 \right) \\
& + e^{6t_{\mu A}} \left(-36t_{1,1} t_{3,2} + 243t_{1,1} t_{3,3}^2 + 288t_{3,3}^2 x_1 + 714t_{1,1} t_{3,3} x_3 + 1404t_{3,3} x_1 x_3 + 504t_{1,1} x_3^2 + 1104x_1 x_3^2 \right) \\
& + e^{7t_{\mu A}} \left(-304t_{2,2}^2 t_{3,3} - 2352t_{2,2} t_{3,3} x_2 - 2520t_{3,3} x_2^2 - 192t_{2,2}^2 x_3 - 2880t_{2,2} x_2 x_3 - 2592x_2^2 x_3 \right) \\
& + e^{8t_{\mu A}} \left(224t_{1,1} t_{2,2} - 1608t_{1,1} x_2 + 816x_1 x_2 \right) + e^{9t_{\mu A}} \left(216t_{3,2} t_{3,3} - 1386t_{3,3}^3 - 1824t_{3,2} x_3 - 6912t_{3,3}^2 x_3 \right. \\
& - 11808t_{3,3} x_3^2 - 7680x_3^3 \Big) + e^{11t_{\mu A}} \left(13808t_{2,2} t_{3,3} + 35400t_{3,3} x_2 + 17472t_{2,2} x_3 \right. \\
& \left. + 33120x_2 x_3 \right) + 10224e^{12t_{\mu A}} t_{1,1} + e^{15t_{\mu A}} (-160704t_{3,3} - 149376x_3),
\end{aligned}$$

$$\begin{aligned}
\phi_{88}^2 = & -t_{1,1} x_1 x_2 - 2x_1^2 x_2 + e^{2t_{\mu A}} \left(-4t_{2,1} t_{3,2} - \frac{2}{3} t_{2,2}^2 t_{3,2} - t_{2,1} t_{3,3}^2 - \frac{1}{6} t_{2,2}^2 t_{3,3}^2 - 6t_{2,1} t_{3,3} x_3 - t_{2,2}^2 t_{3,3} x_3 \right. \\
& + 12t_{2,2} t_{3,3} x_2 x_3 + 18t_{3,3} x_2^2 x_3 - 8t_{2,1} x_3^2 - \frac{4}{3} t_{2,2}^2 x_3^2 + 16t_{2,2} x_2 x_3^2 + 24x_2^2 x_3^2 \Big) + e^{3t_{\mu A}} \left(-24t_{1,1} t_{2,2} t_{3,3} \right. \\
& - 24t_{2,2} t_{3,3} x_1 - 18t_{1,1} t_{3,3} x_2 - 52t_{3,3} x_1 x_2 - 32t_{1,1} t_{2,2} x_3 - 32t_{2,2} x_1 x_3 - 144x_1 x_2 x_3 \Big) \\
& + e^{4t_{\mu A}} \left(-48t_{1,1}^2 - 16t_{2,1} t_{2,2} - \frac{8t_{2,2}^3}{3} + 12t_{3,2}^2 + 6t_{3,2} t_{3,3}^2 + \frac{3t_{3,3}^4}{4} - 48t_{1,1} x_1 \right. \\
& + 90t_{3,2} t_{3,3} x_3 + \frac{45}{2} t_{3,3}^3 x_3 + 120t_{3,2} x_3^2 + 138t_{3,3}^2 x_3^2 + 288t_{3,3} x_3^3 + 192x_3^4 \Big) \\
& + e^{6t_{\mu A}} \left(204t_{2,2} t_{3,2} + 99t_{2,2} t_{3,3}^2 + 432t_{3,2} x_2 + 252t_{3,3}^2 x_2 + 298t_{2,2} t_{3,3} x_3 - 324t_{3,3} x_2 x_3 + 312t_{2,2} x_3^2 - 432x_2 x_3^2 \right) \\
& + e^{7t_{\mu A}} \left(1872t_{1,1} t_{3,3} + 2808t_{3,3} x_1 + 2112t_{1,1} x_3 + 2208x_1 x_3 \right) + e^{8t_{\mu A}} \left(2096t_{2,1} - \frac{536t_{2,2}^2}{3} + 1728t_{2,2} x_2 + 3888x_2^2 \right) \\
& + e^{10t_{\mu A}} \left(-8976t_{3,2} - 13428t_{3,3}^2 - 20928t_{3,3} x_3 - 10752x_3^2 \right) + e^{12t_{\mu A}} \left(-10704t_{2,2} - 68256x_2 \right) + 295680e^{16t_{\mu A}},
\end{aligned}$$

$$\begin{aligned}
\phi_{88}^3 = & e^{t_{\mu A}} \left(-t_{1,1} t_{2,1} - \frac{1}{6} t_{1,1} t_{2,2}^2 - 2t_{2,1} x_1 - \frac{1}{3} t_{2,2}^2 x_1 - 2t_{1,1} t_{2,2} x_2 - 4t_{2,2} x_1 x_2 - 3t_{1,1} x_2^2 - 6x_1 x_2^2 \right) + e^{2t_{\mu A}} \left(-4t_{2,2} t_{3,1} \right. \\
& - t_{2,2} t_{3,2} t_{3,3} - \frac{1}{24} t_{2,2} t_{3,3}^3 - 6t_{3,1} x_2 - \frac{3}{2} t_{3,2} t_{3,3} x_2 - \frac{1}{16} t_{3,3}^3 x_2 + 12t_{3,2} x_2 x_3 + 3t_{3,3}^2 x_2 x_3 + 18t_{3,3} x_2 x_3^2 + 24x_2 x_3^3 \Big) \\
& + e^{3t_{\mu A}} \left(-21t_{1,1} t_{3,2} - \frac{21}{4} t_{1,1} t_{3,3}^2 - 18t_{3,2} x_1 - \frac{9}{2} t_{3,3}^2 x_1 - 18t_{1,1} t_{3,3} x_3 - 24t_{1,1} x_3^2 \right) + e^{4t_{\mu A}} \left(-52t_{2,1} t_{3,3} \right. \\
& + \frac{10}{3} t_{2,2}^2 t_{3,3} - 14t_{2,2} t_{3,3} x_2 - 48t_{3,3} x_2^2 - 48t_{2,1} x_3 + 8t_{2,2}^2 x_3 - 48t_{2,2} x_2 x_3 \Big) + e^{5t_{\mu A}} \left(23t_{1,1} t_{2,2} + 78t_{2,2} x_1 \right. \\
& + 30t_{1,1} x_2 - 36x_1 x_2 \Big) + e^{6t_{\mu A}} \left(-36t_{3,1} + 195t_{3,2} t_{3,3} + \frac{405t_{3,3}^3}{8} + 144t_{3,2} x_3 + 180t_{3,3}^2 x_3 + 192t_{3,3} x_3^2 \right) \\
& + e^{8t_{\mu A}} \left(-156t_{2,2} t_{3,3} + 1314t_{3,3} x_2 + 48t_{2,2} x_3 + 336x_2 x_3 \right) + e^{9t_{\mu A}} \left(240t_{1,1} + 1152x_1 \right) + e^{12t_{\mu A}} \left(-9300t_{3,3} - 4464x_3 \right).
\end{aligned}$$

A.8. ψ for $A = (2, 3, 4)$.

$$\psi^1 = x_1 + e^{3t_{\mu A}} (18t_{3,3} + 48x_3), \quad \psi^2 = -48e^{4t_{\mu A}}, \quad \psi^3 = 0.$$

A.9. Frobenius potential for $A = (2, 3, 5)$.

$$\begin{aligned}
& \mathcal{F}(t_1, t_{1,1}, t_{2,1}, t_{2,2}, t_{3,1}, t_{3,2}, t_{3,3}, t_{3,4}, t_{\mu_A}) \\
&= \frac{t_1^2 t_{\mu_A}}{2} + t_1 \left(\frac{t_{1,1}^2}{4} + \frac{t_{2,1} t_{2,2}}{3} + \frac{t_{3,2} t_{3,3}}{5} + \frac{t_{3,1} t_{3,4}}{5} \right) + \left(\frac{t_{2,1}^3}{18} + \frac{t_{3,1} t_{3,2}^2}{10} + \frac{t_{3,1}^2 t_{3,3}}{10} \right) \\
&+ \left(-\frac{t_{1,1}^4}{96} - \frac{t_{2,1}^2 t_{2,2}^2}{36} - \frac{t_{3,2}^2 t_{3,3}^2}{50} - \frac{t_{3,1} t_{3,3}^3}{150} - \frac{t_{3,2}^3 t_{3,4}}{150} - \frac{1}{25} t_{3,1} t_{3,2} t_{3,3} t_{3,4} - \frac{t_{3,1}^2 t_{3,4}^2}{100} \right) \\
&+ \left(\frac{t_{2,1} t_{2,2}^4}{648} + \frac{t_{3,3}^5}{3000} + \frac{1}{250} t_{3,2} t_{3,3}^3 t_{3,4} + \frac{1}{250} t_{3,2}^2 t_{3,3} t_{3,4}^2 + \frac{1}{500} t_{3,1} t_{3,3}^2 t_{3,4}^2 + \frac{1}{750} t_{3,1} t_{3,2} t_{3,4}^3 \right) \\
&+ \left(-\frac{t_{2,2}^6}{19440} - \frac{3 t_{3,3}^4 t_{3,4}^2}{10000} - \frac{t_{3,2} t_{3,3}^2 t_{3,4}^3}{1500} - \frac{t_{3,2}^2 t_{3,4}^4}{7500} - \frac{t_{3,1} t_{3,3} t_{3,4}^4}{15000} \right) + \left(\frac{7 t_{3,3}^3 t_{3,4}^4}{150000} + \frac{11 t_{3,2} t_{3,3} t_{3,4}^5}{375000} + \frac{t_{3,1} t_{3,4}^6}{2250000} \right) \\
&+ \left(-\frac{13 t_{3,3}^2 t_{3,4}^6}{4500000} - \frac{t_{3,2} t_{3,4}^7}{3150000} \right) + \frac{7 t_{3,3} t_{3,4}^8}{90000000} - \frac{7 t_{3,4}^{10}}{810000000} + e^{t_{\mu_A}} \left(t_{1,1} t_{2,1} t_{3,1} + \frac{1}{6} t_{1,1} t_{2,2}^2 t_{3,1} + \frac{1}{10} t_{1,1} t_{2,1} t_{3,3}^2 \right. \\
&+ \frac{1}{60} t_{1,1} t_{2,2}^2 t_{3,3}^2 + \frac{1}{5} t_{1,1} t_{2,1} t_{3,2} t_{3,4} + \frac{1}{30} t_{1,1} t_{2,2}^2 t_{3,2} t_{3,4} + \frac{1}{50} t_{1,1} t_{2,1} t_{3,3} t_{3,4}^2 + \frac{1}{300} t_{1,1} t_{2,2}^2 t_{3,3} t_{3,4}^2 \\
&+ \frac{t_{1,1} t_{2,1} t_{3,4}^4}{3000} + \frac{t_{1,1} t_{2,2}^2 t_{3,4}^4}{18000} \left. \right) + e^{2 t_{\mu_A}} \left(\frac{t_{2,2} t_{3,1}^2}{2} + \frac{t_{2,1}^2 t_{3,2}}{2} + \frac{1}{2} t_{1,1}^2 t_{2,2} t_{3,2} + \frac{1}{6} t_{2,1} t_{2,2}^2 t_{3,2} + \frac{t_{2,2}^4 t_{3,2}}{72} \right. \\
&+ \frac{1}{10} t_{2,2} t_{3,1} t_{3,3}^2 + \frac{t_{2,2} t_{3,3}^4}{200} + \frac{1}{5} t_{2,2} t_{3,1} t_{3,2} t_{3,4} + \frac{1}{5} t_{2,1}^2 t_{3,3} t_{3,4} + \frac{1}{5} t_{1,1}^2 t_{2,2} t_{3,3} t_{3,4} + \frac{1}{15} t_{2,1} t_{2,2}^2 t_{3,3} t_{3,4} \\
&+ \frac{1}{180} t_{2,2}^4 t_{3,3} t_{3,4} + \frac{1}{50} t_{2,2} t_{3,2} t_{3,3}^2 t_{3,4} + \frac{1}{50} t_{2,2} t_{3,2}^2 t_{3,4}^2 + \frac{1}{50} t_{2,2} t_{3,1} t_{3,3} t_{3,4}^2 + \frac{1}{500} t_{2,2} t_{3,3}^3 t_{3,4}^2 + \frac{t_{2,1}^2 t_{3,4}^3}{75} \\
&+ \frac{1}{75} t_{1,1}^2 t_{2,2} t_{3,4}^3 + \frac{1}{225} t_{2,1} t_{2,2}^2 t_{3,4}^3 + \frac{t_{2,2}^4 t_{3,4}^3}{2700} + \frac{1}{250} t_{2,2} t_{3,2} t_{3,3} t_{3,4}^3 + \frac{t_{2,2} t_{3,1} t_{3,4}^4}{3000} + \frac{7 t_{2,2} t_{3,3}^2 t_{3,4}^4}{30000} \\
&+ \frac{t_{2,2} t_{3,2} t_{3,4}^5}{15000} + \frac{t_{2,2} t_{3,3} t_{3,4}^6}{150000} + \frac{t_{2,2} t_{3,4}^8}{18000000} \left. \right) + e^{3 t_{\mu_A}} \left(t_{1,1} t_{3,1} t_{3,2} + \frac{t_{1,1}^3 t_{3,3}}{3} + t_{1,1} t_{2,1} t_{2,2} t_{3,3} + \frac{1}{6} t_{1,1} t_{2,2}^3 t_{3,3} \right. \\
&+ \frac{1}{10} t_{1,1} t_{3,2} t_{3,3}^2 + \frac{1}{5} t_{1,1} t_{3,2}^2 t_{3,4} + \frac{2}{5} t_{1,1} t_{3,1} t_{3,3} t_{3,4} + \frac{1}{25} t_{1,1} t_{3,3}^3 t_{3,4} + \frac{t_{1,1}^3 t_{3,4}^2}{10} + \frac{3}{10} t_{1,1} t_{2,1} t_{2,2} t_{3,4}^2 \\
&+ \frac{1}{20} t_{1,1} t_{2,2}^3 t_{3,4}^2 + \frac{1}{10} t_{1,1} t_{3,2} t_{3,3} t_{3,4}^2 + \frac{2}{75} t_{1,1} t_{3,1} t_{3,4}^3 + \frac{4}{375} t_{1,1} t_{3,3}^2 t_{3,4}^3 + \frac{17 t_{1,1} t_{3,2} t_{3,4}^4}{3000} + \frac{t_{1,1} t_{3,3} t_{3,4}^5}{1500} \\
&+ \frac{t_{1,1} t_{3,4}^7}{112500} \left. \right) + e^{4 t_{\mu_A}} \left(\frac{t_{2,2}^2 t_{3,2}^2}{4} + t_{2,1} t_{3,1} t_{3,3} + \frac{1}{6} t_{2,2}^2 t_{3,1} t_{3,3} + \frac{t_{2,1} t_{3,3}^3}{10} + \frac{t_{2,2}^2 t_{3,3}^3}{60} + t_{1,1}^2 t_{2,1} t_{3,4} \right. \\
&+ \frac{1}{2} t_{2,1}^2 t_{2,2} t_{3,4} + \frac{2}{3} t_{1,1}^2 t_{2,2}^2 t_{3,4} + \frac{1}{6} t_{2,1} t_{2,2}^3 t_{3,4} + \frac{t_{2,2}^5 t_{3,4}}{72} + \frac{1}{5} t_{2,1} t_{3,2} t_{3,3} t_{3,4} + \frac{7}{30} t_{2,2}^2 t_{3,2} t_{3,3} t_{3,4} \\
&+ \frac{3}{10} t_{2,1} t_{3,1} t_{3,4}^2 + \frac{1}{20} t_{2,2}^2 t_{3,1} t_{3,4}^2 + \frac{1}{20} t_{2,1} t_{3,3}^2 t_{3,4}^2 + \frac{29}{600} t_{2,2}^2 t_{3,3}^2 t_{3,4}^2 + \frac{3}{50} t_{2,1} t_{3,2} t_{3,4}^3 + \frac{7}{300} t_{2,2}^2 t_{3,2} t_{3,4}^3 \\
&+ \frac{19 t_{2,1} t_{3,3} t_{3,4}^4}{3000} + \frac{23 t_{2,2}^2 t_{3,3} t_{3,4}^4}{3600} + \frac{t_{2,1} t_{3,4}^6}{10000} + \frac{7 t_{2,2}^2 t_{3,4}^6}{36000} \left. \right) + e^{5 t_{\mu_A}} \left(t_{1,1} t_{2,1}^2 + t_{1,1}^3 t_{2,2} + \frac{4}{3} t_{1,1} t_{2,1} t_{2,2}^2 \right. \\
&+ \frac{7 t_{1,1} t_{2,2}^4}{36} + t_{1,1} t_{2,2} t_{3,2} t_{3,3} + t_{1,1} t_{2,2} t_{3,1} t_{3,4} + \frac{1}{2} t_{1,1} t_{2,2} t_{3,3}^2 t_{3,4} + \frac{1}{2} t_{1,1} t_{2,2} t_{3,2} t_{3,4}^2 + \frac{1}{6} t_{1,1} t_{2,2} t_{3,3} t_{3,4}^3 \\
&+ \frac{1}{120} t_{1,1} t_{2,2} t_{3,4}^5 \left. \right) + e^{6 t_{\mu_A}} \left(t_{1,1}^2 t_{3,1} + t_{2,1} t_{2,2} t_{3,1} + \frac{t_{2,2}^3 t_{3,1}}{6} + \frac{t_{3,2}^3}{6} + \frac{3 t_{1,1}^2 t_{3,3}^2}{5} + \frac{1}{10} t_{2,1} t_{2,2} t_{3,3}^2 \right. \\
&+ \frac{11 t_{2,2}^3 t_{3,3}^2}{60} + \frac{t_{3,1}^2 t_{3,4}}{2} + \frac{7}{10} t_{1,1}^2 t_{3,2} t_{3,4} + \frac{6}{5} t_{2,1} t_{2,2} t_{3,2} t_{3,4} + \frac{1}{5} t_{2,2}^3 t_{3,2} t_{3,4} + \frac{1}{5} t_{3,2}^2 t_{3,3} t_{3,4} + \frac{1}{10} t_{3,1} t_{3,3}^2 t_{3,4} \\
&+ \frac{t_{3,3}^4 t_{3,4}}{200} + \frac{1}{5} t_{3,1} t_{3,2} t_{3,4}^2 + \frac{13}{25} t_{1,1}^2 t_{3,3} t_{3,4}^2 + \frac{21}{50} t_{2,1} t_{2,2} t_{3,3} t_{3,4}^2 + \frac{17}{100} t_{2,2}^3 t_{3,3} t_{3,4}^2 + \frac{1}{10} t_{3,2} t_{3,3}^2 t_{3,4}^2 \\
&+ \frac{t_{3,2}^2 t_{3,4}^3}{30} + \frac{1}{50} t_{3,1} t_{3,3} t_{3,4}^3 + \frac{19 t_{3,3}^3 t_{3,4}^3}{1500} + \frac{22 t_{1,1}^2 t_{3,4}^4}{375} + \frac{27 t_{2,1} t_{2,2} t_{3,4}^4}{1000} + \frac{39 t_{2,2}^3 t_{3,4}^4}{2000} + \frac{11}{750} t_{3,2} t_{3,3} t_{3,4}^4 \\
&+ \frac{t_{3,1} t_{3,4}^5}{3000} + \frac{7 t_{3,3}^2 t_{3,4}^5}{30000} + \frac{19 t_{3,2} t_{3,4}^6}{45000} + \frac{67 t_{3,3} t_{3,4}^7}{450000} + \frac{521 t_{3,4}^9}{162000000} \left. \right) + e^{7 t_{\mu_A}} \left(t_{1,1} t_{2,1} t_{3,2} + \frac{7}{6} t_{1,1} t_{2,2}^2 t_{3,2} \right. \\
&+ \frac{7}{5} t_{1,1} t_{2,1} t_{3,3} t_{3,4} + \frac{49}{30} t_{1,1} t_{2,2}^2 t_{3,3} t_{3,4} + \frac{49}{150} t_{1,1} t_{2,1} t_{3,4}^3 + \frac{343}{900} t_{1,1} t_{2,2}^2 t_{3,4}^3 \left. \right) + e^{8 t_{\mu_A}} \left(t_{2,2} t_{3,1} t_{3,2} + 2 t_{1,1}^2 t_{2,2} t_{3,3} \right.
\end{aligned}$$

$$\begin{aligned}
& + t_{2,1}t_{2,2}^2t_{3,3} + \frac{t_{2,2}^4t_{3,3}}{6} + \frac{3}{5}t_{2,2}t_{3,2}t_{3,3}^2 + \frac{1}{5}t_{2,2}t_{3,2}^2t_{3,4} + \frac{2}{5}t_{2,2}t_{3,1}t_{3,3}t_{3,4} + \frac{6}{25}t_{2,2}t_{3,3}^3t_{3,4} + \frac{t_{2,1}^2t_{3,4}^2}{2} \\
& + \frac{8}{5}t_{1,1}^2t_{2,2}t_{3,4}^2 + \frac{7}{15}t_{2,1}t_{2,2}^2t_{3,4}^2 + \frac{17t_{2,2}^4t_{3,4}^2}{90} + \frac{2}{5}t_{2,2}t_{3,2}t_{3,3}t_{3,4}^2 + \frac{2}{75}t_{2,2}t_{3,1}t_{3,4}^3 + \frac{18}{125}t_{2,2}t_{3,3}^2t_{3,4}^3 \\
& + \frac{19}{375}t_{2,2}t_{3,2}t_{3,4}^4 + \frac{2}{75}t_{2,2}t_{3,3}t_{3,4}^5 + \frac{34t_{2,2}^2t_{3,4}^7}{28125} \Big) + e^{9t_{\mu A}} \left(t_{1,1}t_{3,1}t_{3,3} + \frac{t_{1,1}t_{3,3}^3}{10} + t_{1,1}^3t_{3,4} + 3t_{1,1}t_{2,1}t_{2,2}t_{3,4} \right. \\
& + \frac{3}{2}t_{1,1}t_{2,2}^3t_{3,4} + \frac{6}{5}t_{1,1}t_{3,2}t_{3,3}t_{3,4} + \frac{3}{10}t_{1,1}t_{3,1}t_{3,4}^2 + \frac{9}{20}t_{1,1}t_{3,3}^2t_{3,4}^2 + \frac{9}{25}t_{1,1}t_{3,2}t_{3,4}^3 + \frac{153t_{1,1}t_{3,3}t_{3,4}^4}{1000} \\
& + \frac{81t_{1,1}t_{3,4}^6}{10000} \Big) + e^{10t_{\mu A}} \left(t_{1,1}^2t_{2,1} + t_{2,1}^2t_{2,2} + \frac{13t_{1,1}^2t_{2,2}^2}{6} + \frac{t_{2,1}t_{2,2}^3}{3} + \frac{23t_{2,2}^5}{180} + t_{2,1}t_{3,2}t_{3,3} + \frac{1}{6}t_{2,2}^2t_{3,2}t_{3,3} \right. \\
& + t_{2,1}t_{3,1}t_{3,4} + \frac{1}{6}t_{2,2}^2t_{3,1}t_{3,4} + \frac{1}{2}t_{2,1}t_{3,3}^2t_{3,4} + \frac{7}{12}t_{2,2}^2t_{3,3}^2t_{3,4} + \frac{1}{2}t_{2,1}t_{3,2}t_{3,4}^2 + \frac{7}{12}t_{2,2}^2t_{3,2}t_{3,4}^2 \\
& + \frac{1}{6}t_{2,1}t_{3,3}t_{3,4}^3 + \frac{19}{36}t_{2,2}^2t_{3,3}t_{3,4}^3 + \frac{t_{2,1}t_{3,4}^5}{120} + \frac{43t_{2,2}^2t_{3,4}^5}{720} \Big) + e^{11t_{\mu A}} \left(t_{1,1}t_{2,2}t_{3,1} + \frac{11}{10}t_{1,1}t_{2,2}t_{3,3}^2 \right. \\
& + \frac{11}{5}t_{1,1}t_{2,2}t_{3,2}t_{3,4} + \frac{121}{50}t_{1,1}t_{2,2}t_{3,3}t_{3,4}^2 + \frac{1331t_{1,1}t_{2,2}t_{3,4}^4}{3000} \Big) + e^{12t_{\mu A}} \left(\frac{t_{3,1}^2}{2} + t_{1,1}^2t_{3,2} + \frac{t_{2,2}^3t_{3,2}}{2} + \frac{t_{3,1}t_{3,3}^2}{10} \right. \\
& + \frac{53t_{3,3}^4}{600} + \frac{t_{3,1}t_{3,2}t_{3,4}}{5} + \frac{7}{5}t_{1,1}^2t_{3,3}t_{3,4} + t_{2,1}t_{2,2}t_{3,3}t_{3,4} + \frac{41}{30}t_{2,2}^3t_{3,3}t_{3,4} + \frac{1}{50}t_{3,2}t_{3,3}^2t_{3,4} + \frac{27t_{3,2}^2t_{3,4}^2}{100} \\
& + \frac{1}{50}t_{3,1}t_{3,3}t_{3,4}^2 + \frac{51t_{3,3}^3t_{3,4}^2}{500} + \frac{62t_{1,1}^2t_{3,4}^3}{75} + \frac{3}{10}t_{2,1}t_{2,2}t_{3,4}^3 + \frac{53t_{2,2}^3t_{3,4}^3}{100} + \frac{51}{250}t_{3,2}t_{3,3}t_{3,4}^3 \\
& + \frac{t_{3,1}t_{3,4}^4}{3000} + \frac{2557t_{3,3}^2t_{3,4}^4}{30000} + \frac{67t_{3,2}t_{3,4}^5}{5000} + \frac{717t_{3,3}t_{3,4}^6}{50000} + \frac{5117t_{3,4}^8}{6000000} \Big) + e^{13t_{\mu A}} \left(t_{1,1}t_{2,1}t_{3,3} + \frac{13}{6}t_{1,1}t_{2,2}^2t_{3,3} \right. \\
& + \frac{13}{10}t_{1,1}t_{2,1}t_{3,4}^2 + \frac{169}{60}t_{1,1}t_{2,2}^2t_{3,4}^2 \Big) + e^{14t_{\mu A}} \left(\frac{t_{2,2}t_{3,2}^2}{2} + \frac{t_{2,1}^2t_{3,4}}{2} + \frac{7}{2}t_{1,1}^2t_{2,2}t_{3,4} + \frac{7}{6}t_{1,1}t_{2,2}^2t_{3,4} \right. \\
& + \frac{49t_{2,2}^4t_{3,4}}{72} + \frac{7}{5}t_{2,2}t_{3,2}t_{3,3}t_{3,4} + \frac{49}{50}t_{2,2}t_{3,3}^2t_{3,4}^2 + \frac{49}{150}t_{2,2}t_{3,2}t_{3,4}^3 + \frac{343}{750}t_{2,2}t_{3,3}t_{3,4}^4 + \frac{2401t_{2,2}t_{3,4}^6}{45000} \Big) \\
& + e^{15t_{\mu A}} \left(\frac{t_{1,1}^3}{3} + 2t_{1,1}t_{2,1}t_{2,2} + \frac{4t_{1,1}t_{2,2}^3}{3} + t_{1,1}t_{3,3}^2t_{3,4} + t_{1,1}t_{3,2}t_{3,4}^2 + t_{1,1}t_{3,3}t_{3,4}^3 + \frac{7t_{1,1}t_{3,4}^5}{60} \right) \\
& + e^{16t_{\mu A}} \left(t_{2,2}^2t_{3,3}^2 + t_{2,1}t_{3,2}t_{3,4} + \frac{2}{3}t_{2,2}^2t_{3,2}t_{3,4} + \frac{2}{5}t_{2,1}t_{3,3}t_{3,4}^2 + \frac{28}{15}t_{2,2}^2t_{3,3}t_{3,4}^2 + \frac{2t_{2,1}t_{3,4}^4}{75} \right. \\
& + \frac{148t_{2,2}^2t_{3,4}^4}{225} \Big) + e^{17t_{\mu A}} \left(t_{1,1}t_{2,2}t_{3,2} + \frac{17}{5}t_{1,1}t_{2,2}t_{3,3}t_{3,4} + \frac{289}{150}t_{1,1}t_{2,2}t_{3,4}^3 \right) + e^{18t_{\mu A}} \left(t_{1,1}^2t_{3,3} + t_{2,1}t_{2,2}t_{3,3} \right. \\
& + \frac{t_{2,2}^3t_{3,3}}{2} + \frac{t_{3,2}t_{3,3}^2}{2} + \frac{t_{3,3}^3t_{3,4}}{5} + \frac{9t_{1,1}^2t_{3,4}^2}{5} + \frac{3}{10}t_{2,1}t_{2,2}t_{3,4}^2 + \frac{33t_{2,2}^3t_{3,4}^2}{20} + \frac{3}{10}t_{3,2}t_{3,3}t_{3,4}^2 + \frac{3t_{3,3}^2t_{3,4}^3}{10} \\
& + \frac{9t_{3,2}t_{3,4}^4}{200} + \frac{63t_{3,3}t_{3,4}^5}{500} + \frac{81t_{3,4}^7}{5000} \Big) + e^{19t_{\mu A}} \left(t_{1,1}t_{2,1}t_{3,4} + \frac{19}{6}t_{1,1}t_{2,2}^2t_{3,4} \right) + e^{20t_{\mu A}} \left(\frac{t_{2,1}^2}{2} + t_{1,1}^2t_{2,2} \right. \\
& + \frac{t_{2,1}t_{2,2}^2}{6} + \frac{37t_{2,2}^4}{72} + \frac{1}{2}t_{2,2}t_{3,3}^2t_{3,4} + \frac{1}{2}t_{2,2}t_{3,2}t_{3,4}^2 + \frac{3}{2}t_{2,2}t_{3,3}t_{3,4}^3 + \frac{43t_{2,2}t_{3,4}^5}{120} \Big) + e^{21t_{\mu A}} \left(t_{1,1}t_{3,2}t_{3,4} \right. \\
& + \frac{7}{5}t_{1,1}t_{3,3}t_{3,4}^2 + \frac{49t_{1,1}t_{3,4}^4}{150} \Big) + e^{22t_{\mu A}} \left(\frac{t_{2,2}^2t_{3,2}}{2} + \frac{11}{5}t_{2,2}^2t_{3,3}t_{3,4} + \frac{121t_{2,2}^2t_{3,4}^3}{75} \right) + e^{23t_{\mu A}} \left(t_{1,1}t_{2,2}t_{3,3} \right. \\
& + \frac{23}{10}t_{1,1}t_{2,2}t_{3,4}^2 \Big) + e^{24t_{\mu A}} \left(\frac{t_{3,2}^2}{4} + t_{1,1}^2t_{3,4} + t_{2,2}^3t_{3,4} + \frac{t_{3,2}t_{3,3}t_{3,4}}{5} + \frac{27t_{3,3}^2t_{3,4}^2}{50} + \frac{t_{3,2}t_{3,4}^3}{75} \right. \\
& + \frac{229t_{3,3}t_{3,4}^4}{750} + \frac{977t_{3,4}^6}{11250} \Big) + e^{25t_{\mu A}} t_{1,1}t_{2,2}^2 + e^{26t_{\mu A}} \left(\frac{t_{2,2}t_{3,3}^2}{2} + \frac{13}{10}t_{2,2}t_{3,3}t_{3,4}^2 + \frac{169t_{2,2}t_{3,4}^4}{200} \right) \\
& + e^{27t_{\mu A}} \left(t_{1,1}t_{3,3}t_{3,4} + \frac{3t_{1,1}t_{3,4}^3}{10} \right) + \frac{7}{4}e^{28t_{\mu A}} t_{2,2}^2t_{3,4}^2 + e^{29t_{\mu A}} t_{1,1}t_{2,2}t_{3,4} + e^{30t_{\mu A}} \left(\frac{t_{1,1}^2}{2} + \frac{t_{2,2}^3}{3} + \frac{t_{3,3}t_{3,4}^3}{3} \right. \\
& + \frac{t_{3,4}^5}{5} \Big) + e^{32t_{\mu A}} \left(t_{2,2}t_{3,3}t_{3,4} + \frac{4t_{2,2}t_{3,4}^3}{5} \right) + \frac{1}{2}e^{34t_{\mu A}} t_{2,2}^2t_{3,4} + e^{36t_{\mu A}} \left(\frac{t_{3,3}^2}{6} + \frac{t_{3,3}t_{3,4}^2}{10} + \frac{53t_{3,4}^4}{200} \right) \\
& + \frac{1}{2}e^{38t_{\mu A}} t_{2,2}t_{3,4}^2 + \frac{1}{4}e^{40t_{\mu A}} t_{2,2}^2 + \frac{1}{6}e^{42t_{\mu A}} t_{3,4}^3 + \frac{1}{8}e^{48t_{\mu A}} t_{3,4}^2 + \frac{e^{60t_{\mu A}}}{60}.
\end{aligned}$$

A.10. Coordinate change for $A = (2, 3, 5)$.

$$\begin{aligned}
s_1(\mathbf{t}) &= t_1 + e^{2t_{\mu A}} \left(-2t_{2,2}t_{3,2} - \frac{4}{5}t_{2,2}t_{3,3}t_{3,4} - \frac{4}{75}t_{2,2}t_{3,4}^2 \right) + e^{3t_{\mu A}} \left(-6t_{1,1}t_{3,3} - \frac{9}{5}t_{1,1}t_{3,4}^2 \right) - 12e^{4t_{\mu A}}t_{2,1}t_{3,4} \\
&\quad - 10e^{5t_{\mu A}}t_{1,1}t_{2,2} + e^{6t_{\mu A}} \left(-60t_{3,1} + 12t_{3,3}^2 + 18t_{3,2}t_{3,4} + \frac{108}{5}t_{3,3}t_{3,4}^2 + \frac{12t_{3,4}^4}{5} \right) \\
&\quad + e^{8t_{\mu A}} \left(264t_{2,2}t_{3,3} + \frac{656}{5}t_{2,2}t_{3,4}^2 \right) + 918e^{9t_{\mu A}}t_{1,1}t_{3,4} + e^{10t_{\mu A}}(1380t_{2,1} + 90t_{2,2}^2) \\
&\quad + e^{12t_{\mu A}} \left(-252t_{3,2} - \frac{33204}{5}t_{3,3}t_{3,4} - \frac{66768t_{3,4}^3}{25} \right) - 36566e^{14t_{\mu A}}t_{2,2}t_{3,4} - 72390e^{15t_{\mu A}}t_{1,1} \\
&\quad + e^{18t_{\mu A}}(334020t_{3,3} + 689508t_{3,4}^2) + 2297860e^{20t_{\mu A}}t_{2,2} - 60218748e^{24t_{\mu A}}t_{3,4} + 1594559250e^{30t_{\mu A}}, \\
s_{1,1}(\mathbf{t}) &= t_{1,1} + e^{3t_{\mu A}} \left(-8t_{3,3} - \frac{12t_{3,4}^2}{5} \right) - 24e^{5t_{\mu A}}t_{2,2} + 1000e^{9t_{\mu A}}t_{3,4} - 59400e^{15t_{\mu A}}, \\
s_{2,1}(\mathbf{t}) &= t_{2,1} + \frac{t_{2,2}^2}{6} + e^{2t_{\mu A}} \left(-3t_{3,2} - \frac{6}{5}t_{3,3}t_{3,4} - \frac{2t_{3,4}^3}{25} \right) - 15e^{4t_{\mu A}}t_{2,2}t_{3,4} - 24e^{5t_{\mu A}}t_{1,1} \\
&\quad + e^{8t_{\mu A}} \left(225t_{3,3} + \frac{615t_{3,4}^2}{2} \right) + 1725e^{10t_{\mu A}}t_{2,2} - 56640e^{14t_{\mu A}}t_{3,4} + 2508000e^{20t_{\mu A}}, \\
s_{2,2}(\mathbf{t}) &= t_{2,2} - 27e^{4t_{\mu A}}t_{3,4} + 2376e^{10t_{\mu A}}, \\
s_{3,1}(\mathbf{t}) &= t_{3,1} + \frac{t_{3,3}^2}{10} + \frac{1}{5}t_{3,2}t_{3,4} + \frac{1}{50}t_{3,3}t_{3,4}^2 + \frac{t_{3,4}^4}{3000} + e^{2t_{\mu A}} \left(-3t_{2,2}t_{3,3} - \frac{9}{10}t_{2,2}t_{3,4}^2 \right) - 8e^{3t_{\mu A}}t_{1,1}t_{3,4} \\
&\quad + e^{4t_{\mu A}} \left(-15t_{2,1} + \frac{5t_{2,2}^2}{2} \right) + e^{6t_{\mu A}} \left(-80t_{3,2} + 13t_{3,3}t_{3,4} + \frac{341t_{3,4}^3}{30} \right) + 522e^{8t_{\mu A}}t_{2,2}t_{3,4} \\
&\quad + 1360e^{9t_{\mu A}}t_{1,1} + e^{12t_{\mu A}} \left(1872t_{3,3} - \frac{23967t_{3,4}^2}{5} \right) - 56115e^{14t_{\mu A}}t_{2,2} + 387405e^{18t_{\mu A}}t_{3,4} + 3926880e^{24t_{\mu A}}, \\
s_{3,2}(\mathbf{t}) &= -4e^{2t_{\mu A}}t_{2,2}t_{3,4} + t_{3,2} + \frac{2}{5}t_{3,3}t_{3,4} + \frac{2t_{3,4}^3}{75} - 10e^{3t_{\mu A}}t_{1,1} + e^{6t_{\mu A}}(-120t_{3,3} - 8t_{3,4}^2) \\
&\quad + 865e^{8t_{\mu A}}t_{2,2} + 6496e^{12t_{\mu A}}t_{3,4} - 971000e^{18t_{\mu A}}, \\
s_{3,3}(\mathbf{t}) &= t_{3,3} + \frac{3t_{3,4}^2}{10} - 5e^{2t_{\mu A}}t_{2,2} - 165e^{6t_{\mu A}}t_{3,4} + 23745e^{12t_{\mu A}}, \\
s_{3,4}(\mathbf{t}) &= t_{3,4} - 250e^{6t_{\mu A}}.
\end{aligned}$$

A.11. ϕ for $A = (2, 3, 5)$.

$$\phi_{ij}^a = 0, \quad a = 1, 2, 3, i = 1, j = 1, \dots, 9.$$

$$\phi_{22}^1 = \frac{x_1}{2} + e^{5t_{\mu A}}(40t_{2,2} + 60x_2) + e^{9t_{\mu A}}(-1464t_{3,4} - 480x_3) + 160320e^{15t_{\mu A}},$$

$$\phi_{22}^2 = e^{4t_{\mu A}}(16t_{3,4} + 20x_3) - 3680e^{10t_{\mu A}}, \quad \phi_{22}^3 = 168e^{6t_{\mu A}}.$$

$$\phi_{23}^1 = e^{4t_{\mu A}}(8t_{3,4} + 10x_3) - 1720e^{10t_{\mu A}}, \quad \phi_{23}^2 = 35e^{5t_{\mu A}}, \quad \phi_{23}^3 = -e^{t_{\mu A}}.$$

$$\begin{aligned}
\phi_{24}^1 &= e^{2t_{\mu A}} \left(-2t_{3,2} - \frac{4}{5}t_{3,3}t_{3,4} - \frac{4t_{3,4}^3}{75} - 3t_{3,3}x_3 - \frac{9}{10}t_{3,4}^2x_3 - 4t_{3,4}x_3^2 - 5x_3^3 \right) + e^{4t_{\mu A}} \left(\frac{8}{3}t_{2,2}t_{3,4} - 4t_{3,4}x_2 \right. \\
&\quad \left. + \frac{25}{3}t_{2,2}x_3 - 5x_2x_3 \right) + e^{5t_{\mu A}}(15t_{1,1} - 12x_1) + e^{8t_{\mu A}} \left(208t_{3,3} + \frac{512t_{3,4}^2}{5} + 519t_{3,4}x_3 + 880x_3^2 \right) \\
&\quad + e^{10t_{\mu A}} \left(-\frac{1900t_{2,2}}{3} + 605x_2 \right) + e^{14t_{\mu A}}(-25642t_{3,4} - 56985x_3) + 1658800e^{20t_{\mu A}}, \\
\phi_{24}^2 &= e^{3t_{\mu A}} \left(-3t_{3,3} - \frac{9t_{3,4}^2}{10} - 4t_{3,4}x_3 - 5x_3^2 \right) + e^{5t_{\mu A}} \left(\frac{35t_{2,2}}{3} - 8x_2 \right) + e^{9t_{\mu A}}(324t_{3,4} + 655x_3) - 25215e^{15t_{\mu A}},
\end{aligned}$$

$$\begin{aligned}
\phi_{24}^3 &= e^{t_{\mu A}} \left(-\frac{t_{2,2}}{3} - x_2 \right) + e^{5t_{\mu A}} (7t_{3,4} + 10x_3) - 1091e^{11t_{\mu A}}. \quad \phi_{25}^1 = 24e^{6t_{\mu A}}, \quad \phi_{25}^2 = -e^{t_{\mu A}}, \quad \phi_{25}^3 = 0. \\
\phi_{26}^1 &= e^{2t_{\mu A}} (-2t_{2,2} - 3x_2) + e^{6t_{\mu A}} \left(\frac{174t_{3,4}}{5} - 6x_3 \right) - 2592e^{12t_{\mu A}}, \\
\phi_{26}^2 &= e^{t_{\mu A}} \left(-\frac{t_{3,4}}{5} - x_3 \right) + 68e^{7t_{\mu A}}, \quad \phi_{26}^3 = -3e^{3t_{\mu A}}. \\
\phi_{27}^1 &= e^{2t_{\mu A}} \left(-\frac{4}{5}t_{2,2}t_{3,4} - \frac{6}{5}t_{3,4}x_2 - 2t_{2,2}x_3 - 3x_2x_3 \right) + e^{3t_{\mu A}} (-3t_{1,1} - 4x_1) + e^{6t_{\mu A}} \left(\frac{144t_{3,3}}{5} + \frac{492t_{3,4}^2}{25} \right. \\
&\quad \left. + \frac{258}{5}t_{3,4}x_3 + 24x_3^2 \right) + e^{8t_{\mu A}} (176t_{2,2} + 264x_2) + e^{12t_{\mu A}} \left(-\frac{29544t_{3,4}}{5} - 5382x_3 \right) + 335304e^{18t_{\mu A}}, \\
\phi_{27}^2 &= e^{t_{\mu A}} \left(-\frac{t_{3,3}}{5} - \frac{t_{3,4}^2}{50} - \frac{2}{5}t_{3,4}x_3 - x_3^2 \right) - e^{3t_{\mu A}} t_{2,2} + e^{7t_{\mu A}} \left(\frac{131t_{3,4}}{5} + 88x_3 \right) - 1756e^{13t_{\mu A}}, \\
\phi_{27}^3 &= e^{3t_{\mu A}} \left(-\frac{6t_{3,4}}{5} - 2x_3 \right) + 111e^{9t_{\mu A}}. \\
\phi_{28}^1 &= e^{2t_{\mu A}} \left(-\frac{4}{5}t_{2,2}t_{3,3} - \frac{4}{25}t_{2,2}t_{3,4}^2 - \frac{6}{5}t_{3,3}x_2 - \frac{6}{25}t_{3,4}^2x_2 - \frac{6}{5}t_{2,2}t_{3,4}x_3 - \frac{9}{5}t_{3,4}x_2x_3 - 2t_{2,2}x_3^2 - 3x_2x_3^2 \right) \\
&\quad + e^{3t_{\mu A}} \left(-\frac{9}{5}t_{1,1}t_{3,4} - \frac{12}{5}t_{3,4}x_1 - 3t_{1,1}x_3 - 4x_1x_3 \right) + e^{6t_{\mu A}} \left(\frac{54t_{3,2}}{5} + \frac{744}{25}t_{3,3}t_{3,4} + \frac{864t_{3,4}^3}{125} + \frac{153}{5}t_{3,3}x_3 \right. \\
&\quad \left. + \frac{2091}{50}t_{3,4}^2x_3 + \frac{402}{5}t_{3,4}x_3^2 + 39x_3^3 \right) + e^{8t_{\mu A}} \left(\frac{568}{5}t_{2,2}t_{3,4} + \frac{852}{5}t_{3,4}x_2 + 267t_{2,2}x_3 + 459x_2x_3 \right) + e^{9t_{\mu A}} (207t_{1,1} + 500x_1) \\
&\quad + e^{12t_{\mu A}} \left(-\frac{21744t_{3,3}}{5} - \frac{105504t_{3,4}^2}{25} - \frac{73461}{5}t_{3,4}x_3 - 11712x_3^2 \right) + e^{14t_{\mu A}} (-12738t_{2,2} - 23040x_2) \\
&\quad + e^{18t_{\mu A}} \left(\frac{3865152t_{3,4}}{5} + 1136145x_3 \right) - 41422320e^{24t_{\mu A}}, \\
\phi_{28}^2 &= e^{t_{\mu A}} \left(-\frac{t_{3,2}}{5} - \frac{1}{25}t_{3,3}t_{3,4} - \frac{t_{3,4}^3}{750} - \frac{2}{5}t_{3,3}x_3 - \frac{2}{25}t_{3,4}^2x_3 - \frac{3}{5}t_{3,4}x_3^2 - x_3^3 \right) - \frac{3}{5}e^{3t_{\mu A}} t_{2,2}t_{3,4} + e^{4t_{\mu A}} (-2t_{1,1} - 8x_1) \\
&\quad + e^{7t_{\mu A}} \left(\frac{401t_{3,3}}{5} + \frac{1037t_{3,4}^2}{50} + \frac{644}{5}t_{3,4}x_3 + 243x_3^2 \right) + e^{9t_{\mu A}} (108t_{2,2} + 216x_2) + e^{13t_{\mu A}} \left(-\frac{46348t_{3,4}}{5} - 19920x_3 \right) \\
&\quad + 762485e^{19t_{\mu A}}, \\
\phi_{28}^3 &= e^{3t_{\mu A}} \left(-\frac{6t_{3,3}}{5} - \frac{6t_{3,4}^2}{25} - \frac{6}{5}t_{3,4}x_3 - 2x_3^2 \right) + e^{5t_{\mu A}} (-t_{2,2} + 3x_2) + e^{9t_{\mu A}} \left(\frac{213t_{3,4}}{5} + 202x_3 \right) - 3456e^{15t_{\mu A}}. \\
\phi_{29}^1 &= e^{2t_{\mu A}} \left(-4t_{2,2}t_{3,2} - \frac{8}{5}t_{2,2}t_{3,3}t_{3,4} - \frac{8}{75}t_{2,2}t_{3,4}^3 - 6t_{3,2}x_2 - \frac{12}{5}t_{3,3}t_{3,4}x_2 - \frac{4}{25}t_{3,4}^3x_2 - 6t_{2,2}t_{3,3}x_3 \right. \\
&\quad \left. - \frac{9}{5}t_{2,2}t_{3,4}^2x_3 - 9t_{3,3}x_2x_3 - \frac{27}{10}t_{3,4}^2x_2x_3 - 8t_{2,2}t_{3,4}x_3^2 - 12t_{3,4}x_2x_3^2 - 10t_{2,2}x_3^3 - 15x_2x_3^3 \right) + e^{3t_{\mu A}} \left(-9t_{1,1}t_{3,3} \right. \\
&\quad \left. - \frac{27}{10}t_{1,1}t_{3,4}^2 - 12t_{3,3}x_1 - \frac{18}{5}t_{3,4}^2x_1 - 12t_{1,1}t_{3,4}x_3 - 16t_{3,4}x_1x_3 - 15t_{1,1}x_3^2 - 20x_1x_3^2 \right) + e^{4t_{\mu A}} (10t_{2,2}^2x_3 + 15t_{2,2}x_2x_3) \\
&\quad + e^{5t_{\mu A}} (15t_{1,1}t_{2,2} - 60t_{2,2}x_1 - 96x_1x_2) + e^{6t_{\mu A}} \left(72t_{3,3}^2 + 84t_{3,2}t_{3,4} + \frac{384}{5}t_{3,3}t_{3,4}^2 + \frac{218t_{3,4}^4}{25} - 60t_{3,2}x_3 \right. \\
&\quad \left. + 270t_{3,3}t_{3,4}x_3 + \frac{433}{5}t_{3,4}^3x_3 + 120t_{3,3}x_3^2 + 300t_{3,4}^2x_3^2 + 330t_{3,4}x_3^3 \right) + e^{8t_{\mu A}} \left(1040t_{2,2}t_{3,3} + 344t_{2,2}t_{3,4}^2 + 1632t_{3,3}x_2 \right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{2688}{5} t_{3,4}^2 x_2 + 1800 t_{2,2} t_{3,4} x_3 + 3051 t_{3,4} x_2 x_3 + 3200 t_{2,2} x_3^2 + 4620 x_2 x_3^2 \Big) + e^{9t_{\mu A}} (1275 t_{1,1} t_{3,4} + 4500 t_{3,4} x_1 \\
& + 2700 t_{1,1} x_3 + 3440 x_1 x_3) + e^{10t_{\mu A}} (-60 t_{2,2}^2 + 2610 t_{2,2} x_2 + 4320 x_2^2) + e^{12t_{\mu A}} \left(-6144 t_{3,2} - \frac{236208}{5} t_{3,3} t_{3,4} \right. \\
& \left. - \frac{349576 t_{3,4}^3}{25} - 33936 t_{3,3} x_3 - \frac{396234}{5} t_{3,4}^2 x_3 - 130848 t_{3,4} x_3^2 - 32160 x_3^3 \right) + e^{14t_{\mu A}} (-153740 t_{2,2} t_{3,4} - 303600 t_{3,4} x_2 \\
& - 288330 t_{2,2} x_3 - 447885 x_2 x_3) + e^{15t_{\mu A}} (-135885 t_{1,1} - 445500 x_1) + e^{18t_{\mu A}} \left(3793104 t_{3,3} + \frac{30387216 t_{3,4}^2}{5} \right. \\
& \left. + 16278702 t_{3,4} x_3 + 9527640 x_3^2 \right) + e^{20t_{\mu A}} (13301840 t_{2,2} + 25757760 x_2) + e^{24t_{\mu A}} (-828787440 t_{3,4} - 884891040 x_3) \\
& + 34686244200 e^{30t_{\mu A}}, \\
\phi_{29}^2 = & -x_1 x_2 + e^{3t_{\mu A}} \left(-6 t_{2,2} t_{3,3} - \frac{9}{5} t_{2,2} t_{3,4}^2 - 8 t_{2,2} t_{3,4} x_3 - 10 t_{2,2} x_3^2 \right) + e^{4t_{\mu A}} (-16 t_{1,1} t_{3,4} - 32 t_{3,4} x_1 - 20 t_{1,1} x_3 \\
& - 40 x_1 x_3) + e^{5t_{\mu A}} \left(20 t_{2,1} + \frac{10 t_{2,2}^2}{3} \right) + e^{7t_{\mu A}} \left(96 t_{3,2} + \frac{1462}{5} t_{3,3} t_{3,4} + \frac{1969 t_{3,4}^3}{25} + 304 t_{3,3} x_3 + \frac{2256}{5} t_{3,4}^2 x_3 + 882 t_{3,4} x_3^2 \right. \\
& \left. + 540 x_3^3 \right) + e^{9t_{\mu A}} (984 t_{2,2} t_{3,4} + 864 t_{3,4} x_2 + 1640 t_{2,2} x_3 + 1440 x_2 x_3) + e^{10t_{\mu A}} (1400 t_{1,1} + 6880 x_1) \\
& + e^{13t_{\mu A}} \left(-48136 t_{3,3} - \frac{263324 t_{3,4}^2}{5} - 174608 t_{3,4} x_3 - 157260 x_3^2 \right) + e^{15t_{\mu A}} (-111570 t_{2,2} - 190080 x_2) \\
& + e^{19t_{\mu A}} (10377650 t_{3,4} + 15134160 x_3) - 584059200 e^{25t_{\mu A}}, \\
\phi_{29}^3 = & e^{t_{\mu A}} \left(-t_{2,1} - \frac{t_{2,2}^2}{6} - 2 t_{2,2} x_2 - 3 x_2^2 \right) + e^{3t_{\mu A}} \left(-3 t_{3,2} - \frac{6}{5} t_{3,3} t_{3,4} - \frac{2 t_{3,4}^3}{25} \right) + e^{5t_{\mu A}} (11 t_{2,2} t_{3,4} + 66 t_{3,4} x_2 \\
& + 20 t_{2,2} x_3) + e^{6t_{\mu A}} (48 t_{1,1} - 192 x_1) + e^{9t_{\mu A}} \left(1431 t_{3,3} + \frac{933 t_{3,4}^2}{10} + 1980 t_{3,4} x_3 + 3000 x_3^2 \right) + e^{11t_{\mu A}} (-571 t_{2,2} + 768 x_2) \\
& + e^{15t_{\mu A}} (-161184 t_{3,4} - 377880 x_3) + 15859680 e^{21t_{\mu A}}. \\
\phi_{33}^1 = & 30 e^{5t_{\mu A}}, \phi_{33}^2 = 0, \phi_{33}^3 = 0. \\
\phi_{34}^1 = & e^{3t_{\mu A}} \left(3 t_{3,3} + \frac{9 t_{3,4}^2}{10} + 4 t_{3,4} x_3 + 5 x_3^2 \right) + e^{5t_{\mu A}} (20 t_{2,2} + 45 x_2) + e^{9t_{\mu A}} (-1287 t_{3,4} - 865 x_3) + 129510 e^{15t_{\mu A}}, \\
\phi_{34}^2 = & \frac{x_2}{3} + e^{4t_{\mu A}} (8 t_{3,4} + 10 x_3) - 2090 e^{10t_{\mu A}}, \phi_{34}^3 = 114 e^{6t_{\mu A}}. \phi_{35}^1 = -e^{t_{\mu A}}, \phi_{35}^2 = 0, \phi_{35}^3 = 0. \\
\phi_{36}^1 = & e^{t_{\mu A}} \left(-\frac{t_{3,4}}{5} - x_3 \right) + 35 e^{7t_{\mu A}}, \phi_{36}^2 = -2 e^{2t_{\mu A}}, \phi_{36}^3 = 0. \\
\phi_{37}^1 = & e^{t_{\mu A}} \left(-\frac{t_{3,3}}{5} - \frac{t_{3,4}^2}{50} - \frac{2}{5} t_{3,4} x_3 - x_3^2 \right) + e^{3t_{\mu A}} (-t_{2,2} - 6 x_2) + e^{7t_{\mu A}} (77 t_{3,4} + 120 x_3) - 8001 e^{13t_{\mu A}}, \\
\phi_{37}^2 = & e^{2t_{\mu A}} \left(-\frac{4 t_{3,4}}{5} - 2 x_3 \right) + 120 e^{8t_{\mu A}}, \phi_{37}^3 = -4 e^{4t_{\mu A}}. \\
\phi_{38}^1 = & e^{t_{\mu A}} \left(-\frac{t_{3,2}}{5} - \frac{1}{25} t_{3,3} t_{3,4} - \frac{t_{3,4}^3}{750} - \frac{2}{5} t_{3,3} x_3 - \frac{2}{25} t_{3,4}^2 x_3 - \frac{3}{5} t_{3,4} x_3^2 - x_3^3 \right) + e^{3t_{\mu A}} \left(-\frac{3}{5} t_{2,2} t_{3,4} - \frac{18}{5} t_{3,4} x_2 \right. \\
& \left. - 6 x_2 x_3 \right) + 2 e^{4t_{\mu A}} t_{1,1} + e^{7t_{\mu A}} \left(17 t_{3,3} + \frac{83 t_{3,4}^2}{2} + 124 t_{3,4} x_3 + 165 x_3^2 \right) + e^{9t_{\mu A}} (57 t_{2,2} + 585 x_2) \\
& + e^{13t_{\mu A}} \left(-\frac{53553 t_{3,4}}{5} - 16000 x_3 \right) + 687555 e^{19t_{\mu A}}, \\
\phi_{38}^2 = & e^{2t_{\mu A}} \left(-\frac{4 t_{3,3}}{5} - \frac{4 t_{3,4}^2}{25} - \frac{6}{5} t_{3,4} x_3 - 2 x_3^2 \right) - 9 e^{4t_{\mu A}} x_2 + e^{8t_{\mu A}} (176 t_{3,4} + 330 x_3) - 17000 e^{14t_{\mu A}},
\end{aligned}$$

$$\begin{aligned}
\phi_{38}^3 &= e^{4t_{\mu A}} \left(-\frac{12t_{3,4}}{5} - 3x_3 \right) + 170e^{10t_{\mu A}}. \\
\phi_{39}^1 &= -x_1x_2 + e^{3t_{\mu A}} \left(-6t_{2,2}t_{3,3} - \frac{9}{5}t_{2,2}t_{3,4}^2 - 18t_{3,3}x_2 - \frac{27}{5}t_{3,4}^2x_2 - 8t_{2,2}t_{3,4}x_3 - 24t_{3,4}x_2x_3 - 10t_{2,2}x_3^2 - 30x_2x_3^2 \right) \\
&\quad + 15e^{4t_{\mu A}}x_1x_3 + e^{5t_{\mu A}}(-90t_{2,2}x_2 - 180x_2^2) + e^{7t_{\mu A}} \left(30t_{3,2} + 258t_{3,3}t_{3,4} + \frac{373t_{3,4}^3}{5} + 150t_{3,3}x_3 + 309t_{3,4}^2x_3 \right. \\
&\quad \left. + 450t_{3,4}x_3^2 + 150x_3^3 \right) + e^{9t_{\mu A}}(1800t_{2,2}t_{3,4} + 7290t_{3,4}x_2 + 2450t_{2,2}x_3 + 6900x_2x_3) - 300e^{10t_{\mu A}}t_{1,1} \\
&\quad + e^{13t_{\mu A}} \left(-27876t_{3,3} - \frac{354414t_{3,4}^2}{5} - 136518t_{3,4}x_3 - 62460x_3^2 \right) + e^{15t_{\mu A}}(-179100t_{2,2} - 730890x_2) \\
&\quad + e^{19t_{\mu A}}(12768810t_{3,4} + 10956150x_3) - 648482400e^{25t_{\mu A}}, \\
\phi_{39}^2 &= e^{2t_{\mu A}} \left(-4t_{3,2} - \frac{8}{5}t_{3,3}t_{3,4} - \frac{8t_{3,4}^3}{75} - 6t_{3,3}x_3 - \frac{9}{5}t_{3,4}^2x_3 - 8t_{3,4}x_3^2 - 10x_3^3 \right) + e^{4t_{\mu A}}(-36t_{3,4}x_2 + 10t_{2,2}x_3 \\
&\quad - 60x_2x_3) + e^{5t_{\mu A}}(55t_{1,1} + 30x_1) + e^{8t_{\mu A}}(300t_{3,3} + 506t_{3,4}^2 + 1890t_{3,4}x_3 + 2300x_3^2) + e^{10t_{\mu A}}(-720t_{2,2} + 7920x_2) \\
&\quad + e^{14t_{\mu A}}(-149020t_{3,4} - 267390x_3) + 11540300e^{20t_{\mu A}}, \\
\phi_{39}^3 &= e^{t_{\mu A}}(-t_{1,1} - 2x_1) + e^{4t_{\mu A}} \left(-4t_{3,3} - \frac{6t_{3,4}^2}{5} \right) + e^{6t_{\mu A}}(24t_{2,2} - 270x_2) + e^{10t_{\mu A}}(2960t_{3,4} + 4500x_3) \\
&\quad - 368820e^{16t_{\mu A}}. \\
\phi_{44}^1 &= -\frac{1}{3}x_1x_2 + e^{3t_{\mu A}} \left(-2t_{2,2}t_{3,3} - \frac{3}{5}t_{2,2}t_{3,4}^2 - \frac{8}{3}t_{2,2}t_{3,4}x_3 - \frac{10}{3}t_{2,2}x_3^2 \right) - 5e^{4t_{\mu A}}x_1x_3 + \frac{10}{3}e^{5t_{\mu A}}t_{2,2}^2 \\
&\quad + e^{7t_{\mu A}} \left(36t_{3,2} + \frac{162}{5}t_{3,3}t_{3,4} + \frac{159t_{3,4}^3}{25} + 54t_{3,3}x_3 + \frac{201}{5}t_{3,4}^2x_3 + 102t_{3,4}x_3^2 + 90x_3^3 \right) + e^{9t_{\mu A}} \left(162t_{2,2}t_{3,4} \right. \\
&\quad \left. - 18t_{3,4}x_2 + \frac{1100}{3}t_{2,2}x_3 + 90x_2x_3 \right) + e^{10t_{\mu A}}(-270t_{1,1} + 288x_1) + e^{13t_{\mu A}} \left(-4194t_{3,3} - \frac{15291t_{3,4}^2}{5} - 14592t_{3,4}x_3 \right. \\
&\quad \left. - 16590x_3^2 \right) + e^{15t_{\mu A}}(-13320t_{2,2} - 5220x_2) + e^{19t_{\mu A}}(578790t_{3,4} + 1180860x_3) - 32746050e^{25t_{\mu A}}, \\
\phi_{44}^2 &= \frac{x_2^2}{3} + e^{2t_{\mu A}} \left(-\frac{4t_{3,2}}{3} - \frac{8}{15}t_{3,3}t_{3,4} - \frac{8t_{3,4}^3}{225} - 2t_{3,3}x_3 - \frac{3}{5}t_{3,4}^2x_3 - \frac{8}{3}t_{3,4}x_3^2 - \frac{10x_3^3}{3} \right) + e^{4t_{\mu A}} \left(-4t_{3,4}x_2 \right. \\
&\quad \left. + \frac{10}{3}t_{2,2}x_3 \right) + e^{5t_{\mu A}} \left(\frac{25t_{1,1}}{3} - 10x_1 \right) + e^{8t_{\mu A}} \left(164t_{3,3} + \frac{898t_{3,4}^2}{15} + 342t_{3,4}x_3 + \frac{1820x_3^2}{3} \right) \\
&\quad + e^{10t_{\mu A}}(-60t_{2,2} + 622x_2) + e^{14t_{\mu A}} \left(-\frac{53368t_{3,4}}{3} - 39150x_3 \right) + \frac{3779600}{3}e^{20t_{\mu A}}, \\
\phi_{44}^3 &= e^{t_{\mu A}} \left(-\frac{t_{1,1}}{3} - \frac{2x_1}{3} \right) + e^{4t_{\mu A}} \left(\frac{8t_{3,3}}{3} + \frac{4t_{3,4}^2}{5} + 4t_{3,4}x_3 + 5x_3^2 \right) \\
&\quad + e^{6t_{\mu A}}(4t_{2,2} + 18x_2) + e^{10t_{\mu A}} \left(-\frac{1648t_{3,4}}{3} - 730x_3 \right) + 44640e^{16t_{\mu A}}. \\
\phi_{45}^1 &= e^{t_{\mu A}} \left(-\frac{t_{2,2}}{3} - x_2 \right) + 15e^{5t_{\mu A}}t_{3,4} - 1725e^{11t_{\mu A}}, \quad \phi_{45}^2 = 24e^{6t_{\mu A}}, \quad \phi_{45}^3 = -2e^{2t_{\mu A}} \\
\phi_{46}^1 &= e^{t_{\mu A}} \left(-\frac{1}{15}t_{2,2}t_{3,4} - \frac{1}{5}t_{3,4}x_2 - \frac{1}{3}t_{2,2}x_3 - x_2x_3 \right) - e^{2t_{\mu A}}t_{1,1} + e^{5t_{\mu A}} \left(-t_{3,3} + \frac{27t_{3,4}^2}{10} + 3t_{3,4}x_3 - 15x_3^2 \right) \\
&\quad + e^{7t_{\mu A}} \left(\frac{161t_{2,2}}{3} + 62x_2 \right) + e^{11t_{\mu A}}(-655t_{3,4} + 870x_3) + 543e^{17t_{\mu A}},
\end{aligned}$$

$$\begin{aligned}
\phi_{46}^2 &= e^{2t_{\mu A}} \left(-\frac{2t_{2,2}}{3} - x_2 \right) + e^{6t_{\mu A}} \left(\frac{54t_{3,4}}{5} - 6x_3 \right) - 180e^{12t_{\mu A}}, \quad \phi_{46}^3 = e^{2t_{\mu A}} \left(-\frac{2t_{3,4}}{5} - x_3 \right) + 52e^{8t_{\mu A}}. \\
\phi_{47}^1 &= e^{t_{\mu A}} \left(-\frac{1}{15}t_{2,2}t_{3,3} - \frac{1}{150}t_{2,2}t_{3,4}^2 - \frac{1}{5}t_{3,3}x_2 - \frac{1}{50}t_{3,4}^2x_2 - \frac{2}{15}t_{2,2}t_{3,4}x_3 - \frac{2}{5}t_{3,4}x_2x_3 - \frac{1}{3}t_{2,2}x_3^2 - x_2x_3^2 \right) \\
&+ e^{2t_{\mu A}} \left(-\frac{2}{5}t_{1,1}t_{3,4} - t_{1,1}x_3 - x_1x_3 \right) + e^{3t_{\mu A}} \left(-\frac{1}{3}t_{2,2}^2 + t_{2,2}x_2 \right) + e^{5t_{\mu A}} \left(4t_{3,2} + \frac{21}{5}t_{3,3}t_{3,4} + \frac{43t_{3,4}^3}{150} + 14t_{3,3}x_3 \right. \\
&+ \frac{27}{5}t_{3,4}^2x_3 + 17t_{3,4}x_3^2 + 10x_3^3 \left. \right) + e^{7t_{\mu A}} \left(\frac{142}{15}t_{2,2}t_{3,4} + \frac{109}{5}t_{3,4}x_2 + 42t_{2,2}x_3 + 132x_2x_3 \right) + e^{8t_{\mu A}} (92t_{1,1} + 96x_1) \\
&+ e^{11t_{\mu A}} \left(-1206t_{3,3} - \frac{1929t_{3,4}^2}{5} - 2758t_{3,4}x_3 - 2600x_3^2 \right) + e^{13t_{\mu A}} (-1770t_{2,2} - 5352x_2) \\
&+ e^{17t_{\mu A}} \left(\frac{601871t_{3,4}}{5} + 233253x_3 \right) - 7840445e^{23t_{\mu A}}, \\
\phi_{47}^2 &= e^{2t_{\mu A}} \left(-\frac{4}{15}t_{2,2}t_{3,4} - \frac{2}{5}t_{3,4}x_2 - \frac{2}{3}t_{2,2}x_3 \right) + e^{3t_{\mu A}} (-3t_{1,1} - 2x_1) \\
&+ e^{6t_{\mu A}} \left(\frac{84t_{3,3}}{5} + \frac{162t_{3,4}^2}{25} + \frac{58}{5}t_{3,4}x_3 + 4x_3^2 \right) + e^{8t_{\mu A}} (76t_{2,2} + 75x_2) + e^{12t_{\mu A}} (-2028t_{3,4} - 1160x_3) + 121914e^{18t_{\mu A}}, \\
\phi_{47}^3 &= e^{2t_{\mu A}} \left(-\frac{2t_{3,3}}{5} - \frac{t_{3,4}^2}{25} - \frac{2}{5}t_{3,4}x_3 - x_3^2 \right) + e^{4t_{\mu A}} \left(\frac{2t_{2,2}}{3} + 2x_2 \right) + e^{8t_{\mu A}} \left(\frac{14t_{3,4}}{5} + 93x_3 \right) - 1008e^{14t_{\mu A}}. \\
\phi_{48}^1 &= e^{t_{\mu A}} \left(-\frac{1}{15}t_{2,2}t_{3,2} - \frac{1}{75}t_{2,2}t_{3,3}t_{3,4} - \frac{t_{2,2}t_{3,4}^3}{2250} - \frac{1}{5}t_{3,2}x_2 - \frac{1}{25}t_{3,3}t_{3,4}x_2 - \frac{1}{750}t_{3,4}^3x_2 - \frac{2}{15}t_{2,2}t_{3,3}x_3 \right. \\
&- \frac{2}{75}t_{2,2}t_{3,4}^2x_3 - \frac{2}{5}t_{3,3}x_2x_3 - \frac{2}{25}t_{3,4}^2x_2x_3 - \frac{1}{5}t_{2,2}t_{3,4}x_3^2 - \frac{3}{5}t_{3,4}x_2x_3^2 - \frac{1}{3}t_{2,2}x_3^3 - x_2x_3^3 \left. \right) + e^{2t_{\mu A}} \left(-\frac{2}{5}t_{1,1}t_{3,3} \right. \\
&- \frac{2}{25}t_{1,1}t_{3,4}^2 - \frac{3}{5}t_{1,1}t_{3,4}x_3 - \frac{3}{5}t_{3,4}x_1x_3 - t_{1,1}x_3^2 - x_1x_3^2 \left. \right) + e^{3t_{\mu A}} \left(-\frac{1}{5}t_{2,2}^2t_{3,4} + \frac{3}{5}t_{2,2}t_{3,4}x_2 + 2t_{2,2}x_2x_3 \right) \\
&+ e^{4t_{\mu A}} \left(-\frac{4}{3}t_{1,1}t_{2,2} - t_{1,1}x_2 + 3x_1x_2 \right) + e^{5t_{\mu A}} \left(-\frac{2}{5}t_{3,3}^2 + \frac{27}{5}t_{3,2}t_{3,4} + \frac{34}{25}t_{3,3}t_{3,4}^2 + \frac{3t_{3,4}^4}{50} + \frac{48}{5}t_{3,3}t_{3,4}x_3 \right. \\
&+ \frac{69}{25}t_{3,4}^3x_3 + 2t_{3,3}x_3^2 + 15t_{3,4}^2x_3^2 + 21t_{3,4}x_3^3 \left. \right) + e^{7t_{\mu A}} \left(\frac{442}{15}t_{2,2}t_{3,3} - \frac{38}{75}t_{2,2}t_{3,4}^2 + \frac{124}{5}t_{3,3}x_2 + \frac{79}{25}t_{3,4}^2x_2 + \frac{458}{15}t_{2,2}t_{3,4}x_3 \right. \\
&+ \frac{386}{5}t_{3,4}x_2x_3 + 112t_{2,2}x_3^2 + 192x_2x_3^2 \left. \right) + e^{8t_{\mu A}} \left(\frac{476}{5}t_{1,1}t_{3,4} + \frac{288}{5}t_{3,4}x_1 + 192t_{1,1}x_3 + 246x_1x_3 \right) \\
&+ e^{9t_{\mu A}} (-23t_{2,2}^2 - 201t_{2,2}x_2) + e^{11t_{\mu A}} \left(-689t_{3,2} - \frac{4921}{5}t_{3,3}t_{3,4} - \frac{32063t_{3,4}^3}{150} - 1704t_{3,3}x_3 - \frac{12828}{5}t_{3,4}^2x_3 \right. \\
&- 6239t_{3,4}x_3^2 - 2585x_3^3 \left. \right) + e^{13t_{\mu A}} \left(-124t_{2,2}t_{3,4} - \frac{11796}{5}t_{3,4}x_2 - \frac{23239}{3}t_{2,2}x_3 - 13531x_2x_3 \right) \\
&+ e^{14t_{\mu A}} (-10547t_{1,1} - 12000x_1) + e^{17t_{\mu A}} \left(\frac{508921t_{3,3}}{5} + \frac{5545087t_{3,4}^2}{50} + \frac{2898999}{5}t_{3,4}x_3 + 489433x_3^2 \right) \\
&+ e^{19t_{\mu A}} (128058t_{2,2} + 367230x_2) + e^{23t_{\mu A}} (-18359274t_{3,4} - 34006980x_3) + 886406335e^{29t_{\mu A}}, \\
\phi_{48}^2 &= e^{2t_{\mu A}} \left(-\frac{4}{15}t_{2,2}t_{3,3} - \frac{4}{75}t_{2,2}t_{3,4}^2 - \frac{2}{5}t_{3,3}x_2 - \frac{2}{25}t_{3,4}^2x_2 - \frac{2}{5}t_{2,2}t_{3,4}x_3 - \frac{2}{3}t_{2,2}x_3^2 \right) + e^{3t_{\mu A}} \left(-\frac{9}{5}t_{1,1}t_{3,4} - \frac{6}{5}t_{3,4}x_1 \right. \\
&- 3t_{1,1}x_3 - 2x_1x_3 \left. \right) + e^{4t_{\mu A}} \left(-2t_{2,1} - \frac{t_{2,2}^2}{3} - 2t_{2,2}x_2 - 9x_2^2 \right) + e^{6t_{\mu A}} \left(\frac{114t_{3,2}}{5} + \frac{444}{25}t_{3,3}t_{3,4} + \frac{394t_{3,4}^3}{125} + \frac{198}{5}t_{3,3}x_3 \right. \\
&+ \frac{513}{25}t_{3,4}^2x_3 + \frac{282}{5}t_{3,4}x_3^2 + 54x_3^3 \left. \right) + e^{8t_{\mu A}} \left(\frac{574}{15}t_{2,2}t_{3,4} + 178t_{3,4}x_2 + 62t_{2,2}x_3 \right) + e^{9t_{\mu A}} (366t_{1,1} + 492x_1)
\end{aligned}$$

$$\begin{aligned}
& + e^{12t\mu_A} \left(-3954t_{3,3} - \frac{14043t_{3,4}^2}{5} - 8680t_{3,4}x_3 - 10130x_3^2 \right) + e^{14t\mu_A} \left(-\frac{20816t_{2,2}}{3} - 20149x_2 \right) \\
& + e^{18t\mu_A} \left(\frac{3055182t_{3,4}}{5} + 756580x_3 \right) - 33881640e^{24t\mu_A}, \\
\phi_{48}^3 = & e^{2t\mu_A} \left(-\frac{2t_{3,2}}{5} - \frac{2}{25}t_{3,3}t_{3,4} - \frac{t_{3,4}^3}{375} - \frac{2}{5}t_{3,3}x_3 - \frac{2}{25}t_{3,4}^2x_3 - \frac{3}{5}t_{3,4}x_3^2 - x_3^3 \right) + e^{4t\mu_A} \left(\frac{2}{5}t_{2,2}t_{3,4} + \frac{6}{5}t_{3,4}x_2 \right. \\
& \left. + 2t_{2,2}x_3 \right) + e^{5t\mu_A} (t_{1,1} + 6x_1) + e^{8t\mu_A} \left(-\frac{16t_{3,3}}{5} - \frac{346t_{3,4}^2}{25} + \frac{134}{5}t_{3,4}x_3 + 138x_3^2 \right) + e^{10t\mu_A} \left(-\frac{550t_{2,2}}{3} - 172x_2 \right) \\
& + e^{14t\mu_A} \left(\frac{20256t_{3,4}}{5} - 4417x_3 \right) - 190116e^{20t\mu_A}. \\
\phi_{49}^1 = & -\frac{1}{3}t_{2,2}x_1x_2 - x_1x_2^2 + e^{2t\mu_A} \left(-2t_{1,1}t_{3,2} - \frac{4}{5}t_{1,1}t_{3,3}t_{3,4} - \frac{4}{75}t_{1,1}t_{3,4}^3 - 3t_{1,1}t_{3,3}x_3 - \frac{9}{10}t_{1,1}t_{3,4}^2x_3 - 3t_{3,3}x_1x_3 \right. \\
& \left. - \frac{9}{10}t_{3,4}^2x_1x_3 - 4t_{1,1}t_{3,4}x_3^2 - 4t_{3,4}x_1x_3^2 - 5t_{1,1}x_3^3 - 5x_1x_3^3 \right) + e^{3t\mu_A} \left(-2t_{2,2}^2t_{3,3} - \frac{3}{5}t_{2,2}^2t_{3,4}^2 - \frac{8}{3}t_{2,2}^2t_{3,4}x_3 - \frac{10}{3}t_{2,2}^2x_3^2 \right) \\
& + e^{4t\mu_A} (-8t_{1,1}t_{2,2}t_{3,4} - 12t_{1,1}t_{3,4}x_2 + 27t_{3,4}x_1x_2 - 5t_{1,1}t_{2,2}x_3 - 5t_{2,2}x_1x_3 - 15t_{1,1}x_2x_3 - 60x_1x_2x_3) \\
& + e^{5t\mu_A} \left(5t_{1,1}^2 + 10t_{3,2}t_{3,3} + 4t_{3,3}^2t_{3,4} + 3t_{3,2}t_{3,4}^2 + \frac{22}{15}t_{3,3}t_{3,4}^3 + \frac{2t_{3,4}^5}{25} + 30t_{2,1}x_2 - 5t_{2,2}^2x_2 + 30t_{3,1}x_3 + 45t_{3,3}^2x_3 \right. \\
& \left. - 18t_{3,2}t_{3,4}x_3 + \frac{81}{5}t_{3,3}t_{3,4}^2x_3 + \frac{63}{20}t_{3,4}^4x_3 + 30t_{3,2}x_3^2 + 68t_{3,3}t_{3,4}x_3^2 + \frac{88}{5}t_{3,4}^3x_3^2 + 160t_{3,3}x_3^3 + 48t_{3,4}^2x_3^3 + 120t_{3,4}^4x_3^3 \right. \\
& \left. + 150x_3^5 \right) + e^{7t\mu_A} \left(214t_{2,2}t_{3,2} + \frac{708}{5}t_{2,2}t_{3,3}t_{3,4} + \frac{1688}{75}t_{2,2}t_{3,4}^3 + 174t_{3,2}x_2 + \frac{528}{5}t_{3,3}t_{3,4}x_2 + \frac{386}{25}t_{3,4}^3x_2 \right. \\
& \left. + 186t_{2,2}t_{3,3}x_3 + \frac{1079}{5}t_{2,2}t_{3,4}^2x_3 + 576t_{3,3}x_2x_3 + \frac{1104}{5}t_{3,4}^2x_2x_3 + 528t_{2,2}t_{3,4}x_3^2 + 828t_{3,4}x_2x_3^2 + 260t_{2,2}x_3^3 \right. \\
& \left. + 960x_2x_3^3 \right) + e^{8t\mu_A} \left(296t_{1,1}t_{3,3} + \frac{1844}{5}t_{1,1}t_{3,4}^2 + 288t_{3,3}x_1 + \frac{432}{5}t_{3,4}^2x_1 + 963t_{1,1}t_{3,4}x_3 + 1674t_{3,4}x_1x_3 + 760t_{1,1}x_3^2 \right. \\
& \left. + 1555x_1x_3^2 \right) + e^{9t\mu_A} \left(-360t_{2,1}t_{3,4} - 108t_{2,2}^2t_{3,4} - 882t_{2,2}t_{3,4}x_2 - 450t_{2,1}x_3 - \frac{385}{3}t_{2,2}^2x_3 - 1200t_{2,2}x_2x_3 \right) \\
& + e^{10t\mu_A} (700t_{1,1}t_{2,2} + 1440t_{2,2}x_1 + 1305t_{1,1}x_2 + 1485x_1x_2) + e^{11t\mu_A} \left(-1800t_{3,1} - 2748t_{3,3}^2 - 4016t_{3,2}t_{3,4} \right. \\
& \left. - \frac{20846}{5}t_{3,3}t_{3,4}^2 - \frac{50116t_{3,4}^4}{75} - 810t_{3,2}x_3 - 21180t_{3,3}t_{3,4}x_3 - \frac{38712}{5}t_{3,4}^3x_3 - 25280t_{3,3}x_3^2 - 28150t_{3,4}^2x_3^2 \right. \\
& \left. - 47570t_{3,4}x_3^3 - 39000x_3^4 \right) + e^{13t\mu_A} \left(-25496t_{2,2}t_{3,3} - \frac{25184}{5}t_{2,2}t_{3,4}^2 - 40644t_{3,3}x_2 - \frac{51876}{5}t_{3,4}^2x_2 - 58138t_{2,2}t_{3,4}x_3 \right. \\
& \left. - 109992t_{3,4}x_2x_3 - 71760t_{2,2}x_3^2 - 175590x_2x_3^2 \right) + e^{14t\mu_A} (-85818t_{1,1}t_{3,4} - 108000t_{3,4}x_1 - 85755t_{1,1}x_3 \\
& - 199035x_1x_3) + e^{15t\mu_A} (41400t_{2,1} + 8220t_{2,2}^2 + 24660t_{2,2}x_2 - 124740x_2^2) + e^{17t\mu_A} \left(429246t_{3,2} + \frac{7504862}{5}t_{3,3}t_{3,4} \right. \\
& \left. + \frac{12203119t_{3,4}^3}{25} + 2577924t_{3,3}x_3 + \frac{18217296}{5}t_{3,4}^2x_3 + 7882462t_{3,4}x_3^2 + 6053940x_3^3 \right) + e^{19t\mu_A} (1836892t_{2,2}t_{3,4} \\
& + 6134130t_{3,4}x_2 + 4371240t_{2,2}x_3 + 11865240x_2x_3) + e^{20t\mu_A} (5316640t_{1,1} + 10692000x_1) + e^{23t\mu_A} \left(-116485460t_{3,3} \right. \\
& \left. - 155871716t_{3,4}^2 - 584406000t_{3,4}x_3 - 576147300x_3^2 \right) + e^{25t\mu_A} (-157960260t_{2,2} - 520036830x_2) \\
& + e^{29t\mu_A} (19266865810t_{3,4} + 29950525410x_3) - 786837196800e^{35t\mu_A},
\end{aligned}$$

$$\begin{aligned}
\phi_{49}^2 = & e^{2t_{\mu A}} \left(-\frac{4}{3}t_{2,2}t_{3,2} - \frac{8}{15}t_{2,2}t_{3,3}t_{3,4} - \frac{8}{225}t_{2,2}t_{3,4}^2 - 2t_{3,2}x_2 - \frac{4}{5}t_{3,3}t_{3,4}x_2 - \frac{4}{75}t_{3,4}^2x_2 - 2t_{2,2}t_{3,3}x_3 - \frac{3}{5}t_{2,2}t_{3,4}^2x_3 \right. \\
& - \frac{8}{3}t_{2,2}t_{3,4}x_3^2 - \frac{10}{3}t_{2,2}x_3^3 \Big) + e^{3t_{\mu A}} \left(-9t_{1,1}t_{3,3} - \frac{27}{10}t_{1,1}t_{3,4}^2 - 6t_{3,3}x_1 - \frac{9}{5}t_{3,4}^2x_1 - 12t_{1,1}t_{3,4}x_3 - 8t_{3,4}x_1x_3 - 15t_{1,1}x_3^2 \right. \\
& - 10x_1x_3^2 \Big) + e^{4t_{\mu A}} \left(-8t_{2,1}t_{3,4} - \frac{4}{3}t_{2,2}^2t_{3,4} - 8t_{2,2}t_{3,4}x_2 - 36t_{3,4}x_2^2 - 10t_{2,1}x_3 + \frac{5}{3}t_{2,2}^2x_3 \right) + e^{5t_{\mu A}} \left(-\frac{5}{3}t_{1,1}t_{2,2} \right. \\
& - 10t_{2,2}x_1 - 40t_{1,1}x_2 - 36x_1x_2 \Big) + e^{6t_{\mu A}} \left(36t_{3,3}^2 + 84t_{3,2}t_{3,4} + \frac{276}{5}t_{3,3}t_{3,4}^2 + \frac{137t_{3,4}^4}{25} - 30t_{3,2}x_3 + 198t_{3,3}t_{3,4}x_3 \right. \\
& + \frac{311}{5}t_{3,4}^2x_3 + 60t_{3,3}x_3^2 + 234t_{3,4}^2x_3^2 + 270t_{3,4}x_3^3 \Big) + e^{8t_{\mu A}} \left(440t_{2,2}t_{3,3} + \frac{308}{3}t_{2,2}t_{3,4}^2 + 600t_{3,3}x_2 + 712t_{3,4}^2x_2 \right. \\
& + 630t_{2,2}t_{3,4}x_3 + \frac{4100}{3}t_{2,2}x_3^2 \Big) + e^{9t_{\mu A}} (2616t_{1,1}t_{3,4} + 3348t_{3,4}x_1 + 3795t_{1,1}x_3 + 3110x_1x_3) + e^{10t_{\mu A}} \left(1670t_{2,1} \right. \\
& - \frac{245t_{2,2}^2}{3} + 3110t_{2,2}x_2 + 7920x_2^2 \Big) + e^{12t_{\mu A}} \left(-11730t_{3,2} - 34248t_{3,3}t_{3,4} - \frac{55018t_{3,4}^3}{5} - 38460t_{3,3}x_3 - 49128t_{3,4}^2x_3 \right. \\
& - 91320t_{3,4}x_3^2 - 39150x_3^3 \Big) + e^{14t_{\mu A}} \left(-\frac{223078}{3}t_{2,2}t_{3,4} - 245636t_{3,4}x_2 - 124050t_{2,2}x_3 - 41580x_2x_3 \right) \\
& + e^{15t_{\mu A}} (-292425t_{1,1} - 398070x_1) + e^{18t_{\mu A}} \left(3320334t_{3,3} + \frac{22793661t_{3,4}^2}{5} + 10242792t_{3,4}x_3 + 7544340x_3^2 \right) \\
& + e^{20t_{\mu A}} \left(\frac{23541890t_{2,2}}{3} + 17836720x_2 \right) + e^{24t_{\mu A}} (-615527100t_{3,4} - 590410650x_3) + 25783674900e^{30t_{\mu A}}, \\
\phi_{49}^3 = & e^{t_{\mu A}} \left(-\frac{1}{3}t_{1,1}t_{2,2} - \frac{2}{3}t_{2,2}x_1 - t_{1,1}x_2 - 2x_1x_2 \right) + e^{2t_{\mu A}} \left(-2t_{3,1} - \frac{t_{3,3}^2}{5} - \frac{2}{5}t_{3,2}t_{3,4} - \frac{1}{25}t_{3,3}t_{3,4}^2 - \frac{t_{3,4}^4}{1500} \right) \\
& + e^{4t_{\mu A}} \left(\frac{14}{3}t_{2,2}t_{3,3} + \frac{7}{5}t_{2,2}t_{3,4}^2 + 14t_{3,3}x_2 + \frac{21}{5}t_{3,4}^2x_2 + 8t_{2,2}t_{3,4}x_3 + 10t_{2,2}x_3^2 + 30x_2x_3^2 \right) + e^{5t_{\mu A}} (19t_{1,1}t_{3,4} \\
& + 54t_{3,4}x_1 - 60x_1x_3) + e^{6t_{\mu A}} (54t_{2,1} + 7t_{2,2}^2 + 42t_{2,2}x_2 - 108x_2^2) + e^{8t_{\mu A}} \left(472t_{3,2} - \frac{406}{5}t_{3,3}t_{3,4} - \frac{5131t_{3,4}^3}{75} \right. \\
& + 960t_{3,3}x_3 + 120t_{3,4}^2x_3 + 990t_{3,4}x_3^2 + 1500x_3^3 \Big) + e^{10t_{\mu A}} \left(-\frac{6760}{3}t_{2,2}t_{3,4} - 604t_{3,4}x_2 - 2930t_{2,2}x_3 - 1800x_2x_3 \right) \\
& + e^{11t_{\mu A}} (-641t_{1,1} + 2970x_1) + e^{14t_{\mu A}} \left(-48924t_{3,3} + \frac{91584t_{3,4}^2}{5} - 121722t_{3,4}x_3 - 244440x_3^2 \right) \\
& + e^{16t_{\mu A}} (157950t_{2,2} - 94122x_2) + e^{20t_{\mu A}} (2898126t_{3,4} + 15130620x_3) - 377238060e^{26t_{\mu A}}.
\end{aligned}$$

$$\phi_{55}^1 = 0, \phi_{55}^2 = 0, \phi_{55}^3 = 0. \quad \phi_{56}^1 = 3e^{3t_{\mu A}}, \phi_{56}^2 = 0, \phi_{56}^3 = 0.$$

$$\phi_{57}^1 = \frac{6}{5}e^{3t_{\mu A}}t_{3,4} - 225e^{9t_{\mu A}}, \phi_{57}^2 = 8e^{4t_{\mu A}}, \phi_{57}^3 = 0.$$

$$\phi_{58}^1 = e^{3t_{\mu A}} \left(\frac{6t_{3,3}}{5} + \frac{6t_{3,4}^2}{25} \right) + e^{5t_{\mu A}} (15t_{2,2} + 27x_2) - 615e^{9t_{\mu A}}t_{3,4} + 56640e^{15t_{\mu A}},$$

$$\phi_{58}^2 = \frac{24}{5}e^{4t_{\mu A}}t_{3,4} - 1000e^{10t_{\mu A}}, \phi_{58}^3 = \frac{x_3}{5} + 54e^{6t_{\mu A}}.$$

$$\begin{aligned}
\phi_{59}^1 &= -x_1x_3 + e^{3t_{\mu A}} \left(-6t_{3,2} - \frac{12}{5}t_{3,3}t_{3,4} - \frac{4t_{3,4}^3}{25} - 18t_{3,3}x_3 - \frac{27}{5}t_{3,4}^2x_3 - 24t_{3,4}x_3^2 - 30x_3^3 \right) + e^{5t_{\mu A}} (12t_{2,2}t_{3,4} \\
&\quad - 36t_{3,4}x_2 - 30t_{2,2}x_3 - 180x_2x_3) + 60e^{6t_{\mu A}}t_{1,1} + e^{9t_{\mu A}} (1080t_{3,3} + 660t_{3,4}^2 + 6210t_{3,4}x_3 + 6000x_3^2) \\
&\quad + e^{11t_{\mu A}} (690t_{2,2} + 12240x_2) + e^{15t_{\mu A}} (-341088t_{3,4} - 712530x_3) + 29208000e^{21t_{\mu A}}, \\
\phi_{59}^2 &= e^{t_{\mu A}} (-t_{1,1} - 2x_1) + e^{4t_{\mu A}} \left(-4t_{3,3} - \frac{6t_{3,4}^2}{5} - 48t_{3,4}x_3 - 60x_3^2 \right) + 84e^{6t_{\mu A}}t_{2,2} \\
&\quad + e^{10t_{\mu A}} (2420t_{3,4} + 12000x_3) - 474660e^{16t_{\mu A}}, \\
\phi_{59}^3 &= e^{2t_{\mu A}} (-4t_{2,2} - 6x_2) + e^{6t_{\mu A}} (36t_{3,4} - 300x_3) + 14976e^{12t_{\mu A}}. \\
\\
\phi_{66}^1 &= e^{3t_{\mu A}} \left(\frac{6t_{3,4}}{5} + 6x_3 \right) - 480e^{9t_{\mu A}}, \quad \phi_{66}^2 = 12e^{4t_{\mu A}}, \quad \phi_{66}^3 = 0. \\
\phi_{67}^1 &= e^{3t_{\mu A}} \left(\frac{3t_{3,3}}{5} + \frac{3t_{3,4}^2}{10} + \frac{12}{5}t_{3,4}x_3 + 3x_3^2 \right) + e^{5t_{\mu A}} (19t_{2,2} + 42x_2) + e^{9t_{\mu A}} (-858t_{3,4} - 585x_3) + 90144e^{15t_{\mu A}}, \\
\phi_{67}^2 &= e^{4t_{\mu A}} \left(\frac{32t_{3,4}}{5} + 14x_3 \right) - 1810e^{10t_{\mu A}}, \quad \phi_{67}^3 = \frac{x_3}{5} + 60e^{6t_{\mu A}}. \\
\\
\phi_{68}^1 &= -\frac{1}{5}x_1x_3 + e^{3t_{\mu A}} \left(-\frac{9t_{3,2}}{5} - \frac{3}{5}t_{3,3}t_{3,4} - \frac{3t_{3,4}^3}{250} - \frac{6}{5}t_{3,3}x_3 - \frac{3}{5}t_{3,4}^2x_3 - 3t_{3,4}x_3^2 - 3x_3^3 \right) \\
&\quad + e^{5t_{\mu A}} \left(\frac{24}{5}t_{2,2}t_{3,4} + \frac{9}{5}t_{3,4}x_2 + 9t_{2,2}x_3 + 9x_2x_3 \right) + 9e^{6t_{\mu A}}t_{1,1} + e^{9t_{\mu A}} \left(159t_{3,3} + \frac{45t_{3,4}^2}{2} + 255t_{3,4}x_3 + 705x_3^2 \right) \\
&\quad + e^{11t_{\mu A}} (-1106t_{2,2} - 930x_2) + e^{15t_{\mu A}} \left(-\frac{13326t_{3,4}}{5} - 37866x_3 \right) + 375135e^{21t_{\mu A}}, \\
\phi_{68}^2 &= e^{t_{\mu A}} \left(-\frac{t_{1,1}}{5} - \frac{2x_1}{5} \right) + e^{4t_{\mu A}} \left(-\frac{4t_{3,3}}{5} + \frac{6t_{3,4}^2}{25} - \frac{6}{5}t_{3,4}x_3 - 6x_3^2 \right) \\
&\quad + e^{6t_{\mu A}} \left(\frac{114t_{2,2}}{5} + 27x_2 \right) + e^{10t_{\mu A}} (-274t_{3,4} + 410x_3) + 16148e^{16t_{\mu A}}, \\
\phi_{68}^3 &= \frac{x_3^2}{5} + e^{2t_{\mu A}} \left(-\frac{4t_{2,2}}{5} - \frac{6x_2}{5} \right) + e^{6t_{\mu A}} \left(\frac{54t_{3,4}}{5} - 9x_3 \right) - \frac{3804}{5}e^{12t_{\mu A}}. \\
\phi_{69}^1 &= -\frac{1}{5}t_{3,4}x_1x_3 - x_1x_3^2 + e^{2t_{\mu A}} (-2t_{1,1}t_{2,2} - 3t_{1,1}x_2 - 3x_1x_2) + e^{3t_{\mu A}} \left(-\frac{6}{5}t_{3,2}t_{3,4} - \frac{12}{25}t_{3,3}t_{3,4}^2 - \frac{4t_{3,4}^4}{125} + 6t_{3,2}x_3 \right. \\
&\quad - \frac{6}{5}t_{3,3}t_{3,4}x_3 - \frac{23}{25}t_{3,4}^3x_3 - \frac{24}{5}t_{3,4}^2x_3^2 - 6t_{3,4}x_3^3) + e^{5t_{\mu A}} \left(10t_{2,2}t_{3,3} + \frac{27}{5}t_{2,2}t_{3,4}^2 + 42t_{3,3}x_2 + \frac{27}{5}t_{3,4}^2x_2 - 18t_{2,2}t_{3,4}x_3 \right. \\
&\quad - 90t_{2,2}x_3^2 - 90x_2x_3^2) + e^{6t_{\mu A}} (30t_{1,1}t_{3,4} - 60t_{1,1}x_3 + 20x_1x_3) + e^{7t_{\mu A}} (168t_{2,2}^2 + 522t_{2,2}x_2 + 540x_2^2) \\
&\quad + e^{9t_{\mu A}} \left(-60t_{3,2} - 42t_{3,3}t_{3,4} + \frac{301t_{3,4}^3}{5} + 360t_{3,3}x_3 + 1230t_{3,4}^2x_3 + 4650t_{3,4}x_3^2 + 1500x_3^3 \right) + e^{11t_{\mu A}} (-8474t_{2,2}t_{3,4} \\
&\quad - 16302t_{3,4}x_2 + 5670t_{2,2}x_3 + 3240x_2x_3) - 2712e^{12t_{\mu A}}t_{1,1} + e^{15t_{\mu A}} \left(4500t_{3,3} + \frac{236862t_{3,4}^2}{5} - 399738t_{3,4}x_3 \right. \\
&\quad \left. - 487740x_3^2 \right) + e^{17t_{\mu A}} (683538t_{2,2} + 1398690x_2) + e^{21t_{\mu A}} (-6361830t_{3,4} + 31002000x_3) + 73886880e^{27t_{\mu A}},
\end{aligned}$$

$$\begin{aligned}
\phi_{69}^2 &= e^{t\mu_A} \left(-\frac{1}{5}t_{1,1}t_{3,4} - \frac{2}{5}t_{3,4}x_1 - t_{1,1}x_3 - 2x_1x_3 \right) + e^{2t\mu_A} \left(-2t_{2,1} - \frac{t_{2,2}^2}{3} \right) + e^{4t\mu_A} \left(6t_{3,2} + \frac{8}{5}t_{3,3}t_{3,4} \right. \\
&\quad \left. - \frac{2t_{3,4}^2}{25} + 14t_{3,3}x_3 - \frac{27}{5}t_{3,4}^2x_3 - 36t_{3,4}x_3^2 - 30x_3^3 \right) + e^{6t\mu_A} \left(\frac{354}{5}t_{2,2}t_{3,4} + 108t_{3,4}x_2 + 54t_{2,2}x_3 + 180x_2x_3 \right) \\
&\quad + e^{7t\mu_A} (68t_{1,1} + 40x_1) + e^{10t\mu_A} (-370t_{3,3} - 683t_{3,4}^2 + 2270t_{3,4}x_3 + 9000x_3^2) + e^{12t\mu_A} (-12570t_{2,2} \\
&\quad - 23760x_2) + e^{16t\mu_A} (165932t_{3,4} - 452490x_3) - 3828000e^{22t\mu_A}, \\
\phi_{69}^3 &= e^{2t\mu_A} \left(-\frac{4}{5}t_{2,2}t_{3,4} - \frac{6}{5}t_{3,4}x_2 - 2t_{2,2}x_3 \right) + e^{3t\mu_A} (-9t_{1,1} - 6x_1) + e^{6t\mu_A} (36t_{3,3} + 18t_{3,4}^2 - 18t_{3,4}x_3 - 300x_3^2) \\
&\quad + e^{8t\mu_A} (836t_{2,2} + 1200x_2) + e^{12t\mu_A} \left(-\frac{63324t_{3,4}}{5} + 20988x_3 \right) + 492660e^{18t\mu_A}.
\end{aligned}$$

$$\begin{aligned}
\phi_{77}^1 &= -\frac{1}{5}x_1x_3 + e^{3t\mu_A} \left(-\frac{12t_{3,2}}{5} - \frac{12}{25}t_{3,3}t_{3,4} - \frac{2t_{3,4}^2}{125} - \frac{18}{5}t_{3,3}x_3 - \frac{3}{25}t_{3,4}^2x_3 - \frac{12}{5}t_{3,4}x_3^2 - 6x_3^3 \right) + e^{5t\mu_A} \left(\frac{28}{5}t_{2,2}t_{3,4} \right. \\
&\quad \left. + \frac{24}{5}t_{3,4}x_2 + 26t_{2,2}x_3 + 12x_2x_3 \right) + e^{6t\mu_A} (36t_{1,1} + 32x_1) + e^{9t\mu_A} \left(102t_{3,3} - \frac{369t_{3,4}^2}{5} - 90t_{3,4}x_3 + 750x_3^2 \right) \\
&\quad + e^{11t\mu_A} (-1854t_{2,2} - 1260x_2) + e^{15t\mu_A} \left(\frac{126678t_{3,4}}{5} - 12474x_3 \right) - 1220640e^{21t\mu_A}, \\
\phi_{77}^2 &= e^{t\mu_A} \left(-\frac{t_{1,1}}{5} - \frac{2x_1}{5} \right) + e^{4t\mu_A} \left(-\frac{12t_{3,3}}{5} + \frac{14t_{3,4}^2}{25} + \frac{8}{5}t_{3,4}x_3 + 4x_3^2 \right) + \frac{44}{5}e^{6t\mu_A}t_{2,2} \\
&\quad + e^{10t\mu_A} (-172t_{3,4} - 420x_3) + 16884e^{16t\mu_A}, \\
\phi_{77}^3 &= \frac{x_3^2}{5} + e^{2t\mu_A} \left(-\frac{4t_{2,2}}{5} - \frac{6x_2}{5} \right) + e^{6t\mu_A} (12t_{3,4} - 8x_3) - \frac{4104}{5}e^{12t\mu_A}.
\end{aligned}$$

$$\begin{aligned}
\phi_{78}^1 &= -\frac{1}{25}t_{3,4}x_1x_3 - \frac{1}{5}x_1x_3^2 + e^{2t\mu_A} \left(-\frac{2}{5}t_{1,1}t_{2,2} - \frac{3}{5}t_{1,1}x_2 - \frac{6}{5}x_1x_2 \right) + e^{3t\mu_A} \left(\frac{6t_{3,3}^2}{25} - \frac{6}{25}t_{3,2}t_{3,4} - \frac{9}{125}t_{3,3}t_{3,4}^2 \right. \\
&\quad \left. - \frac{4t_{3,4}^4}{625} + \frac{6}{25}t_{3,3}t_{3,4}x_3 - \frac{3}{125}t_{3,4}^3x_3 + \frac{6}{5}t_{3,3}x_3^2 \right) + e^{5t\mu_A} \left(-\frac{3}{5}t_{2,2}t_{3,3} + \frac{11}{10}t_{2,2}t_{3,4}^2 + \frac{27}{5}t_{3,3}x_2 + \frac{159}{50}t_{3,4}^2x_2 \right. \\
&\quad \left. + \frac{24}{5}t_{2,2}t_{3,4}x_3 + \frac{54}{5}t_{3,4}x_2x_3 + 7t_{2,2}x_3^2 + 15x_2x_3^2 \right) + e^{6t\mu_A} \left(\frac{84}{5}t_{1,1}t_{3,4} + \frac{96}{5}t_{3,4}x_1 + 15t_{1,1}x_3 + 47x_1x_3 \right) \\
&\quad + e^{7t\mu_A} \left(\frac{3t_{2,2}^2}{5} - \frac{63}{5}t_{2,2}x_2 \right) + e^{9t\mu_A} \left(87t_{3,2} - \frac{384}{5}t_{3,3}t_{3,4} - \frac{866t_{3,4}^3}{25} - 156t_{3,3}x_3 - \frac{849}{5}t_{3,4}^2x_3 - 228t_{3,4}x_3^2 + 105x_3^3 \right) \\
&\quad + e^{11t\mu_A} \left(-\frac{2531}{5}t_{2,2}t_{3,4} - 1365t_{3,4}x_2 - 1670t_{2,2}x_3 - 2370x_2x_3 \right) + e^{12t\mu_A} \left(-\frac{14202t_{1,1}}{5} - 4000x_1 \right) \\
&\quad + e^{15t\mu_A} \left(\frac{68847t_{3,3}}{5} + \frac{217941t_{3,4}^2}{10} + \frac{287124}{5}t_{3,4}x_3 + 8757x_3^2 \right) + e^{17t\mu_A} \left(\frac{404558t_{2,2}}{5} + 141738x_2 \right) \\
&\quad + e^{21t\mu_A} (-3840648t_{3,4} - 3069810x_3) + 175921251e^{27t\mu_A},
\end{aligned}$$

$$\begin{aligned}
\phi_{78}^2 &= e^{t\mu_A} \left(-\frac{1}{25}t_{1,1}t_{3,4} - \frac{2}{25}t_{3,4}x_1 - \frac{1}{5}t_{1,1}x_3 - \frac{2}{5}x_1x_3 \right) + e^{2t\mu_A} \left(-\frac{2t_{2,1}}{5} - \frac{t_{2,2}^2}{15} \right) + e^{4t\mu_A} \left(-2t_{3,2} + \frac{16}{25}t_{3,3}t_{3,4} \right. \\
&\quad \left. + \frac{22t_{3,4}^3}{375} + \frac{22}{25}t_{3,4}^2x_3 - 4x_3^3 \right) + e^{6t\mu_A} \left(\frac{174}{25}t_{2,2}t_{3,4} + \frac{54}{5}t_{3,4}x_2 + \frac{114}{5}t_{2,2}x_3 \right) + e^{7t\mu_A} \left(\frac{353t_{1,1}}{5} + 94x_1 \right) \\
&\quad + e^{10t\mu_A} \left(-298t_{3,3} - \frac{1309t_{3,4}^2}{5} - 572t_{3,4}x_3 + 290x_3^2 \right) + e^{12t\mu_A} (-2134t_{2,2} - 2025x_2) + e^{16t\mu_A} (74992t_{3,4} + 30398x_3) \\
&\quad - 4046580e^{22t\mu_A}, \\
\phi_{78}^3 &= \frac{1}{25}t_{3,4}x_3^2 + \frac{x_3^3}{5} + e^{2t\mu_A} \left(-\frac{4}{25}t_{2,2}t_{3,4} - \frac{6}{25}t_{3,4}x_2 - \frac{4}{5}t_{2,2}x_3 \right) + e^{3t\mu_A} \left(-\frac{12t_{1,1}}{5} - \frac{12x_1}{5} \right) + e^{6t\mu_A} \left(\frac{54t_{3,3}}{5} \right. \\
&\quad \left. + \frac{147t_{3,4}^2}{25} + \frac{66}{5}t_{3,4}x_3 - 17x_3^2 \right) + e^{8t\mu_A} \left(\frac{358t_{2,2}}{5} + 66x_2 \right) + e^{12t\mu_A} \left(-\frac{54654t_{3,4}}{25} - \frac{3079x_3}{5} \right) + 123324e^{18t\mu_A}. \\
\phi_{79}^1 &= -\frac{1}{5}t_{3,3}x_1x_3 - \frac{1}{50}t_{3,4}^2x_1x_3 - \frac{2}{5}t_{3,4}x_1x_3^2 - x_1x_3^3 + e^{2t\mu_A} \left(-\frac{4}{5}t_{1,1}t_{2,2}t_{3,4} - \frac{6}{5}t_{1,1}t_{3,4}x_2 - \frac{6}{5}t_{3,4}x_1x_2 - 2t_{1,1}t_{2,2}x_3 \right. \\
&\quad \left. - 3t_{2,2}x_1x_3 - 3t_{1,1}x_2x_3 - 12x_1x_2x_3 \right) + e^{3t\mu_A} \left(-3t_{1,1}^2 - \frac{6}{5}t_{3,2}t_{3,3} - \frac{12}{25}t_{3,3}^2t_{3,4} - \frac{3}{25}t_{3,2}t_{3,4}^2 - \frac{2}{25}t_{3,3}t_{3,4}^3 - \frac{2t_{3,4}^5}{625} \right. \\
&\quad \left. + 8t_{2,2}^2x_2 + 24t_{2,2}x_2^2 + 18x_2^3 - \frac{18}{5}t_{3,3}^2x_3 + \frac{12}{5}t_{3,2}t_{3,4}x_3 - \frac{12}{25}t_{3,3}t_{3,4}^2x_3 - \frac{11}{250}t_{3,4}^4x_3 + 6t_{3,2}x_3^2 - \frac{12}{5}t_{3,3}t_{3,4}x_3^2 \right. \\
&\quad \left. - \frac{8}{25}t_{3,4}^3x_3^2 - 6t_{3,3}x_3^3 - \frac{3}{5}t_{3,4}^2x_3^3 \right) + e^{5t\mu_A} \left(2t_{2,2}t_{3,2} + \frac{36}{5}t_{2,2}t_{3,3}t_{3,4} + \frac{112}{75}t_{2,2}t_{3,4}^3 + 12t_{3,2}x_2 + \frac{72}{5}t_{3,3}t_{3,4}x_2 \right. \\
&\quad \left. + \frac{116}{25}t_{3,4}^3x_2 + 52t_{2,2}t_{3,3}x_3 + 12t_{2,2}t_{3,4}^2x_3 + 24t_{3,3}x_2x_3 + \frac{144}{5}t_{3,4}^2x_2x_3 + 16t_{2,2}t_{3,4}x_3^2 + 24t_{3,4}x_2x_3^2 - 10t_{2,2}x_3^3 - 60x_2x_3^3 \right) \\
&\quad + e^{6t\mu_A} \left(96t_{1,1}t_{3,3} + \frac{168}{5}t_{1,1}t_{3,4}^2 + 96t_{3,3}x_1 + \frac{144}{5}t_{3,4}^2x_1 + 90t_{1,1}t_{3,4}x_3 + 257t_{3,4}x_1x_3 + 60t_{1,1}x_3^2 + 300x_1x_3^2 \right) \\
&\quad + e^{7t\mu_A} \left(-\frac{84}{5}t_{2,2}^2t_{3,4} - \frac{1656}{5}t_{2,2}t_{3,4}x_2 - 432t_{3,4}x_2^2 + 118t_{2,2}^2x_3 + 312t_{2,2}x_2x_3 \right) + e^{8t\mu_A} (424t_{1,1}t_{2,2} + 480t_{2,2}x_1 \\
&\quad + 780t_{1,1}x_2 + 1575x_1x_2) + e^{9t\mu_A} \left(-264t_{3,3}^2 - 6t_{3,2}t_{3,4} - \frac{1152}{5}t_{3,3}t_{3,4}^2 - \frac{1456t_{3,4}^4}{25} - 720t_{3,2}x_3 - 516t_{3,3}t_{3,4}x_3 \right. \\
&\quad \left. - 384t_{3,4}^3x_3 - 120t_{3,3}x_3^2 - 24t_{3,4}^2x_3^2 + 1590t_{3,4}x_3^3 \right) + e^{11t\mu_A} \left(-5028t_{2,2}t_{3,3} - \frac{11004}{5}t_{2,2}t_{3,4}^2 - 7362t_{3,3}x_2 \right. \\
&\quad \left. - \frac{14571}{5}t_{3,4}^2x_2 - 15608t_{2,2}t_{3,4}x_3 - 15744t_{3,4}x_2x_3 - 4060t_{2,2}x_3^2 - 510x_2x_3^2 \right) + e^{12t\mu_A} \left(-\frac{120024}{5}t_{1,1}t_{3,4} - 36000t_{3,4}x_1 \right. \\
&\quad \left. - 12912t_{1,1}x_3 - 41388x_1x_3 \right) + e^{13t\mu_A} (-6918t_{2,2}^2 - 3294t_{2,2}x_2 + 16524x_2^2) + e^{15t\mu_A} \left(14292t_{3,2} + \frac{873846}{5}t_{3,3}t_{3,4} \right. \\
&\quad \left. + \frac{2109309t_{3,4}^3}{25} + 87942t_{3,3}x_3 + 286941t_{3,4}^2x_3 - 44814t_{3,4}x_3^2 - 168660x_3^3 \right) + e^{17t\mu_A} \left(\frac{5938786}{5}t_{2,2}t_{3,4} + 1473906t_{3,4}x_2 \right. \\
&\quad \left. + 1595538t_{2,2}x_3 + 1224180x_2x_3 \right) + e^{18t\mu_A} (2077020t_{1,1} + 3564000x_1) + e^{21t\mu_A} (-14782620t_{3,3} - 31873638t_{3,4}^2 \\
&\quad - 42125310t_{3,4}x_3 + 8344500x_3^2) + e^{23t\mu_A} (-97157950t_{2,2} - 110617962x_2) + e^{27t\mu_A} (3797143338t_{3,4} + 1547617680x_3) \\
&\quad - 136685952000e^{33t\mu_A},
\end{aligned}$$

$$\begin{aligned}
\phi_{79}^2 = & e^{t\mu_A} \left(-\frac{1}{5}t_{1,1}t_{3,3} - \frac{1}{50}t_{1,1}t_{3,4}^2 - \frac{2}{5}t_{3,3}x_1 - \frac{1}{25}t_{3,4}^2x_1 - \frac{2}{5}t_{1,1}t_{3,4}x_3 - \frac{4}{5}t_{3,4}x_1x_3 - t_{1,1}x_3^2 - 2x_1x_3^2 \right) \\
& + e^{2t\mu_A} \left(-\frac{4}{5}t_{2,1}t_{3,4} - \frac{2}{15}t_{2,2}^2t_{3,4} - 2t_{2,1}x_3 - \frac{1}{3}t_{2,2}^2x_3 + 4t_{2,2}x_2x_3 + 6x_2^2x_3 \right) + e^{3t\mu_A} (-7t_{1,1}t_{2,2} - 6t_{2,2}x_1 - 6t_{1,1}x_2 \\
& - 16x_1x_2) + e^{4t\mu_A} \left(-\frac{4}{5}t_{3,3}^2 + \frac{12}{5}t_{3,2}t_{3,4} + \frac{16}{25}t_{3,3}t_{3,4}^2 + \frac{t_{3,4}^4}{25} + 18t_{3,2}x_3 + \frac{16}{5}t_{3,3}t_{3,4}x_3 + \frac{6}{5}t_{3,4}^3x_3 + 20t_{3,3}x_3^2 \right. \\
& \left. - \frac{6}{5}t_{3,4}^2x_3^2 - 12t_{3,4}x_3^3 \right) + e^{6t\mu_A} \left(\frac{184}{5}t_{2,2}t_{3,3} + \frac{732}{25}t_{2,2}t_{3,4}^2 + 48t_{3,3}x_2 + \frac{288}{5}t_{3,4}^2x_2 + \frac{458}{5}t_{2,2}t_{3,4}x_3 - 36t_{3,4}x_2x_3 \right. \\
& \left. + 44t_{2,2}x_3^2 \right) + e^{7t\mu_A} \left(\frac{1561}{5}t_{1,1}t_{3,4} + 514t_{3,4}x_1 + 228t_{1,1}x_3 + 600x_1x_3 \right) + e^{8t\mu_A} (120t_{2,1} + 128t_{2,2}^2 + 144t_{2,2}x_2) \\
& + e^{10t\mu_A} \left(-780t_{3,2} - 1956t_{3,3}t_{3,4} - \frac{5166t_{3,4}^3}{5} - 2280t_{3,3}x_3 - 2608t_{3,4}^2x_3 + 260t_{3,4}x_3^2 + 3750x_3^3 \right) \\
& + e^{12t\mu_A} (-18148t_{2,2}t_{3,4} - 23604t_{3,4}x_2 - 25060t_{2,2}x_3 - 3996x_2x_3) + e^{13t\mu_A} (-46692t_{1,1} - 82776x_1) \\
& + e^{16t\mu_A} \left(300708t_{3,3} + \frac{2668822t_{3,4}^2}{5} + 677214t_{3,4}x_3 - 257400x_3^2 \right) + e^{18t\mu_A} (2113848t_{2,2} + 2138400x_2) \\
& + e^{22t\mu_A} (-80787900t_{3,4} - 28500750x_3) + 3363095520e^{28t\mu_A}, \\
\phi_{79}^3 = & e^{2t\mu_A} \left(-\frac{4}{5}t_{2,2}t_{3,3} - \frac{2}{25}t_{2,2}t_{3,4}^2 - \frac{6}{5}t_{3,3}x_2 - \frac{3}{25}t_{3,4}^2x_2 - \frac{4}{5}t_{2,2}t_{3,4}x_3 - 2t_{2,2}x_3^2 \right) + e^{3t\mu_A} \left(-\frac{18}{5}t_{1,1}t_{3,4} - \frac{12}{5}t_{3,4}x_1 \right. \\
& \left. - 6t_{1,1}x_3 \right) + e^{4t\mu_A} \left(-16t_{2,1} + \frac{4t_{2,2}^2}{3} - 2t_{2,2}x_2 - 12x_2^2 \right) + e^{6t\mu_A} \left(24t_{3,2} + \frac{156}{5}t_{3,3}t_{3,4} + \frac{142t_{3,4}^3}{25} - 24t_{3,3}x_3 + \frac{108}{5}t_{3,4}^2x_3 \right. \\
& \left. - 78t_{3,4}x_3^2 - 300x_3^3 \right) + e^{8t\mu_A} \left(\frac{1672}{5}t_{2,2}t_{3,4} + 678t_{3,4}x_2 + 1224t_{2,2}x_3 \right) + e^{9t\mu_A} (2559t_{1,1} + 3150x_1) + e^{12t\mu_A} \left(-\frac{65124t_{3,3}}{5} \right. \\
& \left. - \frac{289362t_{3,4}^2}{25} - \frac{67224}{5}t_{3,4}x_3 + 32988x_3^2 \right) + e^{14t\mu_A} (-101052t_{2,2} - 101232x_2) + e^{18t\mu_A} (2974884t_{3,4} - 24060x_3) \\
& - 146515476e^{24t\mu_A}.
\end{aligned}$$

$$\begin{aligned}
\phi_{88}^1 = & -\frac{1}{25}t_{3,3}x_1x_3 - \frac{1}{250}t_{3,4}^2x_1x_3 - \frac{2}{25}t_{3,4}x_1x_3^2 - \frac{1}{5}x_1x_3^3 + e^{2t\mu_A} \left(-\frac{4}{25}t_{1,1}t_{2,2}t_{3,4} - \frac{6}{25}t_{1,1}t_{3,4}x_2 - \frac{12}{25}t_{3,4}x_1x_2 \right. \\
& \left. - \frac{2}{5}t_{1,1}t_{2,2}x_3 - \frac{3}{5}t_{2,2}x_1x_3 - \frac{3}{5}t_{1,1}x_2x_3 - \frac{12}{5}x_1x_2x_3 \right) + e^{3t\mu_A} \left(-\frac{3}{5}t_{1,1}^2 - \frac{12}{125}t_{3,3}^2t_{3,4} + \frac{6}{125}t_{3,2}t_{3,4}^2 - \frac{6}{625}t_{3,3}t_{3,4}^3 \right. \\
& \left. - \frac{2t_{3,4}^5}{3125} + \frac{8}{5}t_{2,2}^2x_2 + \frac{24}{5}t_{2,2}x_2^2 + \frac{18x_2^3}{5} + \frac{6}{25}t_{3,3}^2x_3 + \frac{12}{125}t_{3,3}t_{3,4}^2x_3 + \frac{21t_{3,4}^4x_3}{1250} + \frac{12}{25}t_{3,3}t_{3,4}x_3^2 + \frac{24}{125}t_{3,4}^3x_3^2 \right. \\
& \left. + \frac{6}{5}t_{3,3}x_3^3 + \frac{9}{25}t_{3,4}^2x_3^3 \right) + e^{5t\mu_A} \left(-\frac{6}{5}t_{2,2}t_{3,2} + \frac{26}{25}t_{2,2}t_{3,3}t_{3,4} + \frac{17}{125}t_{2,2}t_{3,4}^3 - \frac{18}{5}t_{3,2}x_2 + \frac{102}{25}t_{3,3}t_{3,4}x_2 \right. \\
& \left. + \frac{141}{125}t_{3,4}^3x_2 + \frac{16}{5}t_{2,2}t_{3,3}x_3 + \frac{96}{25}t_{2,2}t_{3,4}^2x_3 + 12t_{3,3}x_2x_3 + \frac{36}{5}t_{3,4}^2x_2x_3 + \frac{66}{5}t_{2,2}t_{3,4}x_3^2 + 18t_{3,4}x_2x_3^2 + 12t_{2,2}x_3^3 + 18x_2x_3^3 \right) \\
& + e^{6t\mu_A} \left(\frac{12}{5}t_{1,1}t_{3,3} + \frac{48}{5}t_{1,1}t_{3,4}^2 + \frac{288}{25}t_{3,4}^2x_1 + \frac{144}{5}t_{1,1}t_{3,4}x_3 + \frac{343}{5}t_{3,4}x_1x_3 + 18t_{1,1}x_3^2 + 62x_1x_3^2 \right) \\
& + e^{7t\mu_A} \left(-\frac{414}{25}t_{2,2}^2t_{3,4} - \frac{2826}{25}t_{2,2}t_{3,4}x_2 - \frac{648}{5}t_{3,4}x_2^2 - \frac{42}{5}t_{2,2}^2x_3 - \frac{108}{5}t_{2,2}x_2x_3 \right) + e^{8t\mu_A} \left(\frac{104}{5}t_{1,1}t_{2,2} + 6t_{1,1}x_2 \right. \\
& \left. + 129x_1x_2 \right) + e^{9t\mu_A} \left(-\frac{36}{5}t_{3,3}^2 + \frac{294}{5}t_{3,2}t_{3,4} - \frac{1356}{25}t_{3,3}t_{3,4}^2 - \frac{499t_{3,4}^4}{25} - \frac{1572}{5}t_{3,3}t_{3,4}x_3 - \frac{4218}{25}t_{3,4}^3x_3 - 300t_{3,3}x_3^2 \right. \\
& \left. - 486t_{3,4}^2x_3^2 - 468t_{3,4}x_3^3 \right) + e^{11t\mu_A} \left(-\frac{1338}{5}t_{2,2}t_{3,3} + \frac{1581}{5}t_{2,2}t_{3,4}^2 - 936t_{3,3}x_2 + 702t_{3,4}^2x_2 - \frac{9536}{5}t_{2,2}t_{3,4}x_3 \right. \\
& \left. - 3084t_{3,4}x_2x_3 - 3202t_{2,2}x_3^2 - 3840x_2x_3^2 \right) + e^{12t\mu_A} \left(-\frac{64104}{25}t_{1,1}t_{3,4} - 4800t_{3,4}x_1 - \frac{23952}{5}t_{1,1}x_3 - \frac{57378}{5}x_1x_3 \right)
\end{aligned}$$

$$\begin{aligned}
& + e^{13t_{\mu A}} \left(\frac{12132t_{2,2}^2}{5} + \frac{60246}{5}t_{2,2}x_2 + \frac{57024x_2^2}{5} \right) + e^{15t_{\mu A}} \left(-\frac{35256t_{3,2}}{5} + \frac{584526}{25}t_{3,3}t_{3,4} + \frac{1315567t_{3,4}^3}{125} \right. \\
& + \frac{244866}{5}t_{3,3}x_3 + \frac{2187471}{25}t_{3,4}^2x_3 + \frac{691266}{5}t_{3,4}x_3^2 + 39012x_3^3 \Big) + e^{17t_{\mu A}} \left(-\frac{967924}{25}t_{2,2}t_{3,4} - \frac{459714}{5}t_{3,4}x_2 \right. \\
& + \frac{1304718}{5}t_{2,2}x_3 + 323604x_2x_3 \Big) + e^{18t_{\mu A}} (215772t_{1,1} + 500000x_1) + e^{21t_{\mu A}} \left(-2200020t_{3,3} - \frac{15303492t_{3,4}^2}{5} \right. \\
& - 14407686t_{3,4}x_3 - 8280720x_3^2 \Big) + e^{23t_{\mu A}} \left(-2026918t_{2,2} - \frac{3189792x_2}{5} \right) + e^{27t_{\mu A}} \left(\frac{2026944318t_{3,4}}{5} + 675234492x_3 \right) \\
& - 17094510720e^{33t_{\mu A}}, \\
\phi_{88}^2 = & e^{t_{\mu A}} \left(-\frac{1}{25}t_{1,1}t_{3,3} - \frac{1}{250}t_{1,1}t_{3,4}^2 - \frac{2}{25}t_{3,3}x_1 - \frac{1}{125}t_{3,4}^2x_1 - \frac{2}{25}t_{1,1}t_{3,4}x_3 - \frac{4}{25}t_{3,4}x_1x_3 - \frac{1}{5}t_{1,1}x_3^2 - \frac{2}{5}x_1x_3^2 \right) \\
& + e^{2t_{\mu A}} \left(-\frac{4}{25}t_{2,1}t_{3,4} - \frac{2}{75}t_{2,2}^2t_{3,4} - \frac{2}{5}t_{2,1}x_3 - \frac{1}{15}t_{2,2}^2x_3 + \frac{4}{5}t_{2,2}x_2x_3 + \frac{6}{5}x_2^2x_3 \right) + e^{3t_{\mu A}} \left(-\frac{7}{5}t_{1,1}t_{2,2} - \frac{6}{5}t_{2,2}x_1 \right. \\
& - \frac{6}{5}t_{1,1}x_2 - \frac{24}{5}x_1x_2 \Big) + e^{4t_{\mu A}} \left(\frac{4t_{3,3}^2}{5} + \frac{12}{25}t_{3,2}t_{3,4} + \frac{16}{125}t_{3,3}t_{3,4}^2 + \frac{11t_{3,4}^4}{625} + \frac{6}{5}t_{3,2}x_3 + \frac{76}{25}t_{3,3}t_{3,4}x_3 \right. \\
& + \frac{64}{125}t_{3,4}^3x_3 + 4t_{3,3}x_3^2 + \frac{78}{25}t_{3,4}^2x_3^2 + \frac{24}{5}t_{3,4}x_3^3 \Big) + e^{6t_{\mu A}} \left(\frac{304}{25}t_{2,2}t_{3,3} + \frac{252}{125}t_{2,2}t_{3,4}^2 + \frac{156}{5}t_{3,3}x_2 + \frac{36}{5}t_{3,4}^2x_2 \right. \\
& + \frac{258}{25}t_{2,2}t_{3,4}x_3 - \frac{108}{5}t_{3,4}x_2x_3 + \frac{144}{5}t_{2,2}x_3^2 \Big) + e^{7t_{\mu A}} \left(\frac{1811}{25}t_{1,1}t_{3,4} + \frac{686}{5}t_{3,4}x_1 + \frac{538}{5}t_{1,1}x_3 + 124x_1x_3 \right) \\
& + e^{8t_{\mu A}} \left(124t_{2,1} + \frac{274t_{2,2}^2}{15} + \frac{684}{5}t_{2,2}x_2 + 243x_2^2 \right) + e^{10t_{\mu A}} \left(-256t_{3,2} - \frac{4604}{5}t_{3,3}t_{3,4} - \frac{15854t_{3,4}^3}{75} - 1212t_{3,3}x_3 \right. \\
& - 930t_{3,4}^2x_3 - 1928t_{3,4}x_3^2 - 710x_3^3 \Big) + e^{12t_{\mu A}} \left(-\frac{16552}{5}t_{2,2}t_{3,4} - 7896t_{3,4}x_2 - 4452t_{2,2}x_3 + \frac{9504}{5}x_2x_3 \right) \\
& + e^{13t_{\mu A}} \left(-\frac{65962t_{1,1}}{5} - \frac{114756x_1}{5} \right) + e^{16t_{\mu A}} \left(\frac{660536t_{3,3}}{5} + \frac{3543388t_{3,4}^2}{25} + \frac{1320902}{5}t_{3,4}x_3 + 161904x_3^2 \right) \\
& + e^{18t_{\mu A}} \left(\frac{2245908t_{2,2}}{5} + 705888x_2 \right) + e^{22t_{\mu A}} (-23681508t_{3,4} - 15860810x_3) + 1067265624e^{28t_{\mu A}}, \\
\phi_{88}^3 = & \frac{1}{25}t_{3,3}x_3^2 + \frac{1}{250}t_{3,4}^2x_3^2 + \frac{2}{25}t_{3,4}x_3^3 + \frac{x_3^4}{5} + e^{2t_{\mu A}} \left(-\frac{4}{25}t_{2,2}t_{3,3} - \frac{2}{125}t_{2,2}t_{3,4}^2 - \frac{6}{25}t_{3,3}x_2 - \frac{3}{125}t_{3,4}^2x_2 \right. \\
& - \frac{8}{25}t_{2,2}t_{3,4}x_3 - t_{2,2}x_3^2 \Big) + e^{3t_{\mu A}} \left(-\frac{24}{25}t_{1,1}t_{3,4} - \frac{24}{25}t_{3,4}x_1 - \frac{12}{5}t_{1,1}x_3 \right) + e^{4t_{\mu A}} \left(-\frac{24t_{2,1}}{5} - \frac{18}{5}t_{2,2}x_2 - \frac{36x_2^2}{5} \right) \\
& + e^{6t_{\mu A}} \left(\frac{36t_{3,2}}{5} + \frac{264}{25}t_{3,3}t_{3,4} + \frac{48t_{3,4}^3}{25} + 12t_{3,3}x_3 + \frac{252}{25}t_{3,4}^2x_3 + \frac{53}{5}t_{3,4}x_3^2 - 26x_3^3 \right) + e^{8t_{\mu A}} \left(\frac{1316}{25}t_{2,2}t_{3,4} \right. \\
& + \frac{762}{5}t_{3,4}x_2 + \frac{628}{5}t_{2,2}x_3 \Big) + e^{9t_{\mu A}} \left(\frac{1281t_{1,1}}{5} + 258x_1 \right) + e^{12t_{\mu A}} \left(-\frac{49104t_{3,3}}{25} - \frac{240552t_{3,4}^2}{125} - \frac{83208}{25}t_{3,4}x_3 - 6x_3^2 \right) \\
& + e^{14t_{\mu A}} \left(-\frac{44412t_{2,2}}{5} - \frac{74268x_2}{5} \right) + e^{18t_{\mu A}} \left(\frac{1815468t_{3,4}}{5} + 173128x_3 \right) - \frac{83805096}{5}e^{24t_{\mu A}}.
\end{aligned}$$

$$\begin{aligned}
\phi_{89}^1 = & -\frac{1}{5}t_{3,2}x_1x_3 - \frac{1}{25}t_{3,3}t_{3,4}x_1x_3 - \frac{1}{750}t_{3,4}^3x_1x_3 - \frac{2}{5}t_{3,3}x_1x_3^2 - \frac{2}{25}t_{3,4}^2x_1x_3^2 - \frac{3}{5}t_{3,4}x_1x_3^3 - x_1x_3^4 \\
& + e^{2t_{\mu A}} \left(-\frac{4}{5}t_{1,1}t_{2,2}t_{3,3} - \frac{4}{25}t_{1,1}t_{2,2}t_{3,4}^2 - \frac{6}{5}t_{1,1}t_{3,3}x_2 - \frac{6}{25}t_{1,1}t_{3,4}^2x_2 - \frac{6}{5}t_{3,3}x_1x_2 - \frac{6}{25}t_{3,4}^2x_1x_2 - \frac{6}{5}t_{1,1}t_{2,2}t_{3,4}x_3 \right. \\
& - \frac{9}{5}t_{2,2}t_{3,4}x_1x_3 - \frac{9}{5}t_{1,1}t_{3,4}x_2x_3 - \frac{36}{5}t_{3,4}x_1x_2x_3 - 2t_{1,1}t_{2,2}x_3^2 - 2t_{2,2}x_1x_3^2 - 3t_{1,1}x_2x_3^2 - 12x_1x_2x_3^2 \Big) + e^{3t_{\mu A}} \left(-\frac{6}{5}t_{3,2}^2 \right. \\
& - \frac{9}{5}t_{1,1}^2t_{3,4} - \frac{18}{25}t_{3,2}t_{3,3}t_{3,4} - \frac{12}{125}t_{3,3}^2t_{3,4}^2 - \frac{1}{25}t_{3,2}t_{3,4}^3 - \frac{6}{625}t_{3,3}t_{3,4}^4 - \frac{2t_{3,4}^6}{9375} + \frac{24}{5}t_{2,2}^2t_{3,4}x_2 + \frac{72}{5}t_{2,2}t_{3,4}x_2^2 + \frac{54}{5}t_{3,4}x_2^3 \\
& - 3t_{1,1}^2x_3 - \frac{6}{5}t_{3,2}t_{3,3}x_3 + \frac{6}{25}t_{3,3}^2t_{3,4}x_3 - \frac{3}{5}t_{3,2}t_{3,4}^2x_3 + \frac{2}{125}t_{3,3}t_{3,4}^3x_3 + \frac{7t_{3,4}^5x_3}{1250} - 4t_{1,1}x_1x_3 + 8t_{2,2}^2x_2x_3 + 24t_{2,2}x_2^2x_3 \\
& + 18x_2^3x_3 - \frac{6}{5}t_{3,2}t_{3,4}x_3^2 + \frac{12}{25}t_{3,3}t_{3,4}^2x_3^2 + \frac{8}{125}t_{3,4}^4x_3^2 + \frac{6}{5}t_{3,3}t_{3,4}^3x_3^2 + \frac{3}{25}t_{3,4}^3x_3^3 \Big) + e^{4t_{\mu A}} \left(-4t_{1,1}t_{2,2}^2 + 6t_{1,1}t_{2,2}x_2 \right. \\
& + 3t_{2,2}x_1x_2 + 18t_{1,1}x_2^2 - 9x_1x_2^2 \Big) + e^{5t_{\mu A}} \left(4t_{2,2}t_{3,3}^2 + \frac{18}{5}t_{2,2}t_{3,2}t_{3,4} + \frac{74}{25}t_{2,2}t_{3,3}t_{3,4}^2 + \frac{36}{125}t_{2,2}t_{3,4}^4 + \frac{84}{5}t_{3,3}^2x_2 \right. \\
& + \frac{246}{25}t_{3,3}t_{3,4}^2x_2 + \frac{144}{125}t_{3,4}^4x_2 - 6t_{2,2}t_{3,2}x_3 + \frac{144}{5}t_{2,2}t_{3,3}t_{3,4}x_3 + \frac{236}{25}t_{2,2}t_{3,4}^3x_3 + \frac{288}{5}t_{3,3}t_{3,4}x_2x_3 + \frac{72}{5}t_{3,4}^3x_2x_3 \\
& + 28t_{2,2}t_{3,3}x_3^2 + \frac{216}{5}t_{2,2}t_{3,4}^2x_3^2 + 60t_{3,3}x_2x_3^2 + \frac{288}{5}t_{3,4}^2x_2x_3^2 + 54t_{2,2}t_{3,4}x_3^3 + 72t_{3,4}x_2x_3^3 \Big) + e^{6t_{\mu A}} \left(6t_{1,1}t_{3,2} \right. \\
& + \frac{288}{5}t_{1,1}t_{3,3}t_{3,4} + \frac{412}{25}t_{1,1}t_{3,4}^3 + \frac{288}{5}t_{3,3}t_{3,4}x_1 + \frac{432}{25}t_{3,4}^3x_1 + 105t_{1,1}t_{3,3}x_3 + \frac{1023}{10}t_{1,1}t_{3,4}^2x_3 + 161t_{3,3}x_1x_3 \\
& + \frac{1969}{10}t_{3,4}^2x_1x_3 + 210t_{1,1}t_{3,4}^2x_3^2 + 442t_{3,4}x_1x_3^2 + 135t_{1,1}x_3^3 + 375x_1x_3^3 \Big) + e^{7t_{\mu A}} \left(\frac{226}{5}t_{2,2}^2t_{3,3} - \frac{1089}{25}t_{2,2}^2t_{3,4}^2 \right. \\
& + \frac{264}{5}t_{2,2}t_{3,3}x_2 - \frac{8226}{25}t_{2,2}t_{3,4}^2x_2 + 72t_{3,3}x_2^2 - \frac{1944}{5}t_{3,4}^2x_2^2 - \frac{6}{5}t_{2,2}^2t_{3,4}x_3 - \frac{2304}{5}t_{2,2}t_{3,4}x_2x_3 - 648t_{3,4}x_2^2x_3 + 198t_{2,2}^2x_3^2 \\
& + 252t_{2,2}x_2^3 \Big) + e^{8t_{\mu A}} \left(\frac{1432}{5}t_{1,1}t_{2,2}t_{3,4} + 288t_{2,2}t_{3,4}x_1 + 138t_{1,1}t_{3,4}x_2 + 1065t_{3,4}x_1x_2 + 639t_{1,1}t_{2,2}x_3 \right. \\
& + 1026t_{2,2}x_1x_3 + 1125t_{1,1}x_2x_3 + 4500x_1x_2x_3 \Big) + e^{9t_{\mu A}} \left(105t_{1,1}^2 - 144t_{2,2}^3 + 282t_{3,2}t_{3,3} - 312t_{3,3}^2t_{3,4} + 165t_{3,2}t_{3,4}^2 \right. \\
& - \frac{884}{5}t_{3,3}t_{3,4}^3 - \frac{3722t_{3,4}^5}{125} - 810t_{2,1}x_2 - 1749t_{2,2}^2x_2 - 5112t_{2,2}^2x_2^2 - 4320x_2^3 - 810t_{3,1}x_3 - 543t_{3,3}^2x_3 + 336t_{3,2}t_{3,4}x_3 \\
& - \frac{6867}{5}t_{3,3}t_{3,4}^2x_3 - \frac{7157}{20}t_{3,4}^4x_3 - 1410t_{3,2}x_3^2 - 2592t_{3,3}t_{3,4}x_3^2 - \frac{7064}{5}t_{3,4}^3x_3^2 - 3360t_{3,3}x_3^3 - 2394t_{3,4}^2x_3^3 - 3240t_{3,4}^4x_3^4 \\
& - 4050x_3^5 \Big) + e^{11t_{\mu A}} \left(-118t_{2,2}t_{3,2} - \frac{26846}{5}t_{2,2}t_{3,3}t_{3,4} + \frac{31369}{75}t_{2,2}t_{3,4}^3 + 2118t_{3,2}x_2 - \frac{50232}{5}t_{3,3}t_{3,4}x_2 + \frac{50168}{25}t_{3,4}^3x_2 \right. \\
& - 7068t_{2,2}t_{3,3}x_3 - \frac{46044}{5}t_{2,2}t_{3,4}^2x_3 - 16704t_{3,3}x_2x_3 - \frac{37944}{5}t_{3,4}^2x_2x_3 - 22822t_{2,2}t_{3,4}x_3^2 - 29256t_{3,4}x_2x_3^2 + 740t_{2,2}x_3^3 \\
& - 11760x_2x_3^3 \Big) + e^{12t_{\mu A}} \left(-\frac{50424}{5}t_{1,1}t_{3,3} - \frac{369624}{25}t_{1,1}t_{3,4}^2 - 12000t_{3,3}x_1 - 25200t_{3,4}^2x_1 - \frac{242061}{5}t_{1,1}t_{3,4}x_3 \right. \\
& - \frac{550014}{5}t_{3,4}x_1x_3 - 23112t_{1,1}x_3^2 - 77512x_1x_3^2 \Big) + e^{13t_{\mu A}} \left(9720t_{2,1}t_{3,4} + \frac{88006}{5}t_{2,2}^2t_{3,4} + \frac{660918}{5}t_{2,2}t_{3,4}x_2 \right. \\
& + \frac{849672}{5}t_{3,4}x_2^2 + 12150t_{2,1}x_3 - 15533t_{2,2}^2x_3 + 19536t_{2,2}x_2x_3 + 57024x_2^2x_3 \Big) + e^{14t_{\mu A}} \left(-33778t_{1,1}t_{2,2} - 60000t_{2,2}x_1 \right. \\
& - 30264t_{1,1}x_2 - 243192x_1x_2 \Big) + e^{15t_{\mu A}} \left(48600t_{3,1} + 56700t_{3,3}^2 - \frac{353112}{5}t_{3,2}t_{3,4} + \frac{4588314}{25}t_{3,3}t_{3,4}^2 + \frac{4335086t_{3,4}^4}{125} \right. \\
& + 115494t_{3,2}x_3 + \frac{2899374}{5}t_{3,3}t_{3,4}x_3 + \frac{8402551}{25}t_{3,4}^3x_3 + 634248t_{3,3}x_3^2 + \frac{4278966}{5}t_{3,4}^2x_3^2 + 1039614t_{3,4}x_3^3 + 1053000x_3^4 \Big)
\end{aligned}$$

$$\begin{aligned}
& + e^{17t_{\mu A}} \left(\frac{2736506}{5} t_{2,2} t_{3,3} - \frac{2065049}{25} t_{2,2} t_{3,4}^2 + 1223070 t_{3,3} x_2 - \frac{5167743}{5} t_{3,4}^2 x_2 + \frac{11899794}{5} t_{2,2} t_{3,4} x_3 \right. \\
& + 2418840 t_{3,4} x_2 x_3 + 873698 t_{2,2} x_3^2 + 2627220 x_2 x_3^2 \left. \right) + e^{18t_{\mu A}} (2880264 t_{1,1} t_{3,4} + 6638400 t_{3,4} x_1 + 3090885 t_{1,1} x_3 \\
& + 9691725 x_1 x_3) + e^{19t_{\mu A}} (-1117800 t_{2,1} - 688278 t_{2,2}^2 - 7162794 t_{2,2} x_2 - 9575280 x_2^2) + e^{21t_{\mu A}} \left(584970 t_{3,2} \right. \\
& - 38455356 t_{3,3} t_{3,4} - \frac{81542494 t_{3,4}^3}{5} - 57636180 t_{3,3} x_3 - 102794646 t_{3,4}^2 x_3 - 158851140 t_{3,4} x_3^2 - 131244600 x_3^3 \left. \right) \\
& + e^{23t_{\mu A}} \left(-28927128 t_{2,2} t_{3,4} + \frac{253873764}{5} t_{3,4} x_2 - 113054640 t_{2,2} x_3 - 177923952 x_2 x_3 \right) + e^{24t_{\mu A}} (-155768424 t_{1,1} \\
& - 445500000 x_1) + e^{27t_{\mu A}} \left(2424423000 t_{3,3} + \frac{18908795082 t_{3,4}^2}{5} + 12444036816 t_{3,4} x_3 + 10511125560 x_3^2 \right) \\
& + e^{29t_{\mu A}} (2803944506 t_{2,2} + 2255008080 x_2) + e^{33t_{\mu A}} (-388000760040 t_{3,4} - 545906954550 x_3) + 14162630484960 e^{39t_{\mu A}}, \\
\phi_{89}^2 = & e^{t_{\mu A}} \left(-\frac{1}{5} t_{1,1} t_{3,2} - \frac{1}{25} t_{1,1} t_{3,3} t_{3,4} - \frac{1}{750} t_{1,1} t_{3,4}^3 - \frac{2}{5} t_{3,2} x_1 - \frac{2}{25} t_{3,3} t_{3,4} x_1 - \frac{1}{375} t_{3,4}^3 x_1 - \frac{2}{5} t_{1,1} t_{3,3} x_3 \right. \\
& - \frac{2}{25} t_{1,1} t_{3,4}^2 x_3 - \frac{4}{5} t_{3,3} x_1 x_3 - \frac{4}{25} t_{3,4}^2 x_1 x_3 - \frac{3}{5} t_{1,1} t_{3,4} x_3^2 - \frac{6}{5} t_{3,4} x_1 x_3^2 - t_{1,1} x_3^3 - 2 x_1 x_3^3 \left. \right) + e^{2t_{\mu A}} \left(-\frac{4}{5} t_{2,1} t_{3,3} \right. \\
& - \frac{2}{15} t_{2,2}^2 t_{3,3} - \frac{4}{25} t_{2,1} t_{3,4}^2 - \frac{2}{75} t_{2,2}^2 t_{3,4}^2 - \frac{6}{5} t_{2,1} t_{3,4} x_3 - \frac{1}{5} t_{2,2}^2 t_{3,4} x_3 + \frac{12}{5} t_{2,2} t_{3,4} x_2 x_3 + \frac{18}{5} t_{3,4} x_2^2 x_3 - 2 t_{2,1} x_3^2 - \frac{1}{3} t_{2,2}^2 x_3^2 \\
& + 4 t_{2,2} x_2 x_3^2 + 6 x_2^2 x_3^2 \left. \right) + e^{3t_{\mu A}} \left(-\frac{21}{5} t_{1,1} t_{2,2} t_{3,4} - \frac{18}{5} t_{2,2} t_{3,4} x_1 - \frac{18}{5} t_{1,1} t_{3,4} x_2 - \frac{48}{5} t_{3,4} x_1 x_2 - 6 t_{1,1} t_{2,2} x_3 \right. \\
& - 4 t_{2,2} x_1 x_3 - 48 x_1 x_2 x_3 \left. \right) + e^{4t_{\mu A}} \left(-10 t_{1,1}^2 - 4 t_{2,1} t_{2,2} - \frac{2 t_{2,2}^3}{3} + \frac{8}{5} t_{3,2} t_{3,3} + \frac{4}{5} t_{3,3}^2 t_{3,4} + \frac{6}{25} t_{3,2} t_{3,4}^2 + \frac{76}{375} t_{3,3} t_{3,4}^3 \right. \\
& + \frac{7 t_{3,4}^5}{625} - 8 t_{1,1} x_1 + 24 t_{3,1} x_3 + 8 t_{3,3}^2 x_3 + 6 t_{3,2} t_{3,4} x_3 + \frac{142}{25} t_{3,3} t_{3,4}^2 x_3 + \frac{71}{125} t_{3,4}^4 x_3 + 54 t_{3,2} x_3^2 + \frac{168}{5} t_{3,3} t_{3,4} x_3^2 \\
& + \frac{138}{25} t_{3,4}^3 x_3^2 + 92 t_{3,3} x_3^3 + \frac{144}{5} t_{3,4}^2 x_3^3 + 96 t_{3,4} x_3^4 + 120 x_3^5 \left. \right) + e^{6t_{\mu A}} \left(\frac{204}{5} t_{2,2} t_{3,2} + \frac{1164}{25} t_{2,2} t_{3,3} t_{3,4} + \frac{1084}{125} t_{2,2} t_{3,4}^3 \right. \\
& + 54 t_{3,2} x_2 + \frac{468}{5} t_{3,3} t_{3,4} x_2 + \frac{468}{25} t_{3,4}^3 x_2 + \frac{78}{5} t_{2,2} t_{3,3} x_3 + \frac{1113}{25} t_{2,2} t_{3,4}^2 x_3 + 72 t_{3,3} x_2 x_3 - \frac{252}{5} t_{3,4}^2 x_2 x_3 + \frac{42}{5} t_{2,2} t_{3,4} x_3^2 \\
& - 108 t_{3,4} x_2 x_3^2 - 186 t_{2,2} x_3^3 \left. \right) + e^{7t_{\mu A}} \left(\frac{1391}{5} t_{1,1} t_{3,3} + \frac{12587}{50} t_{1,1} t_{3,4}^2 + 322 t_{3,3} x_1 + \frac{1969}{5} t_{3,4}^2 x_1 + \frac{2774}{5} t_{1,1} t_{3,4} x_3 \right. \\
& + 884 t_{3,4} x_1 x_3 + 213 t_{1,1} x_3^2 + 750 x_1 x_3^2 \left. \right) + e^{8t_{\mu A}} \left(596 t_{2,1} t_{3,4} + \frac{482}{15} t_{2,2}^2 t_{3,4} + \frac{2592}{5} t_{2,2} t_{3,4} x_2 + 972 t_{3,4} x_2^2 + 390 t_{2,1} x_3 \right. \\
& + 59 t_{2,2}^2 x_3 - 600 t_{2,2} x_2 x_3 - 1440 x_2^2 x_3 \left. \right) + e^{9t_{\mu A}} (1416 t_{1,1} t_{2,2} + 2052 t_{2,2} x_1 + 2520 t_{1,1} x_2 + 7820 x_1 x_2) \\
& + e^{10t_{\mu A}} \left(-1440 t_{3,1} - 1648 t_{3,3}^2 - 1988 t_{3,2} t_{3,4} - \frac{16518}{5} t_{3,3} t_{3,4}^2 - \frac{47047 t_{3,4}^4}{75} - 6630 t_{3,2} x_3 - 9696 t_{3,3} t_{3,4} x_3 \right. \\
& - 3024 t_{3,4}^3 x_3 - 16950 t_{3,3} x_3^2 - 9465 t_{3,4}^2 x_3^2 - 24190 t_{3,4} x_3^3 - 31200 x_3^4 \left. \right) + e^{12t_{\mu A}} \left(-14532 t_{2,2} t_{3,3} - \frac{72024}{5} t_{2,2} t_{3,4}^2 \right. \\
& - 37224 t_{3,3} x_2 - \frac{178704}{5} t_{3,4}^2 x_2 - 20108 t_{2,2} t_{3,4} x_3 + \frac{95112}{5} t_{3,4} x_2 x_3 + 12050 t_{2,2} x_3^2 + 9504 x_2 x_3^2 \left. \right) \\
& + e^{13t_{\mu A}} \left(-\frac{606656}{5} t_{1,1} t_{3,4} - \frac{1100028}{5} t_{3,4} x_1 - 82696 t_{1,1} x_3 - 155024 x_1 x_3 \right) + e^{14t_{\mu A}} \left(-73424 t_{2,1} - \frac{74872 t_{2,2}^2}{3} \right. \\
& - 166050 t_{2,2} x_2 - 213840 x_2^2 \left. \right) + e^{16t_{\mu A}} (378972 t_{3,2} + \frac{7113124}{5} t_{3,3} t_{3,4} + \frac{16080922 t_{3,4}^3}{25} + 1685018 t_{3,3} x_3
\end{aligned}$$

$$\begin{aligned}
& + 1682971t_{3,4}^2x_3 + 3127632t_{3,4}x_3^2 + 3654120x_3^3 \Big) + e^{18t_{\mu A}} \left(\frac{24792324}{5}t_{2,2}t_{3,4} + 10311552t_{3,4}x_2 + 2003564t_{2,2}x_3 \right. \\
& - 554400x_2x_3 \Big) + e^{19t_{\mu A}} (10436151t_{1,1} + 19383450x_1) + e^{22t_{\mu A}} (-119109720t_{3,3} - 202718352t_{3,4}^2 - 297398170t_{3,4}x_3 \\
& - 269136900x_3^2) + e^{24t_{\mu A}} (-363099900t_{2,2} - 646842240x_2) + e^{28t_{\mu A}} (23176371060t_{3,4} + 15870520800x_3) \\
& - 860105264700e^{34t_{\mu A}}, \\
\phi_{89}^3 = & e^{2t_{\mu A}} \left(-\frac{4}{5}t_{2,2}t_{3,2} - \frac{4}{25}t_{2,2}t_{3,3}t_{3,4} - \frac{2}{375}t_{2,2}t_{3,4}^3 - \frac{6}{5}t_{3,2}x_2 - \frac{6}{25}t_{3,3}t_{3,4}x_2 - \frac{1}{125}t_{3,4}^3x_2 - \frac{4}{5}t_{2,2}t_{3,3}x_3 \right. \\
& - \frac{4}{25}t_{2,2}t_{3,4}^2x_3 - \frac{6}{5}t_{2,2}t_{3,4}x_3^2 - 2t_{2,2}x_3^3 \Big) + e^{3t_{\mu A}} \left(-\frac{18}{5}t_{1,1}t_{3,3} - \frac{18}{25}t_{1,1}t_{3,4}^2 - \frac{12}{5}t_{3,3}x_1 - \frac{12}{25}t_{3,4}^2x_1 - \frac{18}{5}t_{1,1}t_{3,4}x_3 \right. \\
& - 6t_{1,1}x_3^2 + 24x_1x_3^2 \Big) + e^{4t_{\mu A}} \left(-\frac{48}{5}t_{2,1}t_{3,4} + \frac{4}{5}t_{2,2}^2t_{3,4} - \frac{6}{5}t_{2,2}t_{3,4}x_2 - \frac{36}{5}t_{3,4}x_2^2 - 36t_{2,1}x_3 - 48t_{2,2}x_2x_3 \right. \\
& - 72x_2^2x_3 \Big) + e^{5t_{\mu A}} (-t_{1,1}t_{2,2} + 6t_{2,2}x_1 + 3t_{1,1}x_2 - 18x_1x_2) + e^{6t_{\mu A}} \left(-18t_{3,1} + \frac{63t_{2,3}^2}{5} + 18t_{3,2}t_{3,4} + \frac{351}{25}t_{3,3}t_{3,4}^2 \right. \\
& + \frac{129t_{3,4}^4}{100} + 36t_{3,2}x_3 + \frac{324}{5}t_{3,3}t_{3,4}x_3 + 12t_{3,4}^3x_3 - 84t_{3,3}x_3^2 + 12t_{3,4}^2x_3^2 - 138t_{3,4}x_3^3 - 300x_3^4 \Big) + e^{8t_{\mu A}} \left(\frac{1522}{5}t_{2,2}t_{3,3} \right. \\
& + \frac{1447}{25}t_{2,2}t_{3,4}^2 + 534t_{3,3}x_2 + 231t_{3,4}^2x_2 + \frac{5152}{5}t_{2,2}t_{3,4}x_3 + 1296t_{3,4}x_2x_3 + 1614t_{2,2}x_3^2 - 810x_2x_3^2 \Big) \\
& + e^{9t_{\mu A}} \left(\frac{7857}{5}t_{1,1}t_{3,4} + 2130t_{3,4}x_1 + 4824t_{1,1}x_3 + 180x_1x_3 \right) + e^{10t_{\mu A}} \left(6470t_{2,1} - \frac{23t_{2,2}^2}{3} + 8548t_{2,2}x_2 + 14820x_2^2 \right) \\
& + e^{12t_{\mu A}} \left(-\frac{68424t_{3,2}}{5} - \frac{568974}{25}t_{3,3}t_{3,4} - \frac{660829t_{3,4}^3}{125} - \frac{119124}{5}t_{3,3}x_3 - \frac{679824}{25}t_{3,4}^2x_3 - \frac{36186}{5}t_{3,4}x_3^2 + 46488x_3^3 \right) \\
& + e^{14t_{\mu A}} \left(-\frac{588876}{5}t_{2,2}t_{3,4} - \frac{1659996}{5}t_{3,4}x_2 - 249308t_{2,2}x_3 - 65448x_2x_3 \right) + e^{15t_{\mu A}} (-433896t_{1,1} - 486384x_1) \\
& + e^{18t_{\mu A}} \left(3490344t_{3,3} + \frac{19796394t_{3,4}^2}{5} + 6493320t_{3,4}x_3 - 1275720x_3^2 \right) + e^{20t_{\mu A}} (16938594t_{2,2} + 28878030x_2) \\
& + e^{24t_{\mu A}} \left(-\frac{3397991478t_{3,4}}{5} - 292688964x_3 \right) + 29981411520e^{30t_{\mu A}}. \\
\phi_{99}^1 = & -e^{-t_{\mu A}}x_1^2x_2x_3 + e^{2t_{\mu A}} \left(-4t_{1,1}t_{2,2}t_{3,2} - \frac{8}{5}t_{1,1}t_{2,2}t_{3,3}t_{3,4} - \frac{8}{75}t_{1,1}t_{2,2}t_{3,4}^3 - 6t_{1,1}t_{3,2}x_2 - \frac{12}{5}t_{1,1}t_{3,3}t_{3,4}x_2 \right. \\
& - \frac{4}{25}t_{1,1}t_{3,4}^3x_2 - 12t_{3,2}x_1x_2 - \frac{24}{5}t_{3,3}t_{3,4}x_1x_2 - \frac{8}{25}t_{3,4}^3x_1x_2 - 6t_{1,1}t_{2,2}t_{3,3}x_3 - \frac{9}{5}t_{1,1}t_{2,2}t_{3,4}^2x_3 - 12t_{2,2}t_{3,3}x_1x_3 \\
& - \frac{18}{5}t_{2,2}t_{3,4}^2x_1x_3 - 9t_{1,1}t_{3,3}x_2x_3 - \frac{27}{10}t_{1,1}t_{3,4}^2x_2x_3 - 54t_{3,3}x_1x_2x_3 - \frac{81}{5}t_{3,4}^2x_1x_2x_3 - 8t_{1,1}t_{2,2}t_{3,4}x_3^2 - 16t_{2,2}t_{3,4}x_1x_3^2 \\
& - 12t_{1,1}t_{3,4}x_2x_3^2 - 72t_{3,4}x_1x_2x_3^2 - 10t_{1,1}t_{2,2}x_3^3 - 20t_{2,2}x_1x_3^3 - 15t_{1,1}x_2x_3^3 - 90x_1x_2x_3^3 \Big) + e^{3t_{\mu A}} \left(-9t_{1,1}^2t_{3,3} \right. \\
& - \frac{27}{10}t_{1,1}^2t_{3,4}^2 + 24t_{2,2}^2t_{3,3}x_2 + \frac{36}{5}t_{2,2}^2t_{3,4}^2x_2 + 72t_{2,2}t_{3,3}x_2^2 + \frac{108}{5}t_{2,2}t_{3,4}^2x_2^2 + 54t_{3,3}x_2^3 + \frac{81}{5}t_{3,4}^2x_2^3 + 24t_{3,2}^2x_3 \\
& - 12t_{1,1}^2t_{3,4}x_3 + \frac{96}{5}t_{3,2}t_{3,3}t_{3,4}x_3 + \frac{96}{25}t_{3,3}^2t_{3,4}^2x_3 + \frac{32}{25}t_{3,2}t_{3,4}^3x_3 + \frac{64}{125}t_{3,3}t_{3,4}^4x_3 + \frac{32t_{3,4}^6x_3}{1875} - 24t_{1,1}t_{3,4}x_1x_3 \\
& + 32t_{2,2}^2t_{3,4}x_2x_3 + 96t_{2,2}t_{3,4}x_2^2x_3 + 72t_{3,4}x_3^2x_3 - 15t_{1,1}^2x_3^2 + 72t_{3,2}t_{3,3}x_3^2 + \frac{144}{5}t_{3,3}^2t_{3,4}x_3^2 + \frac{108}{5}t_{3,2}t_{3,4}^2x_3^2 \\
& + \frac{264}{25}t_{3,3}t_{3,4}^3x_3^2 + \frac{72}{125}t_{3,4}^5x_3^2 - 30t_{1,1}x_1x_3^2 + 40t_{2,2}^2x_2x_3^2 + 120t_{2,2}x_2^2x_3^2 + 90x_2^3x_3^2 + 54t_{3,3}^2x_3^3 + 96t_{3,2}t_{3,4}x_3^3
\end{aligned}$$

$$\begin{aligned}
& + \frac{354}{5} t_{3,3} t_{3,4}^2 x_3^3 + \frac{371}{50} t_{3,4}^4 x_3^3 + 120 t_{3,2} x_3^4 + 192 t_{3,3} t_{3,4} x_3^4 + \frac{232}{5} t_{3,4}^3 x_3^4 + 180 t_{3,3} x_3^5 + 150 t_{3,4}^2 x_3^5 + 240 t_{3,4} x_3^6 + 150 x_3^7 \Big) \\
& + e^{4t_{\mu A}} \left(-16 t_{1,1} t_{2,2}^2 t_{3,4} + 24 t_{1,1} t_{2,2} t_{3,4} x_2 + 24 t_{2,2} t_{3,4} x_1 x_2 + 72 t_{1,1} t_{3,4} x_2^2 - 72 t_{3,4} x_1 x_2^2 - 10 t_{1,1} t_{2,2}^2 x_3 - 20 t_{2,2}^2 x_1 x_3 \right. \\
& + 45 t_{1,1} t_{2,2} x_2 x_3 - 90 t_{2,2} x_1 x_2 x_3 + 90 t_{1,1} x_2^2 x_3 - 360 x_1 x_2^2 x_3 \Big) + e^{5t_{\mu A}} \left(-35 t_{1,1}^2 t_{2,2} + 8 t_{2,2} t_{3,2} t_{3,3} + \frac{16}{5} t_{2,2} t_{3,3}^2 t_{3,4} \right. \\
& + \frac{12}{5} t_{2,2} t_{3,2} t_{3,4}^2 + \frac{88}{75} t_{2,2} t_{3,3} t_{3,4}^3 + \frac{8}{125} t_{2,2} t_{3,4}^5 - 30 t_{1,1}^2 x_2 + 40 t_{2,2}^3 x_2 + 120 t_{3,2} t_{3,3} x_2 + 48 t_{3,3}^2 t_{3,4} x_2 + 36 t_{3,2} t_{3,4}^2 x_2 \\
& + \frac{88}{5} t_{3,3} t_{3,4}^3 x_2 + \frac{24}{25} t_{3,4}^5 x_2 + 120 t_{1,1} x_1 x_2 + 120 t_{2,2}^2 x_2^2 + 90 t_{2,2} x_2^3 + 120 t_{2,2} t_{3,3}^2 x_3 - 96 t_{2,2} t_{3,2} t_{3,4} x_3 + \frac{168}{5} t_{2,2} t_{3,3} t_{3,4}^2 x_3 \\
& + \frac{206}{25} t_{2,2} t_{3,4}^4 x_3 + 180 t_{3,3}^2 x_2 x_3 + 288 t_{3,2} t_{3,4} x_2 x_3 + \frac{1116}{5} t_{3,3} t_{3,4}^2 x_2 x_3 + \frac{597}{25} t_{3,4}^4 x_2 x_3 - 240 t_{2,2} t_{3,2} x_2^3 + 64 t_{2,2} t_{3,3} t_{3,4} x_2^3 \\
& + \frac{208}{5} t_{2,2} t_{3,4}^3 x_2^3 + 360 t_{3,2} x_2 x_2^3 + 816 t_{3,3} t_{3,4} x_2 x_2^3 + \frac{1056}{5} t_{3,4}^3 x_2 x_2^3 + 20 t_{2,2} t_{3,3}^3 x_3^3 + 6 t_{2,2} t_{3,4}^2 x_3^3 + 840 t_{3,3} x_2 x_3^3 \\
& + 828 t_{3,4}^2 x_2 x_3^3 - 240 t_{2,2} t_{3,4} x_3^4 + 1440 t_{3,4} x_2 x_3^4 - 300 t_{2,2} x_3^5 + 900 x_2 x_3^5 \Big) + e^{6t_{\mu A}} \left(252 t_{1,1} t_{3,3}^2 - 36 t_{1,1} t_{3,2} t_{3,4} \right. \\
& + \frac{684}{5} t_{1,1} t_{3,3} t_{3,4}^2 + \frac{543}{25} t_{1,1} t_{3,4}^4 + 288 t_{3,3}^2 x_1 + \frac{864}{5} t_{3,3} t_{3,4}^2 x_1 + \frac{648}{25} t_{3,4}^4 x_1 - 480 t_{1,1} t_{3,2} x_3 + 522 t_{1,1} t_{3,3} t_{3,4} x_3 \\
& + \frac{1007}{5} t_{1,1} t_{3,4}^3 x_3 - 120 t_{3,2} x_1 x_3 + 1356 t_{3,3} t_{3,4} x_1 x_3 + 418 t_{3,4}^3 x_1 x_3 + 240 t_{1,1} t_{3,3} x_3^2 + 576 t_{1,1} t_{3,4}^2 x_3^2 + 1440 t_{3,3} x_1 x_3^2 \\
& + 1536 t_{3,4}^2 x_1 x_3^2 + 390 t_{1,1} t_{3,4} x_3^3 + 2340 t_{3,4} x_1 x_3^3 - 300 t_{1,1} x_3^4 + 1200 x_1 x_3^4 \Big) + e^{7t_{\mu A}} \left(576 t_{2,2}^2 t_{3,2} - \frac{368}{5} t_{2,2}^2 t_{3,3} t_{3,4} \right. \\
& - \frac{1896}{25} t_{2,2}^2 t_{3,4}^3 + 1944 t_{2,2} t_{3,2} x_2 - \frac{7872}{5} t_{2,2} t_{3,3} t_{3,4} x_2 - \frac{16344}{25} t_{2,2} t_{3,4}^2 x_2 + 2160 t_{3,2} x_2^2 - 1656 t_{3,3} t_{3,4} x_2^2 - \frac{3492}{5} t_{3,4}^2 x_2^2 \\
& + 464 t_{2,2}^2 t_{3,3} x_3 - \frac{264}{5} t_{2,2}^2 t_{3,4}^2 x_3 + 216 t_{2,2} t_{3,3} x_2 x_3 - \frac{14076}{5} t_{2,2} t_{3,4}^2 x_2 x_3 - 720 t_{3,3} x_2^2 x_3 - 2808 t_{3,4}^2 x_2^2 x_3 + 912 t_{2,2}^2 t_{3,4} x_2^3 \\
& - 3312 t_{2,2} t_{3,4} x_2 x_2^3 - 3240 t_{3,4} x_2^2 x_2^3 + 1590 t_{2,2}^2 x_3^3 + 360 t_{2,2} x_2 x_3^3 \Big) + e^{8t_{\mu A}} \left(2632 t_{1,1} t_{2,2} t_{3,3} + \frac{5068}{5} t_{1,1} t_{2,2} t_{3,4}^2 \right. \\
& + 2880 t_{2,2} t_{3,3} x_1 + 864 t_{2,2} t_{3,4}^2 x_1 + 4560 t_{1,1} t_{3,3} x_2 - 240 t_{1,1} t_{3,4}^2 x_2 + 9360 t_{3,3} x_1 x_2 + 3624 t_{3,4}^2 x_1 x_2 + 4956 t_{1,1} t_{2,2} t_{3,4} x_3 \\
& + 8496 t_{2,2} t_{3,4} x_1 x_3 + 4005 t_{1,1} t_{3,4} x_2 x_3 + 27270 t_{3,4} x_1 x_2 x_3 + 7520 t_{1,1} t_{2,2} x_3^2 + 10720 t_{2,2} x_1 x_3^2 + 8400 t_{1,1} x_2 x_3^2 \\
& + 32400 x_1 x_2 x_3^2 \Big) + e^{9t_{\mu A}} \left(-720 t_{3,2}^2 - 1440 t_{3,3}^3 + 1845 t_{1,1}^2 t_{3,4} - 1536 t_{2,2}^3 t_{3,4} - 360 t_{3,2} t_{3,3} t_{3,4} - \frac{6624}{5} t_{3,3}^2 t_{3,4}^2 \right. \\
& + \frac{132}{5} t_{3,2} t_{3,4}^3 - \frac{9312}{25} t_{3,3} t_{3,4}^4 - \frac{4708 t_{3,4}^6}{125} - 13176 t_{2,2}^2 t_{3,4} x_2 - 25488 t_{2,2} t_{3,4} x_2^2 - 12150 t_{3,4} x_2^3 + 3600 t_{1,1}^2 x_3 - 2760 t_{2,2}^3 x_3 \\
& - 6480 t_{3,2} t_{3,3} x_3 - 9048 t_{3,3}^2 t_{3,4} x_3 - 3768 t_{3,2} t_{3,4}^2 x_3 - \frac{27768}{5} t_{3,3} t_{3,4}^3 x_3 - \frac{17038}{25} t_{3,4}^5 x_3 + 3600 t_{1,1} x_1 x_3 - 23180 t_{2,2}^2 x_2 x_3 \\
& - 54960 t_{2,2} x_2^2 x_3 - 39600 x_2^3 x_3 - 15360 t_{3,3}^2 x_3^2 - 15840 t_{3,2} t_{3,4} x_3^2 - 27360 t_{3,3} t_{3,4} x_3^2 - \frac{26736}{5} t_{3,4}^4 x_3^2 - 24000 t_{3,2} x_3^3 \\
& - 61980 t_{3,3} t_{3,4} x_3^3 - 24034 t_{3,4}^3 x_3^3 - 57600 t_{3,3} x_3^4 - 63600 t_{3,4}^2 x_3^4 - 93900 t_{3,4} x_3^5 - 60000 x_3^6 \Big) + e^{10t_{\mu A}} \left(2680 t_{1,1} t_{2,2}^2 \right. \\
& + 7200 t_{2,2}^2 x_1 + 4890 t_{1,1} t_{2,2} x_2 + 37380 t_{2,2} x_1 x_2 + 3600 t_{1,1} x_2^2 + 67680 x_1 x_2^2 \Big) + e^{11t_{\mu A}} \left(-18048 t_{2,2} t_{3,3}^2 \right. \\
& - 24016 t_{2,2} t_{3,2} t_{3,4} - \frac{33136}{5} t_{2,2} t_{3,3} t_{3,4}^2 + \frac{140824}{75} t_{2,2} t_{3,4}^4 - 44640 t_{3,3}^2 x_2 - 76200 t_{3,2} t_{3,4} x_2 - 21864 t_{3,3} t_{3,4}^2 x_2 \\
& + \frac{22852}{5} t_{3,4}^4 x_2 + 13920 t_{2,2} t_{3,2} x_3 - 76608 t_{2,2} t_{3,3} t_{3,4} x_3 - \frac{34048}{5} t_{2,2} t_{3,4}^3 x_3 - 72000 t_{3,2} x_2 x_3 - 176040 t_{3,3} t_{3,4} x_2 x_3 \\
& - 12972 t_{3,4}^3 x_2 x_3 - 55960 t_{2,2} t_{3,3} x_2^2 - 60164 t_{2,2} t_{3,4}^2 x_2^2 - 261600 t_{3,3} x_2 x_2^2 - 187320 t_{3,4}^2 x_2 x_2^2 - 77860 t_{2,2} t_{3,4} x_2^3 \\
& - 475800 t_{3,4} x_2 x_2^3 + 22200 t_{2,2} x_3^4 - 360000 x_2 x_3^4 \Big) + e^{12t_{\mu A}} \left(10656 t_{1,1} t_{3,2} - \frac{689688}{5} t_{1,1} t_{3,3} t_{3,4} - \frac{803796}{25} t_{1,1} t_{3,4}^3 \right. \\
& - 216000 t_{3,3} t_{3,4} x_1 - 64800 t_{3,4}^3 x_1 - 60816 t_{1,1} t_{3,3} x_3 - \frac{1042074}{5} t_{1,1} t_{3,4}^2 x_3 - 230472 t_{3,3} x_1 x_3 - \frac{2470608}{5} t_{3,4}^2 x_1 x_3
\end{aligned}$$

$$\begin{aligned}
& -321888t_{1,1}t_{3,4}x_3^2 - 910896t_{3,4}x_1x_3^2 - 21360t_{1,1}x_3^3 - 438120x_1x_3^3 \Big) + e^{13t_{\mu A}} \Big(-68024t_{2,2}^2t_{3,3} + \frac{556684}{5}t_{2,2}^2t_{3,4}^2 \\
& - 150792t_{2,2}t_{3,3}x_2 + \frac{3054972}{5}t_{2,2}t_{3,4}^2x_2 - 160128t_{3,3}x_2^2 + \frac{2350008}{5}t_{3,4}^2x_2^2 + 137048t_{2,2}^2t_{3,4}x_3 + 1611144t_{2,2}t_{3,4}x_2x_3 \\
& + 1743696t_{3,4}x_2^2x_3 - 204240t_{2,2}^2x_3^2 + 321480t_{2,2}x_2x_3^2 + 285120x_2^2x_3^2 \Big) + e^{14t_{\mu A}} (-493452t_{1,1}t_{2,2}t_{3,4} - 1080000t_{2,2}t_{3,4}x_1 \\
& - 646416t_{1,1}t_{3,4}x_2 - 3380736t_{3,4}x_1x_2 - 749670t_{1,1}t_{2,2}x_3 - 1260540t_{2,2}x_1x_3 - 665505t_{1,1}x_2x_3 - 3973590x_1x_2x_3) \\
& + e^{15t_{\mu A}} \Big(-193185t_{1,1}^2 + 111840t_{2,2}^3 + 159408t_{3,2}t_{3,3} + \frac{5920416}{5}t_{3,3}^2t_{3,4} + \frac{2103912}{5}t_{3,2}t_{3,4}^2 + \frac{17101296}{25}t_{3,3}t_{3,4}^3 \\
& + \frac{7979808t_{3,4}^5}{125} + 861840t_{2,2}^2x_2 + 1547640t_{2,2}x_2^2 + 614790x_2^3 + 1490160t_{3,3}^2x_3 + 1577424t_{3,2}t_{3,4}x_3 + \frac{23950128}{5}t_{3,3}t_{3,4}^2x_3 \\
& + \frac{25157276}{25}t_{3,4}^4x_3 + 2099160t_{3,2}x_3^2 + 10896144t_{3,3}t_{3,4}x_3^2 + \frac{27179808}{5}t_{3,4}^3x_3^2 + 8679420t_{3,3}^3x_3^3 + 14474376t_{3,4}^2x_3^3 \\
& + 19799760t_{3,4}^4x_3^4 + 11699700x_3^5 \Big) + e^{17t_{\mu A}} \Big(2543256t_{2,2}t_{3,2} + \frac{42484072}{5}t_{2,2}t_{3,3}t_{3,4} - \frac{35982556}{25}t_{2,2}t_{3,4}^3 + 8075160t_{3,2}x_2 \\
& + 19675632t_{3,3}t_{3,4}x_2 - 1700232t_{3,4}^3x_2 + 10344384t_{2,2}t_{3,3}x_3 + \frac{12508056}{5}t_{2,2}t_{3,4}^2x_3 + 24713640t_{3,3}x_2x_3 \\
& + 1013472t_{3,4}^2x_2x_3 + 18909632t_{2,2}t_{3,4}x_3^2 + 54765120t_{3,4}x_2x_3^2 + 9743940t_{2,2}x_3^3 + 56039400x_2x_3^3 \Big) \\
& + e^{18t_{\mu A}} (12174840t_{1,1}t_{3,3} + 18742176t_{1,1}t_{3,4}^2 + 21384000t_{3,3}x_1 + 46915200t_{3,4}^2x_1 + 43516650t_{1,1}t_{3,4}x_3 \\
& + 117255780t_{3,4}x_1x_3 + 25045200t_{1,1}x_3^2 + 86731200x_1x_3^2) + e^{19t_{\mu A}} (-9348032t_{2,2}^2t_{3,4} - 60201816t_{2,2}t_{3,4}x_2 \\
& - 34585560t_{3,4}x_2^2 - 7344120t_{2,2}^2x_3 - 118365480t_{2,2}x_2x_3 - 130798800x_2^2x_3) + e^{20t_{\mu A}} (44237480t_{1,1}t_{2,2} + 106920000t_{2,2}x_1 \\
& + 73369200t_{1,1}x_2 + 322908000x_1x_2) + e^{21t_{\mu A}} \Big(-111676320t_{3,3}^2 - 105057720t_{3,2}t_{3,4} - 372047472t_{3,3}t_{3,4}^2 \\
& - \frac{312471528t_{3,4}^4}{5} - 135330000t_{3,2}x_3 - 1028764080t_{3,3}t_{3,4}x_3 - 520356832t_{3,4}^3x_3 - 938318400t_{3,3}x_3^2 - 1821978240t_{3,4}^2x_3^2 \\
& - 2728752300t_{3,4}x_3^3 - 1595040000x_3^4 \Big) + e^{23t_{\mu A}} \Big(-818066800t_{2,2}t_{3,3} - 251287096t_{2,2}t_{3,4}^2 - 1837633056t_{3,3}x_2 \\
& - \frac{4231559784}{5}t_{3,4}^2x_2 - 790690320t_{2,2}t_{3,4}x_3 - 989596008t_{3,4}x_2x_3 - 1433740200t_{2,2}x_3^2 - 4342613760x_2x_3^2 \Big) \\
& + e^{24t_{\mu A}} (-3018152376t_{1,1}t_{3,4} - 8019000000t_{3,4}x_1 - 2451801360t_{1,1}x_3 - 7485858720x_1x_3) + e^{25t_{\mu A}} (-147207960t_{2,2}^2 \\
& + 698942520t_{2,2}x_2 - 208802880x_2^2) + e^{27t_{\mu A}} (6205115520t_{3,2} + 57329810784t_{3,3}t_{3,4} + 29106573360t_{3,4}^3 \\
& + 61873919280t_{3,3}x_3 + 118547035032t_{3,4}^2x_3 + 236876184000t_{3,4}x_3^2 + 156672016950x_3^3) + e^{29t_{\mu A}} (86399501864t_{2,2}t_{3,4} \\
& + 193415433600t_{3,4}x_2 + 60645688440t_{2,2}x_3 + 100959556500x_2x_3) + e^{30t_{\mu A}} (144023219100t_{1,1} + 396940500000x_1) \\
& + e^{33t_{\mu A}} (-2655547480800t_{3,3} - 5537714448240t_{3,4}^2 - 11844161039400t_{3,4}x_3 - 10402175520000x_3^2) \\
& + e^{35t_{\mu A}} (-4835694794400t_{2,2} - 9566526202560x_2) + e^{39t_{\mu A}} (452466384691680t_{3,4} + 424218905102400x_3) \\
& - 13317074584740000e^{45t_{\mu A}},
\end{aligned}$$

$$\begin{aligned}
\phi_{99}^2 = & -t_{1,1}x_1x_2 - 2x_1^2x_2 + e^{2t_{\mu A}} \Big(-4t_{2,1}t_{3,2} - \frac{2}{3}t_{2,2}^2t_{3,2} - \frac{8}{5}t_{2,1}t_{3,3}t_{3,4} - \frac{4}{15}t_{2,2}^2t_{3,3}t_{3,4} - \frac{8}{75}t_{2,1}t_{3,4}^3 - \frac{4}{225}t_{2,2}^2t_{3,4}^3 \\
& - 6t_{2,1}t_{3,3}x_3 - t_{2,2}^2t_{3,3}x_3 - \frac{9}{5}t_{2,1}t_{3,4}^2x_3 - \frac{3}{10}t_{2,2}^2t_{3,4}^2x_3 + 12t_{2,2}t_{3,3}x_2x_3 + \frac{18}{5}t_{2,2}t_{3,4}^2x_2x_3 + 18t_{3,3}x_2^2x_3 + \frac{27}{5}t_{3,4}^2x_2^2x_3 \\
& - 8t_{2,1}t_{3,4}x_3^2 - \frac{4}{3}t_{2,2}^2t_{3,4}x_3^2 + 16t_{2,2}t_{3,4}x_2x_3^2 + 24t_{3,4}x_2^2x_3^2 - 10t_{2,1}x_3^3 - \frac{5}{3}t_{2,2}^2x_3^3 + 20t_{2,2}x_2x_3^3 + 30x_2^2x_3^3 \Big) \\
& + e^{3t_{\mu A}} \Big(-24t_{1,1}t_{2,2}t_{3,3} - \frac{36}{5}t_{1,1}t_{2,2}t_{3,4}^2 - 24t_{2,2}t_{3,3}x_1 - \frac{36}{5}t_{2,2}t_{3,4}^2x_1 - 18t_{1,1}t_{3,3}x_2 - \frac{27}{5}t_{1,1}t_{3,4}^2x_2 - 52t_{3,3}x_1x_2
\end{aligned}$$

$$\begin{aligned}
& -\frac{78}{5}t_{3,4}^2x_1x_2 - 32t_{1,1}t_{2,2}t_{3,4}x_3 - 32t_{2,2}t_{3,4}x_1x_3 - 144t_{3,4}x_1x_2x_3 - 40t_{1,1}t_{2,2}x_3^2 - 40t_{2,2}x_1x_3^2 - 180x_1x_2x_3^2 \\
& + e^{4t_{\mu A}} \left(12t_{3,2}^2 - 48t_{1,1}^2t_{3,4} - 16t_{2,1}t_{2,2}t_{3,4} - \frac{8}{3}t_{3,2}^2t_{3,4} + \frac{48}{5}t_{3,2}t_{3,3}t_{3,4} + \frac{48}{25}t_{3,3}^2t_{3,4} + \frac{16}{25}t_{3,2}t_{3,4}^2 + \frac{32}{125}t_{3,3}t_{3,4}^2 \right. \\
& + \frac{16t_{3,4}^6}{1875} - 48t_{1,1}t_{3,4}x_1 - 60t_{1,1}^2x_3 - 10t_{2,1}t_{2,2}x_3 - \frac{5}{3}t_{2,2}^3x_3 + 90t_{3,2}t_{3,3}x_3 + 36t_{3,3}^2t_{3,4}x_3 + 27t_{3,2}t_{3,4}^2x_3 + \frac{66}{5}t_{3,3}t_{3,4}^3x_3 \\
& + \frac{18}{25}t_{3,4}^5x_3 - 60t_{1,1}x_1x_3 + 20t_{2,2}^2x_2x_3 + 30t_{2,2}x_2^2x_3 + 108t_{3,3}^2x_3^2 + 120t_{3,2}t_{3,4}x_3^2 + \frac{564}{5}t_{3,3}t_{3,4}^2x_3^2 + \frac{323}{25}t_{3,4}^4x_3^2 \\
& + 150t_{3,2}x_3^3 + 348t_{3,3}t_{3,4}x_3^3 + \frac{452}{5}t_{3,4}^3x_3^3 + 360t_{3,3}^4x_3^4 + 300t_{3,4}^2x_3^4 + 480t_{3,4}x_3^5 + 300x_3^6 \left. \right) + e^{5t_{\mu A}} \left(-20t_{1,1}t_{2,1} \right. \\
& - \frac{130}{3}t_{1,1}t_{2,2}^2 - 40t_{2,2}^2x_1 - 30t_{1,1}t_{2,2}x_2 + 84t_{2,2}x_1x_2 \left. \right) + e^{6t_{\mu A}} \left(48t_{2,2}t_{3,3}^2 + 204t_{2,2}t_{3,2}t_{3,4} + \frac{552}{5}t_{2,2}t_{3,3}t_{3,4}^2 \right. \\
& + \frac{244}{25}t_{2,2}t_{3,4}^4 + 144t_{3,3}^2x_2 + 432t_{3,2}t_{3,4}x_2 + \frac{1296}{5}t_{3,3}t_{3,4}^2x_2 + \frac{612}{25}t_{3,4}^4x_2 - 90t_{2,2}t_{3,2}x_3 + 262t_{2,2}t_{3,3}t_{3,4}x_3 \\
& + 87t_{2,2}t_{3,4}^3x_3 + 720t_{3,2}x_2x_3 - 36t_{3,3}t_{3,4}x_2x_3 - 78t_{3,4}^3x_2x_3 - 280t_{2,2}t_{3,3}x_3^2 + 228t_{2,2}t_{3,4}^2x_3^2 - 432t_{3,4}^2x_2x_3^2 \\
& - 90t_{2,2}t_{3,4}x_3^3 - 540t_{3,4}x_2x_3^3 - 600t_{2,2}x_3^4 \left. \right) + e^{7t_{\mu A}} \left(36t_{1,1}t_{3,2} + \frac{9432}{5}t_{1,1}t_{3,3}t_{3,4} + \frac{14064}{25}t_{1,1}t_{3,4}^2 - 240t_{3,2}x_1 \right. \\
& + 2712t_{3,3}t_{3,4}x_1 + 836t_{3,4}^3x_1 + 1584t_{1,1}t_{3,3}x_3 + \frac{12936}{5}t_{1,1}t_{3,4}^2x_3 + 2880t_{3,3}x_1x_3 + 3072t_{3,4}^2x_1x_3 + 4272t_{1,1}t_{3,4}x_3^2 \\
& + 4680t_{3,4}x_1x_3^2 + 2040t_{1,1}x_3^3 + 2400x_1x_3^3 \left. \right) + e^{8t_{\mu A}} \left(480t_{2,1}t_{3,3} + 976t_{2,2}^2t_{3,3} + 2240t_{2,1}t_{3,4}^2 + \frac{1712}{15}t_{2,2}^2t_{3,4}^2 \right. \\
& + 672t_{2,2}t_{3,3}x_2 + \frac{9648}{5}t_{2,2}t_{3,4}^2x_2 + 3888t_{3,4}^2x_2^2 + 5670t_{2,1}t_{3,4}x_3 + 663t_{2,2}^2t_{3,4}x_3 - 4680t_{2,2}t_{3,4}x_2x_3 - 4050t_{3,4}^2x_2^2x_3 \\
& + 4400t_{2,1}x_3^2 + \frac{5980}{3}t_{2,2}^2x_3^2 - 8800t_{2,2}x_2x_3^2 - 13200x_2^2x_3^2 \left. \right) + e^{9t_{\mu A}} (11028t_{1,1}t_{2,2}t_{3,4} + 16992t_{2,2}t_{3,4}x_1 + 12690t_{1,1}t_{3,4}x_2 \\
& + 35540t_{3,4}x_1x_2 + 15920t_{1,1}t_{2,2}x_3 + 21440t_{2,2}x_1x_3 + 14400t_{1,1}x_2x_3 + 64800x_1x_2x_3) + e^{10t_{\mu A}} \left(6480t_{1,1}^2 + 3580t_{2,1}t_{2,2} \right. \\
& + \frac{1070t_{2,2}^3}{3} - 2820t_{3,2}t_{3,3} - 12312t_{3,3}^2t_{3,4} - 9822t_{3,2}t_{3,4}^2 - \frac{53572}{5}t_{3,3}t_{3,4}^2 - \frac{31712t_{3,4}^5}{25} + 7200t_{1,1}x_1 + 720t_{2,2}^2x_2 \\
& - 21990t_{3,3}^2x_3 - 36810t_{3,2}t_{3,4}x_3 - 48846t_{3,3}t_{3,4}^2x_3 - \frac{92391}{10}t_{3,4}^4x_3 - 37200t_{3,2}x_3^2 - 112800t_{3,3}t_{3,4}x_3^2 - 41120t_{3,4}^3x_3^2 \\
& - 105450t_{3,3}^3x_3^3 - 118515t_{3,4}^2x_3^3 - 187800t_{3,4}^4x_3^4 - 120000x_3^5 \left. \right) + e^{12t_{\mu A}} \left(-37320t_{2,2}t_{3,2} - 134976t_{2,2}t_{3,3}t_{3,4} \right. \\
& - \frac{238568}{5}t_{2,2}t_{3,4}^3 - 95040t_{3,2}x_2 - 218016t_{3,3}t_{3,4}x_2 - \frac{623952}{5}t_{3,4}^3x_2 - 142000t_{2,2}t_{3,3}x_3 - 123218t_{2,2}t_{3,4}^2x_3 \\
& - 187488t_{3,3}x_2x_3 + \frac{97668}{5}t_{3,4}^2x_2x_3 - 118240t_{2,2}t_{3,4}x_3^2 + 275616t_{3,4}x_2x_3^2 + 12150t_{2,2}x_3^3 + 47520x_2x_3^3 \left. \right) \\
& + e^{13t_{\mu A}} \left(-310524t_{1,1}t_{3,3} - \frac{2823906}{5}t_{1,1}t_{3,4}^2 - 460944t_{3,3}x_1 - \frac{4941216}{5}t_{3,4}^2x_1 - 1299552t_{1,1}t_{3,4}x_3 - 1821792t_{3,4}x_1x_3 \right. \\
& - 826440t_{1,1}x_3^2 - 876240x_1x_3^2 \left. \right) + e^{14t_{\mu A}} \left(-767776t_{2,1}t_{3,4} - \frac{743612}{3}t_{2,2}^2t_{3,4} - 1179120t_{2,2}t_{3,4}x_2 - 1710720t_{3,4}^2x_2^2 \right. \\
& - 1018710t_{2,1}x_3 - 313395t_{2,2}^2x_3 + 14940t_{2,2}x_2x_3 + 204930x_2^2x_3 \left. \right) + e^{15t_{\mu A}} (-1454700t_{1,1}t_{2,2} - 2521080t_{2,2}x_1 \\
& - 2105730t_{1,1}x_2 - 6105780x_1x_2) + e^{16t_{\mu A}} \left(2165568t_{3,3}^2 + 3609984t_{3,2}t_{3,4} + \frac{40336272}{5}t_{3,3}t_{3,4}^2 + \frac{54600584t_{3,4}^4}{25} \right. \\
& + 5544810t_{3,2}x_3 + 19824522t_{3,3}t_{3,4}x_3 + 8788881t_{3,4}^3x_3 + 16810920t_{3,3}^2x_3^2 + 21775608t_{3,4}^2x_3^2 + 34558560t_{3,4}^3x_3^3 \\
& + 23399400x_3^4 \left. \right) + e^{18t_{\mu A}} \left(18291996t_{2,2}t_{3,3} + \frac{153341514}{5}t_{2,2}t_{3,4}^2 + 31220640t_{3,3}x_2 + 63970560t_{3,4}^2x_2 \right.
\end{aligned}$$

$$\begin{aligned}
& + 44706608t_{2,2}t_{3,4}x_3 + 13201020t_{3,4}x_2x_3 + 14586360t_{2,2}x_3^2 - 20908800x_2x_3^2 \Big) + e^{19t_{\mu A}} (130053624t_{1,1}t_{3,4} \\
& + 234511560t_{3,4}x_1 + 126689040t_{1,1}x_3 + 173462400x_1x_3) + e^{20t_{\mu A}} \left(70426400t_{2,1} + \frac{107803060t_{2,2}^2}{3} + 154487520t_{2,2}x_2 \right. \\
& \left. + 188179200x_2^2 \right) + e^{22t_{\mu A}} (-322723200t_{3,2} - 1658859840t_{3,3}t_{3,4} - 1104177104t_{3,4}^3 - 1751296950t_{3,3}x_3 \\
& - 2789043165t_{3,4}^2x_3 - 3852264600t_{3,4}x_3^2 - 2730360000x_3^3) + e^{24t_{\mu A}} (-5796453600t_{2,2}t_{3,4} - 10520500320t_{3,4}x_2 \\
& - 3593143350t_{2,2}x_3 - 1178290080x_2x_3) + e^{25t_{\mu A}} (-8480495040t_{1,1} - 14971717440x_1) + e^{28t_{\mu A}} (98859098880t_{3,3} \\
& + 229725744576t_{3,4}^2 + 338692970880t_{3,4}x_3 + 212350641900x_3^2) + e^{30t_{\mu A}} (321584010960t_{2,2} + 508631389200x_2) \\
& + e^{34t_{\mu A}} (-20567085718800t_{3,4} - 13652202240000x_3) + 651652721212800e^{40t_{\mu A}}, \\
\\
\phi_{99}^3 = & e^{t_{\mu A}} \left(-t_{1,1}t_{2,1} - \frac{1}{6}t_{1,1}t_{2,2}^2 - 2t_{2,1}x_1 - \frac{1}{3}t_{2,2}^2x_1 - 2t_{1,1}t_{2,2}x_2 - 4t_{2,2}x_1x_2 - 3t_{1,1}x_2^2 - 6x_1x_2^2 \right) + e^{2t_{\mu A}} \left(-4t_{2,2}t_{3,1} \right. \\
& - \frac{2}{5}t_{2,2}t_{3,3}^2 - \frac{4}{5}t_{2,2}t_{3,2}t_{3,4} - \frac{2}{25}t_{2,2}t_{3,3}t_{3,4}^2 - \frac{1}{750}t_{2,2}t_{3,4}^4 - 6t_{3,1}x_2 - \frac{3}{5}t_{3,3}^2x_2 - \frac{6}{5}t_{3,2}t_{3,4}x_2 - \frac{3}{25}t_{3,3}t_{3,4}^2x_2 - \frac{1}{500}t_{3,4}^4x_2 \\
& + 12t_{3,2}x_2x_3 + \frac{24}{5}t_{3,3}t_{3,4}x_2x_3 + \frac{8}{25}t_{3,4}^2x_2x_3 + 18t_{3,3}x_2x_3^2 + \frac{27}{5}t_{3,4}^2x_2x_3^2 + 24t_{3,4}x_2x_3^3 + 30x_2x_3^4 \Big) + e^{3t_{\mu A}} \left(-21t_{1,1}t_{3,2} \right. \\
& - \frac{42}{5}t_{1,1}t_{3,3}t_{3,4} - \frac{14}{25}t_{1,1}t_{3,4}^3 - 18t_{3,2}x_1 - \frac{36}{5}t_{3,3}t_{3,4}x_1 - \frac{12}{25}t_{3,4}^3x_1 - 18t_{1,1}t_{3,3}x_3 - \frac{27}{5}t_{1,1}t_{3,4}^2x_3 - 24t_{1,1}t_{3,4}x_3^2 \\
& - 30t_{1,1}x_3^3 \Big) + e^{4t_{\mu A}} \left(-52t_{2,1}t_{3,3} + \frac{10}{3}t_{2,2}^2t_{3,3} - \frac{78}{5}t_{2,1}t_{3,4}^2 + t_{2,2}^2t_{3,4}^2 - 14t_{2,2}t_{3,3}x_2 - \frac{21}{5}t_{2,2}t_{3,4}^2x_2 - 48t_{3,3}x_2^2 \right. \\
& - \frac{72}{5}t_{3,4}^2x_2^2 - 48t_{2,1}t_{3,4}x_3 + 8t_{2,2}^2t_{3,4}x_3 - 48t_{2,2}t_{3,4}x_2x_3 - 60t_{2,1}x_3^2 + 10t_{2,2}^2x_3^2 - 90t_{2,2}x_2x_3^2 \Big) + e^{5t_{\mu A}} (23t_{1,1}t_{2,2}t_{3,4} \\
& + 78t_{2,2}t_{3,4}x_1 + 30t_{1,1}t_{3,4}x_2 - 36t_{3,4}x_1x_2 + 90t_{1,1}t_{2,2}x_3 - 120t_{1,1}x_2x_3 - 720x_1x_2x_3) + e^{6t_{\mu A}} \left(84t_{1,1}^2 + 144t_{2,1}t_{2,2} \right. \\
& + 4t_{2,2}^3 + 204t_{3,2}t_{3,3} - 36t_{3,1}t_{3,4} + 78t_{3,3}^2t_{3,4} + 54t_{3,2}t_{3,4}^2 + \frac{146}{5}t_{3,3}t_{3,4}^3 + \frac{81t_{3,4}^5}{50} + 288t_{1,1}x_1 + 90t_{2,1}x_2 + 273t_{2,2}^2x_2 \\
& + 432t_{2,2}x_2^2 + 144t_{3,3}^2x_3 + 144t_{3,2}t_{3,4}x_3 + 144t_{3,3}t_{3,4}^2x_3 + \frac{84}{5}t_{3,4}^4x_3 + 180t_{3,2}x_3^2 + 264t_{3,3}t_{3,4}x_3^2 + \frac{312}{5}t_{3,4}^3x_3^2 + 240t_{3,3}x_3^3 \\
& + 72t_{3,4}^2x_3^3 \Big) + e^{8t_{\mu A}} \left(2324t_{2,2}t_{3,2} + \frac{3868}{5}t_{2,2}t_{3,3}t_{3,4} + \frac{1138}{75}t_{2,2}t_{3,4}^3 + 4080t_{3,2}x_2 + 2946t_{3,3}t_{3,4}x_2 + 503t_{3,4}^3x_2 \right. \\
& + 3552t_{2,2}t_{3,3}x_3 + \frac{5568}{5}t_{2,2}t_{3,4}^2x_3 - 1440t_{3,3}x_2x_3 - 96t_{3,4}^2x_2x_3 + 5436t_{2,2}t_{3,4}x_3^2 - 2970t_{3,4}x_2x_3^2 + 6720t_{2,2}x_3^3 - 6000x_2x_3^3 \Big) \\
& + e^{9t_{\mu A}} \left(12783t_{1,1}t_{3,3} + \frac{40749}{10}t_{1,1}t_{3,4}^2 + 18270t_{3,3}x_1 + 6633t_{3,4}^2x_1 + 19242t_{1,1}t_{3,4}x_3 + 25440t_{1,1}x_3^2 \right) \\
& + e^{10t_{\mu A}} \left(36080t_{2,1}t_{3,4} - \frac{8096}{3}t_{2,2}^2t_{3,4} + 39220t_{2,2}t_{3,4}x_2 + 82320t_{3,4}^2x_2 + 48000t_{2,1}x_3 - 8960t_{2,2}^2x_3 + 10380t_{2,2}x_2x_3 \right) \\
& + e^{11t_{\mu A}} (20699t_{1,1}t_{2,2} + 71310t_{2,2}x_1 + 83568t_{1,1}x_2 + 125856x_1x_2) + e^{12t_{\mu A}} \left(125856t_{3,1} - \frac{384372t_{3,3}^2}{5} - \frac{610044}{5}t_{3,2}t_{3,4} \right. \\
& - \frac{2983272}{25}t_{3,3}t_{3,4}^2 - \frac{1840531t_{3,4}^4}{125} + 72000t_{3,2}x_3 - 196560t_{3,3}t_{3,4}x_3 - 70152t_{3,4}^3x_3 + 100500t_{3,3}x_3^2 - 100230t_{3,4}^2x_3^2 \\
& \left. + 204000t_{3,4}x_3^3 + 450000x_3^4 \right) + e^{14t_{\mu A}} \left(-1057056t_{2,2}t_{3,3} - \frac{2936184}{5}t_{2,2}t_{3,4}^2 - 1362384t_{3,3}x_2 - \frac{9687726}{5}t_{3,4}^2x_2 \right)
\end{aligned}$$

$$\begin{aligned}
& -1909968t_{2,2}t_{3,4}x_3 + 77952t_{3,4}x_2x_3 - 2989260t_{2,2}x_3^2 + 427410x_2x_3^2) + e^{15t_{\mu A}}(-4862592t_{1,1}t_{3,4} - 6648192t_{3,4}x_1 \\
& - 6363810t_{1,1}x_3) + e^{16t_{\mu A}}(-7993620t_{2,1} - 599310t_{2,2}^2 - 11865582t_{2,2}x_2 - 17087760x_2^2) + e^{18t_{\mu A}}(13279500t_{3,2} \\
& + 58919244t_{3,3}t_{3,4} + \frac{118070466t_{3,4}^3}{5} + 16506720t_{3,3}x_3 + 46114704t_{3,4}^2x_3 + 7337760t_{3,4}x_3^2 - 79182000x_3^3) \\
& + e^{20t_{\mu A}}(289168248t_{2,2}t_{3,4} + 582458130t_{3,4}x_2 + 337148640t_{2,2}x_3 - 11652000x_2x_3) + e^{21t_{\mu A}}(598087680t_{1,1} \\
& + 640800000x_1) + e^{24t_{\mu A}}\left(-4658254068t_{3,3} - \frac{42954595602t_{3,4}^2}{5} - 6772067424t_{3,4}x_3 + 3614130720x_3^2\right) \\
& + e^{26t_{\mu A}}(-25117901460t_{2,2} - 36050857920x_2) + e^{30t_{\mu A}}(1001073115680t_{3,4} + 176276934000x_3) - 34893343710720e^{36t_{\mu A}}.
\end{aligned}$$

A.12. ψ for $A = (2, 3, 5)$.

$$\begin{aligned}
\psi^1 &= x_1 + e^{3t_{\mu A}}\left(18t_{3,3} + \frac{27t_{3,4}^2}{5} + 48t_{3,4}x_3 + 90x_3^2\right) + e^{5t_{\mu A}}(-210t_{2,2} - 360x_2) \\
&+ e^{9t_{\mu A}}(13230t_{3,4} - 12000x_3) - 2708910e^{15t_{\mu A}}, \\
\psi^2 &= -48e^{4t_{\mu A}}t_{3,4} + 40800e^{10t_{\mu A}}, \\
\psi^3 &= -1440e^{6t_{\mu A}}.
\end{aligned}$$

REFERENCES

- [1] D. Abramovich, T. Graber and A. Vistoli, *Gromov–Witten theory of Deligne–Mumford stacks*, Amer. J. Math. **130** (2008), no. 5, 1337–1398.
- [2] W. Chen and Y. Ruan, *Orbifold Gromov–Witten Theory*, Orbifolds in mathematics and physics (Madison, WI, 2001), 25–85, Contemp. Math., **310**, Amer. Math. Soc., Providence, RI, 2002.
- [3] A. Douai and C. Sabbah, *Gauss–Manin systems, Brieskorn lattices and Frobenius structures* (I), Proceedings of the International Conference in Honor of Frédéric Pham (Nice, 2002). Ann. Inst. Fourier (Grenoble) 53 (2003), no. 4, 1055–1116.
- [4] A. Douai and C. Sabbah, *Gauss–Manin systems, Brieskorn lattices and Frobenius structures* (II), Frobenius manifolds (Quantum cohomology and singularities), Hertling, C. and Marcolli, M. eds, Aspects of Mathematics, vol. E36, Vieweg, 2004, p. 1–18.
- [5] B. Dubrovin and Y. Zhang, *Extended Affine Weyl Groups and Frobenius Manifolds*, Compositio Math. 111 (1998) 167–219.
- [6] B. Dubrovin, *Geometry of 2d topological field theories*, Integrable systems and quantum groups (Montecatini Terme, 1993), Lecture Notes in Math., vol. 1620, Springer, Berlin, 1996, pp. 120–348.
- [7] C. Hertling *Frobenius manifolds and moduli spaces for singularities*, Cambridge Tracts in Mathematics, Cambridge University Press, Spring 2002.
- [8] C. Hertling and C. Sevenheck *Limits of families of Brieskorn lattices and compactified classifying spaces*, Adv. Math. 223 (2010), no. 4, 11551224.

- [9] Y. Ishibashi, Y. Shiraishi, A. Takahashi, *A Uniqueness Theorem for Frobenius Manifolds and Gromov–Witten Theory for Orbifold Projective Lines*, arXiv:1209.4870.
- [10] Y. Ishibashi, Y. Shiraishi, A. Takahashi, in preparation.
- [11] Todor E. Milanov, Hsian-Hua Tseng, *The spaces of Laurent polynomials, $\mathbb{P}^1$ -orbifolds, and integrable hierarchies*, Journal für die reine und angewandte Mathematik (Crelle’s Journal), Volume 2008, Issue 622, Pages189–235.
- [12] Y. Manin, *Frobenius manifolds, Quantum Cohomology, and Moduli Spaces*, American Mathematical Soc., 1999 - 303.
- [13] P. Rossi, *Gromov-Witten theory of orbicurves, the space of tri-polynomials and Symplectic Field Theory of Seifert fibrations*, arXiv:0808.2626.
- [14] C. Sabbah, *Frobenius manifolds: isomonodromic deformations and infinitesimal period mappings*, Exposition. Math. 16 (1998), no. 1, 1–57.
- [15] K. Saito, *Primitive forms for a universal unfolding of a function with an isolated critical point*. J. Fac. Sci. Univ. Tokyo Sect. IA Math. **28** (1982), no. 3, 775–792.
- [16] K. Saito, *Period mapping associated to a primitive form*, Publ. RIMS, Kyoto Univ. **19** (1983) 1231–1264.
- [17] K. Saito, *The higher residue pairings $K_F^{(k)}$ for a family of hypersurface singular points*, Proceedings of Symposia in Pure Mathematics Vol. 40 (1983), part 2, pp. 441–463.
- [18] M. Saito, *On the structure of Brieskorn lattice*, Annales de l’institut Fourier (1989) Volume: 39, Issue: 1, Publisher: Institut Fourier, page 27–72.
- [19] K. Saito and A. Takahashi, *From Primitive Forms to Frobenius manifolds*, Proceedings of Symposia in Pure Mathematics, **78** (2008) 31–48.
- [20] I. Satake and A. Takahashi, *Gromov–Witten invariants for mirror orbifolds of simple elliptic singularities*, arXiv:1103.0951.
- [21] A. Takahashi, *Weighted projective lines associated to regular systems of weights of dual type*, Adv. Stud. Pure Math. **59** (2010), 371–388.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA OSAKA, 560-0043, JAPAN

E-mail address: y-ishibashi@cr.math.sci.osaka-u.ac.jp

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA OSAKA, 560-0043, JAPAN

E-mail address: sm5021sy@ecs.cmc.osaka-u.ac.jp

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA OSAKA, 560-0043, JAPAN

E-mail address: takahashi@math.sci.osaka-u.ac.jp